



Comparative Efficacy of Electrical and Solar Light Trap with Pheromone Septa and without Pheromone Septa for *Helicoverpa armigera*

Pravin¹, Amit Kumar Sharma^{1*}, Shraddha Tare¹ and Sandeep Patidar²

¹Department of Entomology, College of Agriculture, Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur (Madhya Pradesh), India.

²Department of Entomology, Rajasthan College of Agriculture, Maharana Pratap University of Agriculture and Technology, Udaipur (Rajasthan), India.

(Corresponding author: Amit Kumar Sharma*)

(Received: 03 January 2025; Revised: 06 February 2025; Accepted: 24 February 2025; Published online: 20 March 2025)

(Published by Research Trend)

DOI: <https://doi.org/10.65041/BiologicalForum.2025.17.3.17>

ABSTRACT: The present investigation was carried out at the BSP (Breeder Seed Production) Farm, Adhartal, JNKVV, Jabalpur (MP) during *rabi* (15th November 2023 to 15th April 2024) season 2023-2024. Four light traps design were used in study via. T4 - Electrical light trap with Acrylic transparent sheet baffle plates + pheromone septa, T2- Electrical light trap with Acrylic transparent sheet baffle plates without pheromone septa, T5 - Solar light trap with Acrylic transparent sheet baffle plates + pheromone septa and T1 - Solar light trap with Acrylic transparent sheet baffle plates without pheromone septa. These four light traps were operated every evening and collection was collected every morning for the duration of the investigation for *Helicoverpa armigera* species, the data was subjected to analysis in paired and two sample t-test. Results indicated that pheromone septa with both electrical and solar light traps, featuring acrylic transparent sheet baffle plates, substantially enhances their efficacy in attracting *Helicoverpa armigera*, making this combined approach a more effective method for trapping this species.

Keywords: *Helicoverpa armigera*, Light trap efficacy, Solar traps, Electrical traps, Pheromone septa, Integrated pest management, Acrylic baffle plates.

INTRODUCTION

Utilizing a light trap as a monitoring tool, one may ascertain the economic standing of the area, the kind and quantity of pests, and their natural enemies in order to get a head start on reducing the density of pest populations. The quantity of pests captured establishes the economic threshold in Integrated Pest Management (IPM), and the pests captured on the light trap may be utilized in the early warning system (Baehaki, 2018). Light trap data is needed to be used as data on dispersal, distribution of pests, and anticipation of pest outbreaks either within or beyond countries (Baehaki *et al.*, 2016). Comparative efficiency of different trapping methods for *Zeuzera pyrina* Fab. (Leopard moth) in walnut orchard. The result found indicated that the combined treatment of pheromone trap and light trap yielded the highest number of males captured on all sampling dates. On the other hand, treatment with the light trap alone showed the lowest number of captured males (Saeidi, 2021).

It provided information respective to insect distribution, abundance, flight pattern and exact time for insect management (Singh and Bambawale 2012; Abbas *et al.*, 2019; Dubey, 2021). Many researchers *viz.*, Singh *et al.* (2018); Singh and Sharma (2018); Kurmi *et al.* (2019); Mishra *et al.* (2019); Meena *et al.*

(2018); Sharma *et al.* (2023); Sharma *et al.* (2020); Bhargava *et al.* (2019); Sharma *et al.* (2019); Kakade *et al.* (2018); Sharma *et al.* (2017); Sharma *et al.* (2015); Patil *et al.* (2024) also conducted studies on the various aspects of light traps uses. Farmers should understand that by attracting and eliminating a single adult moth, they have the potential to effectively manage around 300 to 400 offspring of insects (Abbas *et al.*, 2019). Pheromone septa have been useful in timing, calibration and eventually reducing the application of insecticides. Pheromone traps are very effective monitoring devices and are relatively cheap (Prajapati, 2010). Use of pheromone septa with light trap may increase the trap catches of certain insect species.

MATERIALS AND METHODS

The research was conducted at the BSP farm in Adhartal, JNKVV, Jabalpur (MP) from 15th November 2023 to 15th April 2024. Four separate light traps were used for the study and placed in the Breeder seed production (BSP) Unit Adhartal, JNKVV Jabalpur (MP). The traps were installed at the center of the cropped field on a board bund near the electrical pole. The trap was operated by switching on the power to illuminate the 15 W Ultra violet, every day from sun set to sunrise. Insects trapped in the collection chamber were collected by removing the collection tray at the

end of each quarter of night. The distance between each trap is 100 m approximately (Patidar *et al.*, 2019). Four traps were installed in different direction and placed in such way to avoid light illumination along them (Ambulkar, 2022; Patil, 2023). To kill the trapped insects in the collecting chamber, Formalin 70% (as a fumigating agent) was placed in the collection tray (Singh *et al.*, 2022). The rubber septum containing the lure was changed after every 15 to 17 days (Pal *et al.*, 2014).

Four treatments were compared to test the relative efficacy of electrical and solar light trap with pheromone septa and without pheromone septa in light traps. on the basis of major phototactic insect pest species to be observed. The observation was recorded in two traps with the same funnel diameter and baffle plates. For analysis purpose, the trap catches were adjusted to weekly total of 7 day computed in experiment (Ambulkar, 2022; Patil, 2023). For comparison of efficiency of four light trap designs the observed data were analyzed by paired and two sample t- test for testing the significant different between two treatments as per the requirement (Patil, 2023).

T4 - Electrical light trap with Acrylic transparent sheet baffle plates + pheromone septa.

T2- Electrical light trap with Acrylic transparent sheet baffle plates without pheromone septa.

T5 - Solar light trap with Acrylic transparent sheet baffle plates + pheromone septa

T1 - Solar light trap with Acrylic transparent sheet baffle plates without pheromone septa

RESULTS AND DISCUSSION

Comparing the efficiency of electrical and solar light traps with and without pheromone septa in trapping *Helicoverpa armigera*, the observed data were analyzed by paired and two sample t test for testing the significant difference between two treatments. Results are presented below-

Treatment plates

T4 - Electrical light trap with Acrylic transparent sheet baffle plates +pheromone septa.

T2- Electrical light trap with Acrylic transparent sheet baffle plates without pheromone septa.

T5 - Solar light trap with Acrylic transparent sheet baffle plates + pheromone septa

T1 - Solar light trap with Acrylic transparent sheet baffle plates without pheromone septa

Higher response of *H. armigera* (54.66%) was observed in electrical light trap with acrylic transparent sheet baffle plates + pheromone septa as compared to electrical light trap with acrylic transparent sheet baffle plates without pheromone septa.

Similarly, Higher response of *H. armigera* (39.86%) was recorded in solar light trap with acrylic transparent sheet baffle plates+ pheromone septa as compared to solar light trap with acrylic transparent sheet baffle plates without pheromone septa.

Table 1: Comparative efficacy of electrical light trap with pheromone septa and without pheromone septa for *Helicoverpa armigera*.

Sr. No.	Name of Insects	T4 (Electrical with AT+ pheromone septa) Weekly mean per trap	T2 (Electrical with AT without pheromone septa) Weekly mean per trap	Statistically difference	Increase in trapping efficiency over T2(%)
1.	<i>Helicoverpa armigera</i>	16.58 (4.13)	10.72 (3.35)	S*	54.66

Table 2: Comparative efficacy of Solar light trap with pheromone septa and without pheromone septa for *Helicoverpa armigera*.

Sr. No.	Name of Insects	T5 (Solar with AT+ pheromone septa) Weekly mean per trap	T1 (Solar with AT without pheromone septa) Weekly mean per trap	Statistically difference	Increase in trapping efficiency over T1(%)
1.	<i>Helicoverpa armigera</i>	15.05 (3.94)	10.76 (3.36)	S*	39.86

Figures in parentheses are square root transform value. *Analysis by two sample t-test.




Fig. 1. Comparative efficacy of electrical light traps with pheromone septa and without pheromone septa for *Helicoverpa armigera*.




Fig. 2. Comparative efficacy of solar light traps with pheromone septa and without pheromone septa for *Helicoverpa armigera*.

Comparative efficacy electrical light trap with pheromone septa and without pheromone septa for *H. armigera*. Statistical analysis reveals that the combination of electrical light trap with acrylic transparent sheet baffle plates + pheromone septa was more effective in capturing *H. armigera* than electrical light trap with acrylic transparent sheet baffle plates without pheromone septa alone, indicating that the addition of pheromone septa enhances the trapping efficacy for this species.

Comparative efficacy solar light trap with pheromone septa and without pheromone septa. Similarly, in case solar light trap with acrylic transparent sheet baffle plates + pheromone septa outperformed solar light trap with acrylic transparent sheet baffle plates without pheromone septa alone in capturing *H. armigera*, indicating that the addition of pheromone septa significantly enhances the trapping efficacy of solar light traps for this species.

Similar kind of finding was expressed by Saeidi (2021), who reported that combining of light trap with pheromone trap significantly increase the efficiency of *Z. pyrina* mass trapping compare to light trap without pheromone trap in the walnut orchards.

CONCLUSIONS

The integration of pheromone septa with both electrical and solar light traps, featuring acrylic transparent sheet baffle plates, substantially enhances their efficacy in attracting *Helicoverpa armigera*, making this combined approach a more effective method for trapping this species.

Acknowledgements. The authors extend their sincere thanks to the Head, Department of Entomology and Dean, College of Agriculture, Jabalpur, Madhya Pradesh for providing the necessary facilities and encouragement for conducting the research work.

Conflict of Interest. None.

REFERENCES

Abbas, M., Ramzan, M., Hussain, N., Ghaffar, A., Hussain, K., Abbas, S. and Raza, A. (2019). Role of light traps in attracting, killing and biodiversity studies of insect pests in Thal. *Pakistan Journal of Agricultural Research*, 32(4), 684-690.

Ambulkar, P. L. (2022). Studies on phototactic insect fauna of *rabi* vegetable crops collected in light trap at

Chhindwara (M.P.). Ph.D Thesis, JNKVV, Jabalpur, 1-334p.

- Baehaki, H. E. (2018). Fundamental strategy of controlling brown plant hopper in securing national rice production. *Pengembangan Inovasi pertanian*, 4(1), 63-57.
- Baehaki, S. E., Iswanto, E. H., Munawar, D. and Sumaryono, N. (2016). Light traps abilities of mercury (ML-160 Watt) BSE models and light traps of solar cell (CFL-20 Watt) to capture of pests in the rice field. *American Journal of Engineering Research*, 5, 138-144.
- Bhargava Megha, Sharma, A. K., Shukla, A. and Mishra Yogendra Kumar (2019). Taxonomic documentation of total insect fauna of medicinal plants collected through light trap in Jabalpur district. *Journal of Entomology and Zoology Studies*, 7(6), 642-647.
- Dubey, P. (2021). Use of Light Trap for Taxonomic Documentation and Seasonal Study on Phototactic insect fauna of medicinal plants at Jabalpur. M. Sc. Thesis, JNKVV, Jabalpur, 1-94p.
- Kakade, S. H., Sharma, A. K., Shukla, A., Pachori, R. and Singh, S. (2018). Studies on insect fauna collected in light trap during *Rabi* season in vegetable fields at Jabalpur.
- Kurmi, A., Pachori, R., Bhowmick, A. K., Sharma, A. K., Thomas, M. and Sharma, H. L. (2019). Bio-diversity of photo-tactic hemipteran insects in the rice ecosystems Jabalpur district, Madhya Pradesh, India. *Journal of Entomology and Zoology Studies*, 7(3), 1359-1362.
- Meena, S. K., Sharma, A. K. and Aarwe, R. (2018). Seasonal incidence and population dynamics of major insect pest species of paddy collected in light trap in relation to weather parameters. *International Journal of Current Microbiology and Applied Sciences*, 7(8), 1705-1715.
- Mishra, Y., Sharma, A. K., Bhowmick, A. K., Saxena, A. K. and Kurmi, A. (2019). Seasonal Incidence of Insect Pest Species of Paddy Collected through Light Trap. *International Journal of Current Microbiology and Applied Sciences*, 8(4), 381-393.
- Pal, S., Chatterjee, H. and Senapati, S. K. (2014). Monitoring of *Helicoverpa armigera* using pheromone traps and relationship of moth activity with larval infestation on carnation (*Dianthus caryophyllus*) in Darjeeling Hills. *Journal of Entomological Research*, 38(1), 23-26.
- Patidar, S., Vaishampayan, S. and Band, S. S. (2019). Comparative efficiency of 125-watt Mercury lamp and 15-watt UV (Black light) tube against the major insect-pest in paddy ecosystem. *Journal of Entomology and Zoology*, 7(5), 1163-1167.

- Patil, R., Sharma, A. K., Das, S. B., Kumar, N., Choudhary, P. and Patel, D. K. (2024). Comparative Efficacy of different Funnel Diameters in Light Traps against Major Phototactic Insect Pests of *rabi* Season. *Uttar Pradesh Journal of Zoology*, 45(6), 191-196.
- Patil, R. (2023). Comparative efficacy of the different funnel diameter and different power sources used in light trap against major phototactic insect pests of *rabi* season at Jabalpur. M.Sc. Thesis, JNKVV, Jabalpur, 1-130p.
- Prajapati, M. (2010). Studies on the major insects trapped in different traps associated with same *Rabi* vegetable crops at Jabalpur. M.Sc. Thesis, JNKVV, Jabalpur, 1-163p.
- Saeidi, Z. (2021). Integration of pheromone and light traps for mass trapping of leopard moth, *Zeuzera Pyrina* (Lepidoptera: Cossidae) in walnut orchards. *Journal of Entomological Society*, 41(2), 123-133.
- Sharma, A. K., Mandloi, R. and Pachori, R. (2017). Study on Biodiversity of Phototactic Fauna Collected in Light Trap in Chickpea (*Cicer aritenum* Linn.). *International Journal of Agriculture Sciences*, 9(12), 4037-4041.
- Sharma, A. K., Mandloi, R., Bhowmick, A. K. and Thakur, A. S. (2019). Study on biodiversity of phototactic hexapod fauna by light trap in soybean (*Glycine max* L.) ecosystem. *Journal of Entomology and Zoology Studies*, 7(2), 641-646.
- Sharma, A. K., Mandloi, R., Saxena, A. K., Thakur, A. S., Sharma, R. and Ramakrishnan, R. S. (2020). Biodiversity of Phototactic insect pests of chickpea ecosystem and records on population dynamics of *Helicoverpa armigera* (Hubner) and *Agrotis ipsilon* (Hufnagel). *Journal of Pharmacognosy and Phytochemistry*, 9(1), 824-829.
- Sharma, A. K., Pachori, R., Thakur, A. S. and Mandloi, R. (2015). Population Dynamics of Major Phototropic Insect Pests of Medicinal Crops with Special Reference to Effect of Ecological Factors on *Helicoverpa armigera* Hub. and *Agrotis ipsilon* Huf. The *Ecoscan* 8 (Special issue), 921-927.
- Sharma, A. K., Yadav, K. N., Tare, S., Nayak, S. and Seervi, S. (2023). Phototactic Response and Taxonomic Distribution of Predaceous Species of Paddy Ecosystem. *Biological Forum – An International Journal*, 15(3), 91-94.
- Singh, R. P., Böttger, D. and Brehm, G. (2022). Moth light traps perform better with vanes: A comparison of different designs. *Journal of Applied Entomology*, 146, 1343-1352.
- Singh, S. and Sharma, A. K. (2018). Population dynamics of major insect pests of rice. *Indian Journal of Entomology* 80(4): 1700-1702.
- Singh, S., Sharma, A. K., Saxena, A. K., Panday, A. K. and Kakade, S. (2018). Taxonomic analysis of phototactic beneficial insects as biocontrol agents (Predators and parasites) collected in light trap in rice ecosystem at Jabalpur. *Journal of Entomology and Zoology Studies*, 6(3), 850-853.
- Singh, S. K. and Bambawale, O. (2012). Light trap for managing insects. Indian Council of Agriculture Research, Unit National Center for Integrated Pest Management. <http://www.google.com/patent/WO2012098484A1?cl=en>

How to cite this article: Pravin, Amit Kumar Sharma, Shraddha Tare and Sandeep Patidar (2025). Comparative Efficacy of Electrical and Solar Light Trap with Pheromone Septa and without Pheromone Septa for *Helicoverpa armigera*. *Biological Forum*, 17(3): 104-107.