

Determination of Nutritive Contents (Protein and Carbohydrate) of Popular Mulberry Varieties

Aabid Khaliq Tantray¹, Mohd Shahid Raja¹, Naveena Nazim^{1*}, Aabid Ahmad Bhat¹, Mohd Farooq Baqal² and Khursheed Ahmad Sahaf¹

¹Division of Silk Product Science, College of Temperate Sericulture, SKUAST- K, Mirgund.

²Division of Host Plant Production, College of Temperate Sericulture, SKUAST- K, Mirgund.
Post Box No. 674, GPO-Srinagar, Kashmir, India.

(Corresponding author: Naveena Nazim*)

(Received 24 September 2021, Accepted 27 November, 2021)

(Published by Research Trend, Website: www.researchtrend.net)

ABSTRACT: Proteins and carbohydrates are the main nutritive components in mulberry leaves. The present study was undertaken to analyse the protein and carbohydrate content in the leaves of popular mulberry varieties (Goshoerami, Koku-21 and Ichinose) at three different stages (tender, medium and coarse). In general, decreasing trend in terms of total protein and carbohydrate content was recorded from tender to coarse textured leaves of all three varieties. The leaf extract of Goshoerami (tender) and Koku-21 (tender) recorded the highest protein and carbohydrate content to the tune of 28.93 and 18.68% respectively. Among the varieties, Goshoerami recorded highest protein content of 25.24%, whereas highest carbohydrate content (16.91%) was found in Koku-21 mulberry variety.

Keywords: Silkworm, Mulberry leaves, Protein and Carbohydrate.

INTRODUCTION

Mulberry plants are fast growing and found in varied climatic conditions. It is mainly used to feed the silkworms for the production of silk. Presence of good amount of bioactive constituents makes it choice of plant for various pharmaceutical and nutraceutical studies. Mulberry plant is also appreciated for the source of mulberry fruit, which is good in taste and have various health benefits. Fruits can be consumed fresh, fried or in the form of juices, squashes etc. Silkworm larval growth, development and cocoon production is directly proportional to the nutritional and biochemical quality of mulberry leaves (Krishnaswami, 1978). Any deficiency in the quality of leaf causes changes in the metabolic activity of silkworm larvae (Ito, 1972). Protein and carbohydrate are prime components of nutrition in leaf and their quantities greatly influence the growth of silkworm. Mulberry leaf quality also depends on the age, variety and leaf position within same species (Narayanswamy *et al.*, 1996). Present study was undertaken for quantitative analysis of protein and carbohydrate composition of mulberry leaves (tender, medium and coarse) of three mulberry varieties (Goshoerami, Ichinose and Koku-21) grown under temperate climatic conditions of Kashmir.

MATERIALS AND METHODS

Present study was conducted at College of Temperate Sericulture, SKUAST-K, Mirgund, Jammu and Kashmir. Leaf samples comprising of equal proportion of tender, medium and coarse from three mulberry varieties *viz.*, Goshoerami, Ichinose and Koku-21 were taken in early morning during the spring 2021.

Processing of leaf samples. The collected leaf samples were first washed with running tap water, followed by distilled water, to decontaminate leaves from dust and other impurities. The samples were air dried on filter papers and then oven dried at 60-65°C (Chapman, 1964) till constant weight was obtained. After drying, samples were crushed with the help of mortar pestle, followed by sieving and stored in envelopes for further use.

Experimental Material. Ethanol (analytical grade), Folin- Ciocalteu reagent (FCR), Anthrone reagent, Sulphuric acid (H₂SO₄), Trichloroacetic acid (TCA), Standard glucose, Alkaline copper reagent (Sodium potassium tartrate (KNaC₄H₄O₆.4H₂O), Copper sulphate (CuSO₄), Sodium hydroxide (NaOH), Sodium carbonate (Na₂CO₃), Bovine serum albumin (BSA), Hydrochloric acid (HCL), Distilled water and Mulberry leaf (tender, medium & coarse) of three popular varieties *viz.*, Goshoerami, Ichinose and Koku-21.

Estimation of total protein content. Total protein content was measured by colorimetric method described by Lowry *et al.*, 1951. 50mg of dry leaf from each sample was homogenized in 80% ethanol using mortar pestle. Homogenates were centrifuged at 5000rpm for 20 minutes. Supernatants were discarded and residues suspended in 10ml of 10% TCA (Trichloroacetic acid) for 30 minutes. The mixtures were centrifuged at 5000rpm for 10 min and the supernatants discarded. Pellets were washed with 5% TCA and protein precipitates (pellets) were dissolved in 1N NaOH and kept in hot water bath for 30 minutes. The samples were diluted 10 times with distilled water. 1ml was taken as protein sample followed by the addition of 5ml of alkaline copper reagent and allowed to stand for 10 minutes. 0.5ml of 50% FCR reagent was added rapidly and mixed. FCR added mixtures are kept at room temperature for 30 minutes and optical density was taken at 750nm.

Estimation of total carbohydrate content. Carbohydrate content in leaf samples was estimated by Thimmaiah 1999, using the anthrone reagent with slight modification. 0.1g of dry leaf samples were taken in test tubes and added with 5ml of 0.5N HCL. Tubes were kept for 3 hours in hot boiling water bath and cooled at room temperature. Dry sodium carbonate was added to neutralize samples till effervescence ended. Volume of tubes were made 10ml with distilled water and 0.05ml from each sample tubes was taken for analysis. Volume of sample tubes was made up to 1ml with distilled water followed by addition of 4ml anthrone reagent. Tubes were kept in boiling water for 8 minutes, cooled rapidly and absorbance of green to dark green colour was read at 630nm.

RESULTS AND DISCUSSION

Total Protein. The results with respect to total protein content of mulberry leaves are presented Table 1 and Fig. 1. In general, tender leaves of Goshorami recorded significantly highest (28.93%) total protein content followed by tender leaves of Koks-21(27.58%) and Ichinose (23.96%). Obtained results coincides with results put forth by Lokesh *et al.*, 2012; Murthy *et al.*, 2013; Jyothi *et al.*, 2016) who reported that the tender leaves of mulberry contain more protein content than medium and coarse textured leaves. High protein content in tender leaves makes it best feed during early stage of larvae where larvae get more nutrition by consuming even less leaf quantity. Protein and carbohydrate content are the major nutritional component in leaf and directly influences the larval growth. Krishnaswami (1978) reported that the constituents like water, total sugars, soluble sugars, proteins, crude fiber and minerals are efficiently consumed by silkworm larvae.

Total carbohydrate. Results of carbohydrate content in mulberry leaves are presented in Table 2 and Fig. 2. Highest carbohydrate content was observed in Koks-21 tender (18.68), followed by Ichinose tender (16.98%) and Goshorami tender (14.48%) which declined gradually within increasing growth periods. More photosynthetic products to upper leaves can be related to the higher carbohydrate composition in the tender (upper leaves) than lower leaves and its depletion can be attributed to genotypic character of matured leaves. Current results are in conformity with the results obtained by (Lokesh *et al.*, 2012; Murthy *et al.*, 2013) who reported that the tender leaves of mulberry contain more carbohydrate content than medium and coarse textured leaves.

Table 1: Protein content (%) in leaves of different mulberry varieties.

Leaf Stage	Mulberry varieties			
	Goshorami	Koks-21	Ichinose	Mean
Tender	28.93	27.58	23.96	26.82
Medium	25.49	24.58	23.68	24.58
Coarse	21.29	20.36	18.87	20.18
Mean	25.24	24.17	22.17	

C.D (p 0.05)

Varieties = 0.0143; Leaf type = 0.0143; Varieties × leaf type = 0.025

Table 2: Carbohydrate content (%) in leaves of different mulberry varieties.

Leaf Stage	Mulberry varieties			
	Goshorami	Koks-21	Ichinose	Mean
Tender	14.48	18.68	16.98	16.713
Medium	13.93	16.25	16.66	15.613
Coarse	12.66	15.8	14.83	14.430
Mean	13.69	16.91	16.16	

C.D (p 0.05)

Varieties = 0.1043; Leaf type = 0.1043; Varieties × leaf type = 0.181




Fig. 1. Graphical representation of total protein content in different leaf varieties of mulberry.




Fig. 2. Graphical representation of carbohydrate content in different leaf varieties of mulberry.

CONCLUSION

Goshoerami is a popular cultivar of temperate regions like Kashmir. Highest protein content in the tender stage of the leaves surmises about the possibility of its utilization during chawki silkworm rearing. In addition, significant amount of protein and carbohydrate content in the Koku-21 variety suggests its potential for being utilized in chawki rearing of silkworms.

FUTURE SCOPE

Mulberry leaves have since long being utilized as a potential source for nutraceutical and biomedical applications. The present investigation of analysis of protein and carbohydrates in the leaves of the selected mulberry varieties is very important for providing reliable data for the use in many areas of nutrition, food science and health. The study shall benefit the researchers for development of mulberry leaf based nutraceutical products in addition to generate a standardized approach of the analytical method for accurate and reliable determination of the protein and carbohydrate contents.

Acknowledgment. The authors are highly thankful to BIRAC, New Delhi for funding the requisite chemicals.

Conflict of Interest. None.

REFERENCES

- Chapman, H. D. (1964). Suggested foliar sampling and handling techniques for determining the nutritional status of some field, horticultural and plantation crops. *Indian Journal of Horticulture*; 21: 97-119.
- Ito, T. (1972). An approach to nutritional control mechanisms in silkworm *Bombyx mori* L. *Israel Journal of Entomology*, 7: 1-6.
- Jyothi, M., Pratap, M., Seenaiiah, R., and Thimma, N. S. (2016). Biochemical Analysis and Protein Profile through SDS-page in Leaves of four (RC-1, RC-2, G-2 and G-4) Mulberry (*Morus alba* L.) Cultivars. *International Journal of Recent Scientific Research*, 7(4): 10296-10300.
- Krishnaswami, S. (1978). New Technology of silkworm rearing, Bull No-2 CSRTI, Mysore, India: 1-20.
- Lokesh, T. S., Ananthanarayana, S. R. and Shivashankar, M. (2012). Influence of vermicompost on improved mulberry varieties of nutrient status of leaf maturity

- levels. *Asian Journal of Biological Sciences*, 7(1): 9-12.
- Lowry, O. H., Rosebrough, N. F., Farr, A. L., and Randall, R. J. (1951). Protein measurement with folin phenol reagent. *Journal of Biological Chemistry*, 193: 267-275.
- Murthy, Y. V. N., Ramesh, H. L., and Munirajappa (2013). Evaluation of mulberry (*Morus*) variety for leaf yielding parameters and phytochemical analysis under different spacing systems. *Indian Journal of Applied Research*, 3(8): 31-33.
- Narayanswamy, K. C., Geethabai M. and Raghuraman, R. (1996). Mite pests of mulberry- A Review. *Indian Journal of Sericulture*, 35(1): 1-8.

How to cite this article: Aabid Khaliq Tantray, Mohd Shahid Raja, Naveena Nazim, Aabid Ahmad Bhat, Mohd Farooq Baqal and Khursheed Ahmad Sahaf (2021). Determination of Nutritive Contents (Protein and Carbohydrate) of Popular Mulberry Varieties. *Biological Forum – An International Journal*, 13(4): 1303-1306.