

## **Trichoderma as Potential Biocontrol Agent on Diseases of Soybean (*Glycine max* L.): A Comprehensive Review**

**Manjula, Aishwarya, Ajay Kumar Gautam\* and Anupam Kumar**

Department of Plant Pathology, School of Agriculture,  
Abhilashi University Mandi (Himachal Pradesh), India.

(Corresponding author: Ajay Kumar Gautam\*)

(Received: 16 July 2024; Revised: 19 August 2024; Accepted: 15 September 2024; Published: 15 October 2024)

(Published by Research Trend)

**ABSTRACT:** *Trichoderma* primarily treats soil-borne infections and a few leaf and panicle diseases in various plant species. *Trichoderma* not only prevents disease, but it also increases plant resistance, accelerates growth, improves nutrient uptake efficiency, and cleans the environment of agrochemical pollutants. *Trichoderma* spp. is also a safe, cost-effective, efficient, and environmentally friendly biocontrol agent for a wide range of crop species. This study examines the use of *Trichoderma* and its control effects in the management of various plant diseases. *Trichoderma*'s biological control mechanisms in plant fungal and nematode disease, such as competition, antibiosis, antagonism, and mycoparasitism, have been discussed. We also talked about how *Trichoderma* promotes plant growth while inducing plant systemic resistance. Developing a diverse range of *Trichoderma* application technologies is an important avenue for future research and development, given its potential to contribute to the agricultural sector's long-term growth.

**Keywords:** *Trichoderma*, plant diseases, biological control, growth promotion, action mechanism.

### **INTRODUCTION**

Soybean (*Glycine max* L.) is a major crop in India and an important source of vegetable protein and oil (Herridge *et al.*, 2008; Prevost *et al.*, 2010). Soybean is a leguminous oil seed crop grown around the world due to its high protein (40%) and fat (20%) content. Soybean is thought to be a subtropical plant from Southeast Asia. This crop was brought to Europe and the United States in the 1700s and 1800s. Farmers in the Midwest of the United States grow roughly half of all soybeans. East Asia accounts for roughly 45% of soybean production, with the remaining 55% coming from America. Brazil, Argentina, Paraguay, China, and India are the world's leading soybean producers, in addition to the United States (Joy *et al.*, 1998).

In India, it is grown as a rainfed kharif crop on 10.97 million ha, yielding 10.99 million tonnes with a productivity of 1002 kg/ha in 2016-17 (SOPA). Madhya Pradesh is India's leading soybean producer, with 54.01 lakh ha of soybean production, an average productivity of 1020 kg/ha, and a total production of 55.06 lakh tonnes in 2016-17 (SOPA). Persoon (1794) established *Trichoderma* as a genus in Germany, proposing four species: *Trichoderma viride*, *T. nigroscens*, *T. aureum*, and *T. roseum*. *Trichoderma* was first isolated from Madras, India, by Thakur and Norris in 1928.

*Trichoderma* can be found in all temperate and tropical soils, as well as forest, agricultural, prairie, salt marsh, and desert soils. *Trichoderma*, for example, accounted

for up to 3% of total fungal propagules in a wider range of forest soils and 1.5% in pasture soils in a variety of crops (Domsch *et al.*, 1980). *Trichoderma* spp. have been known to parasitize other fungi for approximately 70 years (Table 1).

*Trichoderma* has been identified as saprophytic fungi found in the roots of many plants. It has been reported that antagonistic fungi produce a variety of volatile and non-volatile organic compounds (Siddiquee *et al.*, 2012; Meena *et al.*, 2017), as well as diffusible antibiotics such as *trichodermin*, *gliotoxin*, and *virid* (Mukherjee *et al.*, 2012; Vargas *et al.*, 2014; Sharma *et al.*, 2016), to control plant pathogenic fungi. During antagonistic activity, these fungi primarily compete with pathogenic fungi for nutrients and space (John *et al.*, 2010; Carvalho *et al.*, 2015).

Similarly, fungi such as *Trichoderma* colonize plant roots, allowing them to protect against biotic stresses such as pathogenic infection (John *et al.*, 2010; Carvalho *et al.*, 2015; Jogaiah *et al.*, 2018) while also promoting plant growth. Biological control is primarily used to control harmful organisms in plants by utilizing beneficial organisms and their products to control plant diseases and effectively reduce the use of chemical fertilizers and pesticides. *Trichoderma*, a biological fungus commonly used for plant pest control, lives in the soil, air, plant surface, and other ecological environments and can effectively control a wide range of plant diseases (Haouhach *et al.*, 2020; Zheng *et al.*, 2021; Wang *et al.*, 2022). *Trichoderma* is primarily used to control soil-borne diseases in plants, as well as

certain leaf and spike diseases (Vicente *et al.*, 2020; Abbas *et al.*, 2022). *Trichoderma* has been shown to prevent disease, promote plant growth, improve nutrient utilization, improve plant resistance, and repair agrochemical pollution (Tilocca *et al.*, 2020; Fontana *et al.*, 2021; Sanchez-Montesinos *et al.*, 2021; Al-Surhane, 2022; Tyskiewicz *et al.*, 2022).

*Trichoderma* is a biocontrol fungus that is widely distributed throughout the world. Tyskiewicz *et al.* (2022) found that *Trichoderma* has significant potential for biological disease control in plants. *Trichoderma* use to control plant diseases has been studied all over the world. *T. viride* and *T. harzianum* inhibit 29 species of plant pathogenic fungi from 18 genera, including *Botrytis*, *Fusarium*, and *Rhizoctonia*, in varying degrees. *Trichoderma* controls a wide range of plant pathogenic fungi, including *Rhizoctonia solani*, *Pythium ultimum*, *Fusarium oxysporum*, *Sclerotinia sclerotiorum*, *Botrytis cinerea*, *Pseudocercospora* spp., and *Colletotrichum* spp. (Alvarez-García *et al.*, 2020; Andrade-Hoyos *et al.*, 2020; Carro-Huerta *et al.*, 2020; Damodaran *et al.*, 2020; Zhang *et al.*, 2020, 2021; Al-Askar *et al.*, 2021; Chen *et al.*, 2021; Degani and Dor 2021; Dugassa *et al.*, 2021; Intana *et al.*, 2021; Zhang *et al.*, 2022; Zhang *et*

*al.*, 2022). *Trichoderma* has been widely used for biological control of cotton verticillium wilt, crop grey mold, tomato graymold, melon wilt, potato dry rot, tobacco root rot, and other plant diseases. (Andrade-Hoyos *et al.*, 2020; Alfiky and Weisskopf 2021; Lazazzara *et al.*, 2021; Leal *et al.*, 2021; Manganiello *et al.*, 2021; Degani *et al.*, 2021a; Pollard-Flamand *et al.*, 2022; Rees *et al.*, 2022; Risoli *et al.*, 2022).

Aside from biotic stress protection, *Trichoderma* reduces abiotic stress such as drought and salinity (Mastouri *et al.*, 2010; Contreras-Cornejo *et al.*, 2014). *Trichoderma* strains influence bioactive metabolite production (Garnica-Vergara *et al.*, 2016). *Trichoderma* spp. also improves nutrient availability for plants, allowing them to better withstand biotic and abiotic stresses. Several *Trichoderma* species have been used in leguminous plants as biocontrol agents and growth promoters. John *et al.* (2010) investigated *T. viride* ability to significantly reduce soil-borne pathogens while also improving root systems. Chickpea (*Cicer arietinum*) inoculated with *T. harzianum* and *Aspergillus niger* showed significant increases in shoot and root length and weight (Yadav *et al.*, 2011).

**Table 1: Different *Trichoderma* species effective against plant disease causing Pathogens.**

Sr. No.	Crop/Plant	<i>Trichoderma</i> spp.	Pathogens	References
1.	Tomato	<i>T. viride</i> , <i>T. harzianum</i>	<i>Fusarium solani</i> <i>Rhizoctonia solani</i>	Haggag and El-Gamal (2012)
2.	Soybean	<i>T. viride</i>	<i>Fusarium oxysporum</i> f. sp. <i>adzuki</i>	John <i>et al.</i> (2010)
3.	Rice	<i>T. viride</i> , <i>T. koningii</i> , <i>T. harzianum</i>	<i>Rhizoctonia solani</i> <i>Fusarium</i> spp.	Bhat <i>et al.</i> (2009); Gomathinayagam <i>et al.</i> (2010); Chakravarthy <i>et al.</i> (2011); Bhramaramba and Nagamani (2013); Biswas and Datta (2013); Gangwar and Sharma (2013)
4.	Potato	<i>T. virens</i> <i>T. harzianum</i>	<i>Rhizoctonia solani</i> <i>Fusarium sambucinum</i>	Ru and Di (2012) Basu (2009); Selvakumar (2008); Pandey and Pundhir (2013)
5.	Mungbean	<i>T. harzianum</i> <i>T. viride</i> <i>T. virens</i>	<i>Rhizoctonia bataticola</i>	Dubey <i>et al.</i> (2009)
6.	Onion	<i>T. viride</i> <i>T. harzianum</i> <i>T. reesei</i>	<i>Alternaria alternata</i> <i>Alternaria porri</i> <i>Alternaria tenuissima</i>	Mishra and Gupta (2012); Prakasam and Sharma (2012); Yadav <i>et al.</i> (2013); Shahnaz <i>et al.</i> (2013)
7.	Maize	<i>T. harzianum</i>	<i>Penicillium notatum</i> , <i>Rhizoctonia</i> <i>solani</i> <i>Fusarium oxysporum</i> <i>Alternaria alternata</i>	Bhandari and Vishunavat (2013); Pal <i>et al.</i> (2013)
8.	Chilli	<i>T. viride</i> , <i>T. harzianum</i> <i>T. pseudokoningii</i>	<i>S. rolfsii</i> <i>F. oxysporum</i> <i>Pythium</i> spp., <i>R. solani</i>	Rini and Sulochana (2006); Kapoor (2008); Vasanthakumari and Shivanna (2013)

**Biocontrol Mechanisms of *Trichoderma*.** Nowadays, *Trichoderma* spp. are promising biocontrol agents against fungal phytopathogens. Examples of such interactions include *T. harzianum* action on *Fusarium oxysporum*, *F. roseum*, *F. solani*, *Phytophthora Manjula et al.*,

*colocaciae*, and *Sclerotium rolfsii*. In general, biocontrol agents grow naturally on the root surface and thus affect root disease in particular, but they can also be effective against foliar diseases (Leaf rot) and bark diseases (*Citrus gummosis*). They can act indirectly by

competing for nutrients and space, changing environmental conditions, or promoting healthy plant growth, plant defensive mechanisms, and antibiosis, or directly through mechanisms like mycoparasitism. Increase Dry matter production increased significantly. Provide natural, long-term immunity to crops and soil (Papavizas, 1985, Howell, 2003; Vinale *et al.*, 2008).

#### Benefits of *Trichoderma*

**(a) Disease control:** *Trichoderma* is a biocontrol agent that is widely used for both soil-borne and foliar diseases (Harman *et al.*, 2010). It also produces cell wall degrading enzymes against pathogenic fungi from various genera, including *Rhizoctonia Fusarium*, *Phytophthora*, *Scelerotiinia* and *Colletotrichum*. *Trichoderma* species have been shown to control a variety of diseases (Table 2).

**Table 2: Diseases controlled by *Trichoderma*.**

Sr. No.	Crop	Diseases	Pathogen	References
1.	Rice	Sheath blight, Bacterial leaf blight, Bakanae	<i>Rhizoctonia solani</i> <i>Fusarium moniliforme</i>	Biswas and Datta (2013); Ng <i>et al.</i> (2015)
2.	Wheat	Leaf blight Loose smut	<i>Alternaria triticina</i> <i>Ustilago segetum</i>	Parveen and Kumar (2004); Singh (2004)
3.	Chickpea	Wilt complex, Root rot	<i>Fusarium</i> spp., <i>Sclerotium</i> spp., <i>Rhizoctonia solani</i>	Gupta <i>et al.</i> (2005)
4.	Pigeon pea	Wilt	<i>Fusariumudum</i>	Chaudhary and Prajapati (2004)
5.	Apple	Ring rot White root rot	<i>Botryosphaeria berengeriana</i> <i>Dematophora necatrix</i>	Kexiang <i>et al.</i> (2002) Tapwal <i>et al.</i> (2005)
6.	Guava	Die back	<i>Lasiodiplodia theobromae</i>	Yadav and Majumdar (2005)
7.	Chilli	Dry root rot	<i>Rhizoctonia solani</i>	Bunker and Mathur (2001)
8.	Tomato	Fusarium wilt Crown, stem and root rot diseases, Collar rot of tomato	<i>Fusarium oxysporum</i> f. sp. <i>lycopersici</i> <i>Rhizoctonia solani</i> , <i>Sclerotinia</i> spp. and <i>Pythium</i> <i>Sclerotium rolfsii</i>	Komy <i>et al.</i> (2015); Marzano <i>et al.</i> (2013); Amin <i>et al.</i> (2010)
9.	Potato	Damping off, Black Scurf, Charcoal Rot, Bacterial brown rot.	<i>Rhizoctonia solani</i> <i>Fusarium</i> and <i>Phoma</i> spp.	Gogoi <i>et al.</i> (2007)
10.	Beans	web blight of beans	<i>Sclerotinia sclerotiorum</i>	Amin <i>et al.</i> (2010)

**(b) Plant Growth Promoter:** *Trichoderma* strains dissolve phosphates and micronutrients. The application of *Trichoderma* strains to plants increases the number of deep roots, improving the plant's ability to withstand drought.

**(c) Biochemical Elicitors of Disease:** *Trichoderma* strains have been found to cause plant resistance. Three types of compounds produced by *Trichoderma* that cause plant resistance have been identified. In plant cultivars, these compounds cause ethylene production, hypersensitivity, and other defensive reactions.

**(d) Transgenic Plants:** The introduction of *Trichoderma* endo chitinase gene into plants like tobacco and potatoes has increased their resistance to fungal growth. Selected transgenic lines are highly tolerant to foliar pathogens such as *Alternaria alternata*, *A. solani*, and *Botrytis cinerea*, as well as the soil-borne pathogen *Rhizoctonia* spp.

**(e) Bioremediation:** *Trichoderma* strains play an important role in bioremediation of pesticide and herbicide-contaminated soil. They can degrade a diverse range of insecticides, including organochlorines, organophosphates, and carbonates.

#### Application of *Trichoderma* in biological control of plant fungal diseases:

**1. Seed treatment:** Before sowing, mix 6-10 grams of *Trichoderma* powder per kilogram of seed.

**2. Nursery treatment:** Apply 10 to 25 g of *Trichoderma* powder per 100 square meters of nursery bed. The efficacy of neem cake and FYM is increased when applied prior to treatment.

**3. Cutting and seedling root dip:** Mix 10g of *Trichoderma* powder with 100g of well-rotten FYM per liter of water, then dip the cuttings and seedlings for 10 minutes before planting.

**4. Soil treatment:** Apply 5 kg of *Trichoderma* powder per hectare after incorporating sun hemp into the soil for green manuring. Alternatively, mix 1kg of *Trichoderma* formulation in 100kg of farmyard manure and cover with polythene for 7 days.

**5. Sprinkle the heap with water intermittently:** Turn the mixture in every 3-4 days and then broadcast in the field.

**6. Plant Treatment:** Drench the soil near the stem with 10g *Trichoderma* powder mixed in 1 liter of water.

#### Precautions taken during *Trichoderma* Application:

1. Avoid using chemical fungicides for 4-5 days following *Trichoderma* application.

2. Do not use *Trichoderma* on dry soil. Moisture is essential to its growth and survival.

3. Avoid directly exposing treated seeds to sunlight.

4. Avoid storing treated FYM for extended periods of time.

#### *Trichoderma* as a potential biological control agent.

The term 'biocontrol' was coined in 1914 with a focus

on plant pathogens and insects, respectively. Biocontrol is the process of reducing plant pest populations using naturally occurring organisms as part of integrated disease management. A variety of biocontrol agents or bio-fungicides exist in the ecosystem, and they must be isolated before being used because biocontrol agents are low-cost to produce, have a long-lasting effect on pathogen growth, and have no effect on human health. *Trichoderma* was first described as a bio-control agent by Weindling (1932); *Trichoderma* species are free-living, cosmopolitan fungi found in soils, decaying organic and vegetable matter (Harman *et al.*, 2004a). Several *Trichoderma* species show potential for biological control of plant pathogenic fungi. *Trichoderma viride* inhibited the growth of soil-borne *Pythium debaryanum*, *Sclerotium rolfsii*, *Fusarium lini*, *F. Culmorum*, and *Fomusanosus* (Wright 1956). *Trichoderma viride* is commonly used in commercial orchard oils to control *Armillaria mellea* (Bliss, 1951). *Trichoderma viride* has been shown in vitro and in vivo to be an antagonist against *Venturi inaequalis*, the causative agent of apple scab (Lindow, 1985). *Trichoderma viride* was discovered to be an antagonist to *Drechslera sorokiniana*, the causative agent of wheat root rot, seedling, and foliar blights (Prasad *et al.*, 1978). Singh *et al.* (1991) found that using *Trichoderma viride* culture filtrate reduced germination of *N. indica* teliospores and sporidia significantly. Biocontrol agent to barley seeds reduced *Helminthosporium* infection on coleoptiles by 87%.

*Trichoderma viride* culture filtrate inhibited chlamyospore germination and suppressed the mycelial growth of *Ustilagoseg tum tritici*. Field tests also demonstrated the agent's biocontrol potential against wheat loose smut (Aggarwal *et al.*, 1993). *Trichoderma harzianum* has been shown to inhibit the growth of *S. rolfsii*, the causative agent of root rot in many crops. Wells *et al.* (1972) applied this species' inoculum to rows of tomato seedlings to protect the crop from *S. rolfsii*. It has also demonstrated potential for biocontrol of *Macro phominaphaseolina* (Elad *et al.*, 1986).

*Trichoderma harzianum* has also been shown to have biocontrol potential against *F. solani* (Calvet *et al.*, 1990), *Fusarium oxysporumciceris*, *Pythium aphanidermatum* (Bhardwaj and Gupta 1987), *Rhizoctonia solani* (Wilson *et al.*, 1988; Cole and Zrinyka 1988), and *S. rolfsii* (Deve and Dutta 1988). In a study of *Trichoderma harzianum* effect on fungal pathogens infecting wheat and black oat straw, the antagonist reduced the incidence of *Cochliobolus sativus*. This antagonist's culture filtrate inhibited the germination of teliospores and sporidia of *N. indica* (Singh *et al.*, 1991) and chlamyospores of *U. segetum tritici*. *Trichoderma koningii* has shown biocontrol potential against *U. segetum tritici* and *Drechslera sorokiniana*, *T. reesi* against *D. sorokiniana* and *T. pseudokoningii* against karnal bunt.

**Table 3: Biocontrol agents of some fungal Plant diseases by different *Trichoderma* spp.**

Sr. No.	<i>Trichoderma</i> spp.	Fungal plant pathogen	Plant Diseases
1	<i>Trichoderma harzianum</i> Rifai	<i>Sclerotium rolfsii</i>	Southern stem blight of soybean
2	<i>Trichoderma harzianum</i> Rifai	<i>Sclerotium rolfsii</i>	Rotting of common vegetables
3	<i>Trichoderma harzianum</i> Rifai	<i>Sclerotium rolfsii</i>	Collar rot of lentil
4	<i>Trichoderma harzianum</i> Rifai	<i>Sclerotinia sclerotiorum</i>	Sunflower head rot
5	<i>Trichoderma harzianum</i> Rifai	<i>Macrophomina phaseolina</i>	Root rot of blackgram
6	<i>Trichoderma harzianum</i> Rifai	<i>Fusarium oxysporum</i> f. <i>splycopersici</i>	Fusarium wilt of tomato
7	<i>Trichoderma harzianum</i> Rifai	<i>Fusarium oxysporum</i> f. sp <i>gladioli</i>	Fusarium wilt & corm rot of <i>gladioli</i>
8	<i>Trichoderma viride</i> Pers. Ex Fr.	<i>Fusarium udum</i>	Pigeon pea wilt
9	<i>Trichoderma viride</i> Pers. Ex Fr.	<i>Rhizopus oryzae</i>	Cotton seedling disease
10.	<i>Trichoderma virens</i> (Miller, Giddens & Foster) v.ArX	<i>Serpula lacrymans</i>	Wood decay
11.	<i>Trichoderma virens</i> (Miller, Giddens & Foster) v.ArX	<i>Colletotrichum truncatum</i>	Brown blotch disease of cowpea
12.	<i>Trichoderma lignorum</i> (Tode) Harz	<i>Rhizoctonia solani</i>	Damping-off of bean
13.	<i>Trichoderma Koningii</i> Oudem	<i>Sclerotium cepivorum</i>	White rot disease of onion roots

***Trichoderma* as biocontrol agent for different soybean diseases.** Soybean (*Glycine max* L.) is the third-most important oilseed crop. Several researchers developed integrated management schedules for root, seed, and foliar diseases. According to Singh and Thapliyal (1998), seed treatment with Vitavax 200 plus *Trichoderma harzianum* or *Gliocladium virens* effectively manages pre- and post-emergence seedling rot.


Pant and Mukhopadhyay (2001) described how to manage soybean seed and seedling rot caused by *R.*

*solani* using biocontrol agents *Gliocladium virens* and *Trichoderma harzianum*. (Ray *et al.*, 2007) discovered that they improve seed germination and reduce seedling rot in soybeans. Another study found *Trichoderma viride* to be effective against two fungal pathogens that infect soybeans, *Fusarium oxysporum* and *Pythium arrhenomanes* (John *et al.*, 2010). Khodke and Raut (2010) investigated the management of root rot or collar rot through seed treatment and fungicide application to the soil.

*Trichoderma* and plant growth-promoting rhizobacteria, *P. fluorescens*, were tested under glasshouse and field conditions against many soil-borne plant pathogens, including *R. solani*, *S. rolfisii*, and *M. phaseolina*, which cause root and stem rot disease in soybeans (Mishra *et al.*, 2011). *Trichoderma* species inhibited the growth of oilseed-borne fungi such as *Aspergillus flavus*, *Alternaria alternata*, *Curvularia lunata*, *Fusarium moniliforme*, *Fusarium oxysporum*, *Rhizopus nigricans*, *Penicillium notatum*, and *Penicillium chrysogenum*, which harm oil seed crops such as soybean, sesame, and sunflower (Jat and Agalave 2013).

**Antibiosis effect of *Trichoderma*:** The term "antibiosis" refers to *Trichoderma* ability to secrete antagonistic substances that prevent plant pathogenic fungi from growing (Kottb *et al.*, 2015; Izquierdo-Garcia *et al.*, 2020; Moran-Diez *et al.*, 2020; Shobha *et al.*, 2020; El-Hasan *et al.*, 2022). *Trichoderma* produces numerous antimicrobial secondary metabolites, including gelatinomycin, trichomycin, chlorotrichomycin, and antibacterial peptides (Maruyama *et al.*, 2020). According to Nawrocka *et al.* (2018), secondary metabolites can promote plant growth, act as antibacterial agents, and provide valuable resources for the development of agricultural antibiotics.

With a 54.81% inhibition rate, Naglot *et al.* (2015) discovered that *T. viride* metabolites significantly inhibited the wilt-specific form of *F. oxysporum*. Manganiello *et al.* (2018) discovered that when exposed to *T. viride* TG050 609's volatile secondary metabolites, *P. nicotianae* mycelium can grow erratically, fracture, or even dissolve. This suggests that *T. viride* has antibiosis activity against *P. nicotianae*. Furthermore, the majority of *Trichoderma* strains can produce antimicrobial compounds such as pentaibols, which can inhibit a variety of plant pathogenic fungi and work in tandem with their cell wall-degrading enzymes to effectively stop their growth (Debode *et al.*, 2018; Mayo-Prieto *et al.*, 2019; Kovacs *et al.*, 2021; Martinez-Salgado *et al.*, 2021).



**Fig. 1.** Schematic diagram of the mechanism of action of *Trichoderma* in plant disease control.

## CONCLUSION AND FUTURE SCOPE

Chemical control is currently the primary method for controlling plant diseases, and it is accomplished by misting fungicides and pesticides. Despite the fact that chemical pesticides have a positive and beneficial effect on agricultural productivity, their improper application has seriously contaminated the environment and increased pathogen resistance. Numerous studies have shown that *Trichoderma* can reduce the use of chemical pesticides while also providing beneficial biological control effects. More efficient and appropriate strains must be discovered to join the biocontrol team, as there are currently few *Trichoderma* biocontrol agents on the market (Nieto-Jacobo *et al.*, 2017; Fiorentino *et al.*, 2018; Lopez *et al.*, 2019; Nawrocka *et al.*, 2019; Poveda *et al.*, 2019; Cabral-Miramonte *et al.*, 2019).

While *Trichoderma* has many applications in agriculture, there are still some issues with its development and application (Rubio *et al.*, 2014; Zhang *et al.*, 2018; Phoka *et al.*, 2020; Santos *et al.*, 2020; Wang *et al.*, 2022). When applied in the field, the *Trichoderma* spore preparation is typically a living fungal preparation that is frequently influenced by various natural factors such as humidity, temperature, soil acidity, alkalinity, and the soil microbial community, reducing the biological control effect and making field test performance unstable. Furthermore, biological control agents have a limited shelf life, and some microorganisms must be refrigerated to maintain a viable concentration at the time of application.

A prospective investigation into the biological and environmental safety of transgenic *Trichoderma* should be conducted concurrently with additional research on the organism (Li *et al.*, 2021). The identification of *Trichoderma* elicitors to recognize plant targets or receptors, the balance regulation of *Trichoderma* colonizing host and plant immune response, the long-distance and trans-growth period transduction mechanism of systematically induced plant disease resistance and its defence signals, and the mechanism of *Trichoderma*-induced plant endophytic microbiome to synergistically stimulate plant immune response have all recently attracted attention from researchers. Research is beginning to emerge on the mechanism of cross-border miRNA transduction between pathogenic microorganisms, plants, and *Trichoderma*, as well as the regulation of the host process and plant immune response to *Trichoderma* colonization.

Combining *Trichoderma* and other microorganisms has made it possible to broaden the target spectrum of microbial metabolites, develop new biopesticides and biostimulants based on metabolites, and discover new metabolites with specific microorganism functions (Wang *et al.*, 2022). It is expected that developing new plant immune-activating protein pesticides and molecularly modifying the *Trichoderma* multi-stimulator fusion protein will open up new avenues for the development of macromolecular biopesticides. Currently, there is an urgent need to identify the synergistic relationships that exist between *Trichoderma*, plants, and pathogenic microorganisms in

order to induce disease resistance on a cross-genome level. Furthermore, new biostimulant or products based on *Trichoderma* and other microbial symbiotic agents must be developed in order to treat diseases and pests. Compound biocontrol fungi outperform single-life biocontrol fungi in terms of disease resistance, environmental adaptation, and control efficacy. Although there are numerous preparations containing various *Trichoderma* species that are used in sustainable agricultural crops, their application is still expensive and not available to all farmers. In the process of developing biocontrol agents, the use of compatible or affinity multiple microorganisms for compounding has grown popular. *Trichoderma* can form alliances with a variety of microorganisms, including fungi and bacteria, to improve plants ability to manage and prevent disease. The primary areas of research for *Trichoderma* as a biocontrol fungus may be as follows.

The ability of the biocontrol agent *Trichoderma* to withstand stressors such as high temperatures, drying, UV radiation, and storage conditions such as more than a year at room temperature is critical for commercial application. At the moment, two primary technologies exist. There are two ways to induce *Trichoderma* to produce stress-resistant chlamydospores: one involves lowering the acidity and controlling oxygen utilization, and the other involves adding chemical additives (such as copper) to the inoculum. The secret to understanding how *Trichoderma* induces plant immunity is to examine how its effectors interact with plant cell receptors. Prospective studies on the biological and environmental safety of transgenic *Trichoderma* should take place concurrently with the advancement of transgenic research.

## REFERENCES

- Abbas, A., Mubeen, M., Zheng, H., Sohail, M. A., Shakeel, Q., and Solanki, M. K. (2022). *Trichoderma* spp. genes involved in the biocontrol activity against *Rhizoctonia solani*. *Frontiers in Microbiology*, 13, 884469.
- Aggarwal, R., Srivastava, K. D., Singh, D. V., Bahadur, P. and Nagarajan, S. (1993). Possible biocontrol of loose smut of wheat. *Indian Journal of Biological Control*, 6, 111-112.
- Al-Surhane, A. A. (2022). Protective role of antifusarial eco-friendly agents (*Trichoderma* and salicylic acid) to improve resistance performance of tomato plants. *Saudi Journal of Biological Sciences*, 29, 2933–2941.
- Alvarez-Garcia, S., Mayo-Prieto, S., Gutierrez, S. and Casquero, P. A. (2020). Self-inhibitory activity of *Trichoderma* soluble metabolites and their antifungal effects on *Fusarium oxysporum*. *Journal of Fungi*, 6, 176.
- Al-Askar, A. A., Saber, W., Ghoneem, K. M., Hafez, E. E. and Ibrahim, A. A. (2021). Crude citric acid of *Trichoderma asperellum*: tomato growth promoter and suppressor of *Fusarium oxysporum* f. sp. *lycopersici*. *Plants* 10, 222.
- Alfiky, A. and Weisskopf, L. (2021). Deciphering *Trichoderma*-plant-pathogen interactions for better development of biocontrol applications. *Journal of Fungi*, 7, 61.
- Andrade-Hoyos, P., Silva-Rojas, H. V. and Romero-Arenas, O. (2020). Endophytic *Trichoderma* species isolated from *Persea americana* and *Cinnamomum verum* roots reduce symptoms caused by *Phytophthora cinnamomi* in avocado. *Planting Theory*, 9, 1220.
- Amin, F., Razdan, V. K., Mohiddin, F. A., Bhat, K. A. and Banday, S. (2010). Potential of *Trichoderma* species as Bio-control agents of Soil borne fungal propagules. *Journal of Phytology*, 2(10), 38–41.
- Basu, A. (2009). Employing eco-friendly potato disease management allows organic tropical Indian production systems to prosper. *Asian Journal of Food & Agro-Industry*, S80-S87.
- Bhandari, P. C. and Vishunavat, K. (2013). Screening of different isolates of *Trichoderma harzianum* and *Pseudomonas fluorescens* against *Fusarium moniliforme*. *Pantnagar. Journal of Research*, 11(2), 243-247.
- Bhardwaj, S. S. and Gupta, P. K. (1987). *In vitro* antagonism of *Trichoderma* spp. against fungal pathogens associated with rhizome rot of ginger. *Indian Journal of Plant Pathology*, 5, 41-42.
- Bhat, K. A., Ali, A. and Wani, A. H. (2009). Evaluation of biocontrol agents against *Rhizoctonia solani* Kuhn and sheath blight disease of rice under temperate ecology. *Plant Disease Research*, 24(1), 15-18.
- Bhramaramba, S. and Nagamani, A. (2013). Antagonistic *Trichoderma* Isolates to control bakane pathogen of rice. *Agricultural Science Digest*, 33(2), 104-108.
- Biswas, S. and Datta, M. (2013). Evaluation of Biological Control Agents against Sheath Blight of Rice in Tripura. *Indian Phytopathology*, 66 (1), 77-80.
- Bliss, D. E. (1951). The destruction of *Armillaria mellea* in citrus soils: *Phytopathology*, 41, 655-683.
- Bunker, R. N. and Mathur, K. (2001). Antagonism of local biocontrol agents to *Rhizoctonia solani* inciting dry rot of chilli. *Journal of Mycology and Plant Pathology*, 31, 50-53.
- Calvet, C., Pera, J. and Brea, J. M. (1990). Interaction of *Trichoderma* spp. with *Glomus mosseae* and two wilt pathogenic fungi. *Agriculture Ecosystem Environment*, 29, 59-65.
- Carvalho, D. D. C., Mello, S. C. M. de, Martins, I. and Lobo, M. (2015). *Tropical Plant Pathology*, 40, 375-381
- Carro-Huerga, G., Compant, S., Gorfer, M., Cardoza, R. E., Schmoll, M., and Gutierrez, S. (2020). Colonization of *Vitis vinifera* L. by the endophyte *Trichoderma* sp. strain T154: biocontrol activity against *Phaeoacremonium minimum*. *Frontiers in Plant Science*, 11, 1170.
- Chakravarthy, S., Nagamani, K., Ratnakumari, A. R. and Bramarambha, Y. S. (2011). Antagonistic ability against *Rhizoctonia solani* and pesticide tolerance of *Trichoderma* strains. *Advances in Environmental Biology*, 5(9), 2631-2638.
- Chaudhary, R. G. and Prajapati, R. K. (2004). Comparative efficacy of Fungal Bio-agents against *Fusarium udum*. *Annals of Plant Protection Science*, 12, 75-79.
- Chen, J., Zhou, L., Din, I. U., Arafat, Y., Li, Q., and Wang, J. (2021). Antagonistic activity of *Trichoderma* spp. against *Fusarium oxysporum* in rhizosphere of radix pseudostellariae triggers the expression of host defense genes and improves its growth under long-term monoculture system. *Frontiers in Microbiology*, 12, 579920.
- Cole, J. and Zrinyika, S. (1988). Integrated control of *Rhizoctonia solani* and *Fusarium solani* in tobacco transplant with *Trichoderma harzianum* and triadimenol. *Plant Pathology* (London), 37, 271-277.

- Contreras-Cornejo, H. A., Macias-Rodriguez, L., Alfaro-Cuevas, R. and Lopez-Bucio, J. (2014). *Molecular Plant-Microbe Interactions*, 27, 503-514.
- Cabral-Miramontes, J. P., Olmedo-Monfil, V., Lara-Banda, M., Zuniga-Romo, E. R. and Arechiga-Carvajal, E. T. (2022). Promotion of plant growth in arid zones by selected *Trichoderma* spp. strains with adaptation plasticity to alkaline pH. *Biology*, 11, 1206.
- Degani, O., Khatib, S., Becher, P., Gordani, A. and Harris, R. (2021a). *Trichoderma asperellum* secreted 6-pentyl- $\alpha$ -pyrone to control *Magnaportheopsis maydis*, the maize late wilt disease agent. *Biology*, 10, 897.
- Deve, P. R. and Dutta, B. K. (1988). *Trichoderma* as a biocontrol agent against *Sclerotium rolfsii*. *Microbios. Letters*, 37, 107-114.
- Domsch, K. H., Gams, W. and Anderson, T. H. (1980). Compendium of Soil fungi. Academic Press, London. p. 794-809.
- Damodaran, T., Rajan, S., Muthukumar, M., Gopal, R., Yadav, K., and Kumar, S. (2020). Biological management of banana Fusarium wilt caused by *Fusarium oxysporum* f. sp. cubense tropical race 4 using antagonistic fungal isolate CSR-T-3 (*Trichoderma reesei*). *Frontiers in Microbiology*, 11, 595845.
- Dubey, S. C., Bhavani, R. and Singh, B. (2009). Development of Pusa 5SD for seed dressing and Pusa Biopellet 10G for soil application formulations of *Trichoderma harzianum* and their evaluation for integrated management of dry root rot of mungbean (*Vigna radiata*). *Biological Control*, 50, 231-242.
- Debode, J., De Tender, C., Cremelie, P., Lee, A. S., Kyndt, T., and Muylle, H. (2018). *Trichoderma*-inoculated miscanthus straw can replace peat in strawberry cultivation, with beneficial effects on disease control. *Frontiers Plant Science*, 9, 213.
- Degani, O. and Dor, S. (2021). *Trichoderma* biological control to protect sensitive maize hybrids against late wilt disease in the field. *Journal of Fungi*, 7, 315.
- Dugassa, A., Alemu, T. and Woldehawariat, Y. (2021). In-vitro compatibility assay of indigenous *Trichoderma* and *Pseudomonas* species and their antagonistic activities against black root rot disease (*Fusarium solani*) of faba bean (*Vicia faba* L.). *BMC Microbiology*, 21, 115.
- Elad, Y., Harder, Y., Chet, I. and Henis, Y. (1986). Prevention with *Trichoderma harzianum* Rifaiagr. infection by *Sclerotium rolfsii* sacc and *Rhizoctonia solanii* kuhn of soil fumigated with methyl bromide, and improvement of disease control in tomatoes and peanuts. *Crop Protection*, 1, 199-211.
- El-Hasan, A., Walker, F., Klaiher, I., Schöne, J., Pfannstiel, J. and Voegelé, R. T. (2022). New approaches to manage Asian soybean rust (*Phakopsora pachyrhizi*) using *Trichoderma* spp. or their antifungal secondary metabolites. *Meta*, 12, 507.
- Fernandez, M. R. (1992). The effect of *Trichoderma harzianum* on fungal pathogens infesting wheat and blackoat straw. *Soil Biology and Biochemistry*, 24, 1031-1034.
- Fiorentino, N., Ventrino, V., Woo, S. L., Pepe, O., De Rosa, A., and Gioia, L. (2018). *Trichoderma*-based biostimulants modulate rhizosphere microbial populations and improve N uptake efficiency, yield, and nutritional quality of leafy vegetables. *Frontiers in Plant Science*, 9, 743.
- Fontana, D. C., de Paula, S., Torres, A. G., de Souza, V., Pascholati, S. F., and Schmidt, D. (2021). Endophytic fungi: biological control and induced resistance to phytopathogens and abiotic stresses. *Pathogens*, 10, 570.
- Gangwar, O. P. and Sharma, P. (2013). A growth and survival of *Trichoderma harzianum* and *Pseudomonas fluorescens* on different substrates and their temporal and spatial population dynamics in irrigated rice ecosystem. *Annals of Plant Protection Science*, 21(1), 104-108.
- Garnica-Vergara, A., Barrera-Ortiz, S., Munoz-Parra, E., Raya-Gonzalez, J., Mendez-Bravo, A., Macias-Rodriguez, L., Ruiz-Herrera, L. F. and Lopez-Bucio, J. (2016). *New Phytologist*, 209, 1496-1512.
- Gogoi, R., Saikia, M., Helim, R. and Ullah, Z. (2007). Management of Potato Diseases using *Trichoderma viride* formulations. *Journal of Mycology and Plant Pathology*, 37, 227-230.
- Gomathinayagam, S., Rekha, M., Murugan, S. S. and Jagessar, J. C. (2010). The biological control of paddy disease brown spot (*Bipolaris oryzae*) by using *Trichoderma viride* in vitro condition. *Journal of Biopesticides*, 3(1), 93-95.
- Gupta, S. B., Thakur, K. S., Singh, A., Tamrakar, D. K. and Thakur, M. P. (2005). Efficacy of *Trichoderma viride* and *Rhizobium* against Wilt complex of Chickpea in field. *Journal of Mycology and Plant Pathology*, 35, 89-91.
- Haggag, K. H. E. and El-Gamal, N. G. (2012). In vitro study on *Fusarium solani* and *Rhizoctonia solani* isolates causing the damping off and root rot diseases in tomatoes. *Nature and Science*, 10, 16-25.
- Harman, G. E., Howell, C. R., Viterbo, A., Chet, I. and Lorito, M. (2004a). *Trichoderma* species-opportunistic, a virulent plant symbiont. *Nature Reviews Microbiology*, 2, 43-56.
- Harman, G. E., Obregon, M. A., Samuels, G. and Lorito, M. (2010). Changing models of biocontrol in the developing and developed world. *Plant Disease*, 94 (8), 928-939.
- Herridge, D. F., Peoples, M. B. and Boddey, R. M. (2008). Global inputs of biological nitrogen fixation in agricultural systems. *Plant Soil*, 311(1), 1-18.
- Haouhach, S., Karkachi, N., Oguiba, B., Sidaoui, A., Chamorro, I., and Kihal, M. (2020). Three new reports of *Trichoderma* in Algeria: *T. atrobrunneum*, (South) *T. longibrachiatum* (South), and *T. afroharzianum* (Northwest). *Microorganisms*, 8, 1455.
- Howell, C. R. (2003). Mechanisms employed by *Trichoderma* species in the biological control of plant diseases: the history and evolution of current concepts. *Plant Disease*, 87, 4-10.
- Intana, W., Kheawlang, S. and Sunpapao, A. (2021). *Trichoderma asperellum* T76-14 released volatile organic compounds against postharvest fruit rot in muskmelons (*Cucumis melo*) caused by *Fusarium incarnatum*. *Journal of Fungi*, 7, 46.
- Izquierdo-García, L. F., Gonzalez-Almario, A., Cotes, A. M. and Moreno-Velandia, C. A. (2020). *Trichoderma virens* G1006 and *Bacillus velezensis* Bs006: a compatible interaction controlling *Fusarium* wilt of cape gooseberry. *Scientific Reports*, 10, 6857.
- Jat, J. G. and Agalave, H. R. (2013). Antagonistic properties of *Trichoderma* species against oilseed-borne fungi. *Science Research Reporter*, 3(2), 171-174.
- Jogaiah, S., Abdelrahman, M., Tran, L. S. P. and Ito, S. I. (2018). *Molecular Plant Pathology*, 19, 870-882.
- John, P. R., Tyagi, R. D., Prevost, D., Brar, S. K., Pouleur, S. and Surampalli, R. Y. (2010). Mycoparasitic *Trichoderma viride* as a biocontrol agent against *Fusarium oxysporum* f. sp. *adzuki* and *Pythium*

- arrhenomanes* and as a growth promoter of soybean. *Crop Protection*, 29, 1452-1459.
- John, R. P., Tyagi, R., Prevost, D., Brar, S. K., Pouleur, S. and Surampalli, R. (2010). *Crop Protection*, 29, 1452-1459.
- Kapoor, A. S. (2008). Biocontrol potential of *Trichoderma* spp. against important soil borne diseases of vegetable crops. *Indian Phytopathology*, 61(4), 492-498.
- Kexiang, G., Xiaoguang, L., Yonghong, L., Tianbo, Z. and Shuliang, W. (2002). Potential of *Trichoderma harzianum* and *T. atroviride* to control *Botryosphaeria berengeriana* f. sp. *piricola*, the cause of apple ring rot. *Journal of Phytopathology*, 150, 271-276.
- Khodke, S. W. and Raut, B. T. (2010). Management of root rot collar rot of soybean. *Indian Phytopathology*, 63(3), 298-301.
- Komy, M. H. E., Saleh, A. A., Eranthodi, A. and Molan, Y. Y. (2015). Characterization of Novel *Trichoderma asperellum* Isolates to Select Effective Biocontrol Agents Against Tomato Fusarium Wilt. *Plant Pathology Journal*, 31(1), 50-60.
- Krivoshchekova, T. G. and Mishchenk, V. S. (1990). The effectiveness of trichodermin. *Zashchita rastenii* (Moskova) No. 11, 22 (Ru).
- Kottb, M., Gigolashvili, T., Grobkinsky, D. K. and Piechulla, B. (2015). *Trichoderma volatiles* effecting Arabidopsis: from inhibition to protection against phytopathogenic fungi. *Front. Microbiol.*, 6, 995.
- Kovacs, C., Csoto, A., Pál, K., Nagy, A., Fekete, E., Karaffa, L., et al. (2021). The biocontrol potential of endophytic *Trichoderma* fungi isolated from Hungarian grapevines. Part I. isolation, identification and in vitro studies. *Pathogens*, 10, 1612.
- Lazazzara, V., Vicelli, B., Bueschl, C., Parich, A., Pertot, I. and Schuhmacher, R. (2021). *Trichoderma* spp. volatile organic compounds protect grapevine downy by activating defense-related processes against downy mildew. *Physiologia Plantarum*, 172, 1950-1965.
- Leal, C., Richet, N., Guise, J. F., Gramaje, D., Armengol, J. and Fontaine, F. (2021). Cultivar contributes to the beneficial effects of *Bacillus subtilis* PTA-271 and *Trichoderma atroviride* SC1 to protect grapevine against *Neofusicoccum parvum*. *Frontiers in Microbiology*, 12, 726132.
- Lindow, S. E. (1985). Foliar Antagonists: Status and Prospects. In: *Biological Control in Agricultural IPM Systems* (Eds. Hoy, M.A. and Herzog, D.C.), Academic Press, Inc. (London), pp. 395-413.
- Li, W. C., Lin, T. C., Chen, C. L., Liu, H. C., Lin, H. N., and Chao, J. L. (2021). Complete genome sequences and genome-wide characterization of *Trichoderma* biocontrol agents provide new insights into their evolution and variation in genome organization, sexual development, and fungal-plant interactions. *Microbiology Spectrum*, 9, e0066321.
- Lopez, A. C., Alvarenga, A. E., Zapata, P. D., Luna, M. F. and Villalba, L. L. (2019). *Trichoderma* spp. from Misiones, Argentina: effective fungi to promote plant growth of the regional crop *Ilex paraguariensis* St. Hil. *Mycology*, 10, 210-221.
- Marzano, M., Gallo, A. and Altomare, C. (2013). Improvement of biocontrol efficacy of *Trichoderma harzianum* vs. *Fusarium oxysporum* f. sp. *lycopersici* through UV induced tolerance to fusaric acid. *Biological control*, 67, 397-408.
- Mastouri, F., Bjorkman, T. and Harman, G. E. (2010). *Phytopathology*, 100, 1213-1221
- Manganiello, G., Nicastro, N., Caputo, M., Zaccardelli, M., Cardi, T. and Pane, C. (2021). Functional hyperspectral imaging by high-related vegetation indices to track the wide-spectrum *Trichoderma* biocontrol activity against soil-borne diseases of baby-leaf vegetables. *Frontiers in Plant Science*, 12, 630059.
- Meena, M., Swapnil, P., Zehra, A., Dubey, M. K. and Upadhyay, R. S. (2017). *Archives Phytopathology Plant Protection*, 50, 629-648.
- Mishra, D. S., Gupta, A. K., Prajapati, C. R. and Singh, U. S. (2011). Combination of fungal and bacterial antagonists for management of root and stem rot disease of soybean. *Pakistan Journal of Botany*, 43(5), 2569-2574.
- Mishra, R. K. and Gupta, R. P. (2012). In vitro evaluation of plant extracts, bioagents and fungicides against purple blotch and *Stemphylium* blight of onion. *Journal of Medicinal Plants Research*, 6(48), 5840-5843.
- Mukherjee, P. K., Horwitz, B. A. and Kenerley, C. M. (2012). *Microbiology*, 158, 35-45.
- Mayo-Prieto, S., Marra, R., Vinale, F., Rodríguez-Gonzalez, A., Woo, S. L., and Lorito, M. (2019). Effect of *Trichoderma velutinum* and *Rhizoctonia solani* on the metabolome of bean plants (*Phaseolus vulgaris* L.). *International Journal of Molecular Sciences*, 20, 549.
- Martinez-Salgado, S. J., Andrade-Hoyos, P., Parraguirre Lezama, C., Rivera-Tapia, A., Luna-Cruz, A. and Romero-Arenas, O. (2021). Biological control of charcoal rot in peanut crop through strains of *Trichoderma* spp., in Puebla, Mexico. *Plan. Theory*, 10, 2630.
- Moran-Diez, M. E., Tranque, E., Bettiol, W., Monte, E. and Hermosa, R. (2020). Differential response of tomato plants to the application of three *Trichoderma* species when evaluating the control of *Pseudomonas syringae* populations. *Plan. Theory*, 9, 626.
- Manganiello, G., Sacco, A., Ercolano, M. R., Vinale, F., Lanzuise, S., and Pascale, A. (2018). Modulation of tomato response to *Rhizoctonia solani* by *Trichoderma harzianum* and its secondary metabolite harzianic acid. *Frontiers in Microbiology*, 9, 1966.
- Maruyama, C. R., Bilesky-Jose, N., de Lima, R. and Fraceto, L. F. (2020). Encapsulation of *Trichoderma harzianum* preserves enzymatic activity and enhances the potential for biological control. *Frontiers in Bioengineering and Biotechnology*, 8, 225.
- Ng, L. C., Ngadin, A., Azhari, M. and Zahari, N. A. (2015). Potential of *Trichoderma* spp. as Biological Control Agents against Bakanae Pathogen (*Fusarium fujikuroi*) in Rice. *Asian Journal of Plant Pathology*, 9, 46-58.
- Nawrocka, J., Małolepsza, U., Szymczak, K. and Szczech, M. (2018). Involvement of metabolic components, volatile compounds, PR proteins, and mechanical strengthening in multilayer protection of cucumber plants against *Rhizoctonia solani* activated by *Trichoderma atroviride* TRS25. *Protoplasma*, 255, 359-373.
- Naglot, A., Goswami, S., Rahman, I., Shrimali, D. D., Yadav, K. K., and Gupta, V. K. (2015). Antagonistic potential of native *Trichoderma viride* strain against potent tea fungal pathogens in North East India. *Plant Pathology Journal*, 31, 278-289.
- Nawrocka, J., Gromek, A. and Małolepsza, U. (2019). Nitric oxide as a beneficial signaling molecule in *Trichoderma atroviride* TRS25- induced systemic defense responses of cucumber plants against *Rhizoctonia solani*. *Frontiers of Plant Science*, 10, 421.

- Nieto-Jacobo, M. F., Steyaert, J. M., Salazar-Badillo, F. B., Nguyen, D. V., Rostas, M., and Braithwaite, M. (2017). Environmental growth conditions of *Trichoderma* spp. affect Indole acetic acid derivatives, volatile organic compounds, and plant growth promotion. *Frontiers of Plant Science*, 8, 102.
- Pal, G. K., Pooja and Kumar, P. (2013). Enhancing seed germination of maize and soybean by using botanical extracts and *Trichoderma harzianum*. *Current Disease*, 2(1), 72-75.
- Pandey, S. and Pundhir, V. S. (2013). Mycoparasitism of potato black scurf pathogen (*Rhizoctonia solani* Kuhn) by biological control agents to sustain production. *Indian Journal of Horticulture*, 70(1), 71-75.
- Pant, R. and Mukhopadhyay, A. N. (2001). Integrated management of seed and seedling rot complex of Soybean. *Indian Phytopathology*, 54(3), 346-350.
- Papavizas, G. C. (1985). *Trichoderma* and *Gliocladium*: Biology, ecology and potential for biocontrol. *Annual Review of Phytopathology*, 22, 23-54.
- Parveen, S. and Kumar, V. R. (2004). Antagonism by *Trichoderma viride* against leaf blight pathogen of wheat. *Journal of Mycology and Plant Pathology*, 34, 220-222.
- Persoon, C. H. (1794). Neuer Veersucheinersystematische Eintheilung der Schwamme (Dispositio methodical fungorum). *Romer's News Magazine Bot.*, 1, 63-128.
- Prakasam, V. and Sharma, P. (2012). *Trichoderma harzianum* (Th-3) a potential strain to manage the purple blotch of onion (*Allium cepa* L.) caused by *Alternaria porri* under North Indian plains. *Journal of Agriculture Science*, 10, 266-272.
- Prasad, K. S. K., Kulkarni, S. and Siddaramaiah, A. L. (1978). Antagonistic action of *Trichoderma* spp. and *Streptomyces* sp. on *Drechslera sativum* (*Helminthosporium sativum*) Pam. King and Bakke) Subram. and Jain. *Current Research*, 12, 202-203.
- Prevost, D., Bertrand, A., Juge, C. and Chalifour, F. P. (2010). Elevated CO<sub>2</sub> induces differences in nodulation of soybean depending on brady rhizobium strain and method of inoculation. *Plant Soil*, 331(1), 115-127.
- Pollard-Flamand, J., Boulé, J., Hart, M. and Urbez-Torres, J. R. (2022). Biocontrol activity of *Trichoderma* species isolated from grapevines in British Columbia against botryosphera dieback fungal pathogens. *Journal of Fungi*, 8, 409.
- Poveda, J., Hermosa, R., Monte, E. and Nicolás, C. (2019). *Trichoderma harzianum* favours the access of arbuscular mycorrhizal fungi to non-host Brassicaceae roots and increases plant productivity. *Scientific Reports*, 9, 11650.
- Phoka, N., Suwannarach, N., Lumyong, S., Ito, S. I., Matsui, K., Arikiti, S. (2020). Role of volatiles from the endophytic fungus *Trichoderma asperelloides* PSU-P1 in biocontrol potential and in promoting the plant growth of *Arabidopsis thaliana*. *Journal of Fungi*, 6, 341.
- Ray, A., Kumar, P. and Tripathi, H. S. (2007). Evaluation of bioagents against *Rhizoctonia solani* Kuhn the cause of aerial blight of soybean. *Indian Phytopathology*, 60(4), 532-534.
- Rees, H. J., Drakulic, J., Cromey, M. G., Bailey, A. M. and Foster, G. D. (2022). Endophytic *Trichoderma* spp. can protect strawberry and privet plants from infection by the fungus *Armillaria mellea*. *PLoS One*, 17, e0271622.
- Risoli, S., Cotrozzi, L., Sarrocco, S., Nuzzaci, M., Pellegrini, E. and Vitti, A. (2022). *Trichoderma* induced resistance to *Botrytis cinerea* in *Solanum* species: a meta-analysis. *Planning Theory*, 11, 180.
- Rini, C. R. and Sulochana, K. K. (2006). Management of seedling rots of chilli (*Capsicum annuum* L.) using *Trichoderma* spp. and fluorescent pseudomonads (*Pseudomonas fluorescens*). *Journal of Tropical Agriculture*, 44(1-2), 79-82.
- Ru, Z. and Di, W. (2012). *Trichoderma* spp. from rhizosphere soil and their antagonism against *Fusarium sambucinum*. *African Journal of Biotechnology* 11, 4180-4186.
- Rubio, M. B., Quijada, N. M., Pérez, E., Domínguez, S., Monte, E. and Hermosa, R. (2014). Identifying beneficial qualities of *Trichoderma parareesei* for plants. *Applied and Environmental Microbiology*, 80, 1864-1873.
- Sanchez-Montesinos, B., Santos, M., Moreno-Gavira, A., Marin-Rodulfo, T., Gea, F. J. and Diane, F. (2021). Biological control of fungal diseases by *Trichoderma aggressivum* f. *europaeum* and its compatibility with fungicides. *Journal of Fungi*, 7, 598.
- Selvakumar, R. (2008). Bioformulations for management of late light of potato in North eastern India. Third International Late Blight Conference, Beijing, April 3-5, 2008.
- Shahnaz, E., Razdan, V. K., Rizvi, S. E. H., Rather, T. R., Gupta, S. and Andrabi, M. (2013). Integrated disease management of foliar blight disease of onion: A case study of application of confounded factorials. *Journal of Agricultural Science*, 5(1), 17-22.
- Sharma, M., Sharma, P., Raja, M., Kumar, K., Chandra, S. and Sharma, R. (2016). *International Journal of Current Microbiology and Applied Sciences*, 5, 382-386.
- Siddiquee, S., Cheong, B. E., Taslima, K., Kausar, H. and Hasan, M. M. (2012). *Journal of Chromatographic Science*, 50, 358-367.
- Singh, D. P. (2004). Use of reduced dose of fungicides and seed treatment with *Trichoderma viride* to control wheat loose smut. *Journal of Mycology and Plant Pathology*, 34, 396-397.
- Singh, V., Srivastava, K. D. and Aggarwal, R. (1991). Significant findings under PL-480 Karnal bunt Project at Delhi centre. 30th All India Wheat Res. Workers' Workshop. Kanpur (India), pp. 1-17.
- Singh, U. and Thapliyal, P. N. (1998). Effect of inoculum density, host cultivars and seed treatment on the seed and seedling rot of soybean caused by *Sclerotium rolfsii*. *Indian Phytopathology*, 51(3), 244-246.
- Santos, M., Santos, L., Costa, D., Vieira, T. A. and Lustosa, D. C. (2020). *Trichoderma* spp. on treatment of *Handroanthus serratifolius* seeds: effect on seedling germination and development. *Heliyon*, 6, e04044.
- SOPA report (2018). Application of fungicides and *Trichoderma viride* in management of soybean pod blight complex. *International Journal of Chemical Studies*, 6(4), 3208-3210.
- Shobha, B., Lakshmeesha, T. R., Ansari, M. A., Almatroudi, A., Alzohairy, M. A., and Basavaraju, S. (2020). Mycosynthesis of Zinc oxide nanoparticles using *Trichoderma* spp. isolated from rhizosphere soils and its synergistic antibacterial effect against *Xanthomonas oryzae* pv. *oryzae*. *Journal of Fungi*, 6, 181.
- Tilocca, B., Cao, A. and Migheli, Q. (2020). Scent of a killer: microbial volatiles and its role in the biological control of plant pathogens. *Frontiers in Microbiology*, 11, 41.

- Tapwal, A., Sharma, Y. P. and Lakhanpal, T. N. (2005). Use of biocontrol agents against white root rot of apple. *Indian Journal of Mycology and Plant Pathology*, 35, 67-69.
- Tyszkiewicz, R., Nowak, A., Ozimek, E. and Jaroszuk-Sciseł, J. (2022). *Trichoderma*: the current status of its application in agriculture for the biocontrol of fungal phytopathogens and stimulation of plant growth. *International Journal of Molecular Sciences*, 23, 2329.
- Vargas, W. A., Mukherjee, P. K., Laughlin, D., Wiest, A., Moran-Diez, M. E. and Kenerley, C. M. (2014). *Microbiology*, 160, 2319-2330.
- Vasanthakumari, M. M. and Shivanna, M. B. (2013). Biological control of anthracnose of chilli with rhizosphere and rhizoplane fungal isolates from grasses. *Archives of Phytopathology and Plant Protection*, 46(14), 1641-1666.
- Vicente, I., Baroncelli, R., Morán-Diez, M. E., Bernardi, R., Puntoni, G., and Hermosa, R. (2020). Combined comparative genomics and gene expression analyses provide insights into the terpene synthases inventory in *Trichoderma*. *Microorganisms*, 8, 1603.
- Vinale, Sivasithamparam, F. K., Ghisalberti, L. E., Marra, R., Woo, L. S. and Lorito, M. (2008). *Trichoderma* plant-pathogen interactions. *Soil Biology and Biochemistry*, 40, 1-10.
- Wang, R., Liu, C., Jiang, X., Tan, Z., Li, H., and Xu, S. (2022). The newly identified *Trichoderma harzianum* partiti virus (ThPV2) does not diminish spore production and biocontrol activity of its host. *Viruses*, 14, 1532.
- Wang, Y., Chen, H., Ma, L., Gong, M., Wu, Y. and Bao, D. (2022). Use of CRISPR-Cas tools to engineer *Trichoderma* species. *Microbial Biotechnology*, 15, 2521–2532.
- Weindling R. (1932). *Trichoderma lignorum* a parasite of other soil fungi. *Phytopathology*, 22, 837-845.
- Weindling, R. (1934). Studies on lethal principle effective in the parasitic action of *Trichoderma lignorum* on *Rhizoctinia solani* and other soil fungi. *Phytopathology*, 24, 1153-1179.
- Wells, H. D., Bell, D.K. and Casimir, A. J. (1972). Efficacy of *Trichoderma harzianum* as a biocontrol for *Sclerotium rofsii*. *Phytopathology*, 62, 442-447.
- Wilson, M., Crawford, E. K. and Campbell, R. (1988). Biocontrol by *Trichoderma harzianum* of damping off lettuce caused by *Rhizoctonia solani*. *Bulletin OEPP*, 18, 83-90.
- Wright, J. M. (1956). The production of antibiotics in soil III. Production of gliotoxin in wheat straw buried in soil. *Annals of Applied Biology*, 44, 461-466.
- Yadav, J., Verma, J. P. and Tiwari, K. N. (2011). *Asian Journal of Biological Sciences*, 4, 291-299.
- Yadav, R. K. and Majumdar, V. L. (2005). Efficacy of plant extracts, biological agents and fungicides against *Lasiodiplodia theobromae* causing die back of guava (*Psidium guajava* L.). *Journal of Mycology and Plant Pathology*, 35: 352-353.
- Yadav, S. K., Dave, A., Sarkar, A., Singh, H. B. and Sharma, B. K. (2013). Co-inoculated biopriming with *Trichoderma*, *Pseudomonas* and *Rhizobium* improves crop growth in *Cicer arietinum* and *Phaseolus vulgaris*. *International Journal of Agriculture Environment & Biotechnology*, 6(2), 255.
- Zhang, F., Huo, Y., Cobb, A. B., Luo, G., Zhou, J. and Yang, G. (2018). *Trichoderma* biofertilizer links to altered soil chemistry, altered microbial communities, and improved grassland biomass. *Frontiers in Microbiology*, 9, 848.
- Zhang, S., Gan, Y., Liu, J., Zhou, J. and Xu, B. (2020). Optimization of the fermentation media and parameters for the bio-control potential of *Trichoderma longibrachiatum* T6 against nematodes. *Frontiers in Microbiology*, 11, 574601.
- Zhang, C., Wang, W., Xue, M., Liu, Z., Zhang, Q., and Hou, J. (2021). The combination of a biocontrol agent *Trichoderma asperellum* SC012 and hymexazol reduces the effective fungicide dose to control *Fusarium* wilt in cowpea. *Journal of Fungi*, 7, 685.
- Zhang, C., Wang, W., Hu, Y., Peng, Z., Ren, S., and Xue, M. (2022). A novel salt-tolerant strain *Trichoderma atroviride* HN082102.1 isolated from marine habitat alleviates salt stress and diminishes cucumber root rot caused by *Fusarium oxysporum*. *BMC Microbiology*, 22, 67.
- Zhang, Y., Xiao, J., Yang, K., Wang, Y., Tian, Y. and Liang, Z. (2022). Transcriptomic and metabolomic insights into the biocontrol mechanism of *Trichoderma asperellum* M45a against watermelon *Fusarium* wilt. *PLoS One*, 17, e0272702.
- Zheng, H., Qiao, M., Lv, Y., Du, X., Zhang, K. Q. and Yu, Z. (2021). New species of *Trichoderma* isolated as endophytes and saprobes from Southwest China. *Journal of Fungi*, 7, 467.

**How to cite this article:** Manjula, Aishwarya, Ajay Kumar Gautam and Anupam Kumar (2024). *Trichoderma* as Potential Biocontrol Agent on Diseases of Soybean (*Glycine max* L.): A Comprehensive Review. *Biological Forum – An International Journal*, 16(10): 117-126.