



Effect of Different Level of Micronutrients on Growth, Flowering and Corm Yield of *Gladiolus* cv. Malaviya Shatabdi

Anjana Sisodia, Devendra Kumar*, Anil K. Singh, Anurag Srivastava,
Sahil and Arun Kumar Maurya

Department of Horticulture, Institute of Agricultural Sciences,
Banaras Hindu University, Varanasi (Uttar Pradesh), India.

(Corresponding author: Devendra Kumar*)

(Received: 15 February 2025; Revised: 25 March 2025; Accepted: 16 April 2025; Published online: 09 May 2025)

(Published by Research Trend)

DOI: <https://doi.org/10.65041/BiologicalForum.2025.17.5.6>

ABSTRACT: An investigation was carried out to evaluate the effect of different levels of micronutrients on the growth, flowering and corm yield of gladiolus (*Gladiolus* spp.) cv. Malaviya Shatabdi. The experiment was conducted at Horticulture Research Farm, Department of Horticulture, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi, Uttar Pradesh, during 2023-24 season. Treatments include a control (distilled water) and varying levels of Fe and B, applied as foliar sprays at 3rd and 6th leaf stage. The results demonstrated that the combined application of 0.2% B + 0.4% Fe significantly improved plant growth, flowering characters and yield parameters. This treatment produced the highest number of leaves per plant (16.04), and maximum leaf width (2.73 cm), widest scape (2.54 cm) whereas, flowering parameters showed promising results at application of 0.2 % B + 0.4 % Fe for days to colour show, days to opening of 3rd and 5th floret, diameter of 3rd and 5th floret, length of 3rd and 5th floret, number of opened florets per spike (9.27) rachis length (44.18 cm) and internodal length (4.87). Similarly, corm production was also significantly influenced by Fe and B applications. The highest number of cormels per hill (9.60) and maximum weight of corm per hill (46.75 g). These findings suggest that foliar application of 0.2% B + 0.4% Fe optimizes vegetative growth, floral characteristics and corm yield in gladiolus cv. Malaviya Shatabdi.

Keywords: Foliar spray, gladiolus corms, micro nutrient, growth, flowering, corm yield.

INTRODUCTION

Gladiolus (*Gladiolus grandiflorus* L.), commonly referred to as "Glad" is a well-known bulbous cut flower plant that originated in South Africa. It is a member of the Iridaceae and sub-family Ixiodeae. Because of its sword-shaped leaves, gladiolus is also known as the Corn Lily or Sword Lily. Being a significant bulbous decorative plant, it holds a prominent place among commercial flower harvests and is much sought after in both domestic and foreign markets (Ahmad *et al.*, 2010). Iron acts as activation for several biochemical processes such as respiration, photosynthesis and symbiotic nitrogen fixation. These results validate the substantial role of iron accumulation of photo assimilates for growth and development of plants (Singh *et al.*, 2017). Whereas, boron plays a crucial role in ensuring normal plant growth (Shireen *et al.*, 2018). This micronutrient is required in trace amounts to support optimal growth, development and metabolic activities in plants (Gowthami *et al.*, 2022). In the case of cut flowers, boron is particularly important for enhancing shelf life, ensuring their longevity and marketability. Additionally, genetic variability exists among different plant genotypes,

leading to variations in growth patterns and adaptability (Swaroop *et al.*, 2019).

In the Varanasi region, the application of iron (Fe) and boron (B) in gladiolus cultivation is essential due to their critical roles in plant growth and flower quality. Iron is vital for chlorophyll synthesis, enhancing photosynthesis and improving flower yield, while boron supports cell wall formation, pollen germination and nutrient transport, leading to better spike and floret development. The alluvial soils of Varanasi often exhibit micronutrient deficiencies, particularly Fe and B, necessitating their supplementation to maximize gladiolus production (Singh *et al.*, 2017 and 2024).

MATERIALS AND METHODS

The field experiment was conducted during the winter season of 2023-24 at the Horticulture Research Farm, Department of Horticulture, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi, Uttar Pradesh, India, using a Randomized Block Design (RBD). Disease free corms of gladiolus cv. Malaviya Shatabdi were planted at a spacing of 30 cm × 20 cm at a depth of 8-10 cm in well prepared field incorporated with farmyard manure and NPK fertilizers applied as per recommendations. The study included eight

treatments with three replications. Iron (FeSO₄) and boron (boric acid) were applied *via* foliar spray at 3rd and 6th leaf stage with treatments comprising control (distilled water), 0.2% B, 0.4% B, 0.2% Fe, 0.4% Fe, 0.2% B + 0.2% Fe, 0.2% B + 0.4% Fe and 0.4% B + 0.4% Fe. All the growth flowering and corm parameters were recorded and analysed statistically.

RESULTS AND DISCUSSION

Growth parameters: Maximum number of leaves per hill (15.81), leaf width (2.73 cm) and scape width (2.54 cm) were recorded in plants treated with 0.2% B + 0.4% Fe treatment, which was statistically similar to the 0.2% B + 0.2% Fe, 0.4% B + 0.4% Fe and 0.4% Fe treatments but showed significant differences when compared to the other treatments. These findings align with the previous work of Soni and Godara (2015) in gerbera and Singh *et al.* (2012 and 2024) in gladiolus. Micronutrients play a crucial role in plant growth by facilitating key processes such as cell division, multiplication and differentiation, which contribute to enhanced photosynthesis and efficient nutrient transport. The observed increase in scape width can be attributed to these micronutrient-induced physiological improvements. Various growth parameters that have been influenced by different levels of iron and boron has been presented in Table 1. The vegetative attributes like number of leaves/hills, leaf width, scape width was significantly influenced due to the treatments. Increase in number of leaves may be due to transport of starch and sugars by boron affecting physiological processes, these findings supported to those of Singh *et al.* (2018 and 2024); Somkuwar *et al.* (2023) in gladiolus. Whereas, increase in leaf width and scape width with treatment 0.2 % B + 0.2% Fe is due to accumulation of boron and iron in leaf tissues causing cell elongation.

Also, cell wall synthesis and stability increase with components of cell wall like pectins, polyols and polyhydroxyl polymers.

Flowering Characters: Present study revealed that iron and boron treatments played a significant role in determining the various flowering parameters (Table 1 and 2; Fig. 1). Plants treated with 0.2% B + 0.4% Fe exhibited the earliest onset of flower colouration (89.93 days), likely due to the promotion of reproductive growth through optimal micronutrient application. These findings are in agreement with Singh *et al.* (2016 and 2024); Somkuwar *et al.* (2023) in gladiolus crop. The earliest floret opening with 0.2 % B + 0.4 % Fe treatment, may be attributed to iron's role in regulating plant hormone levels, promoting early maturation and floret development. Similar effects were observed in marigold by Kumar *et al.* (2010) following foliar iron application and in gladiolus by Memon *et al.* (2013). Whereas, increase in floret diameter and length is due to the reason that iron and boron play a vital role in promoting vegetative growth, which subsequently enhances flowering and may contribute to an increase in floret diameter. Similar results were reported by Bhandari *et al.* (2022) in calendula, Chopde *et al.* (2016) in annual chrysanthemum, Balkrishnan *et al.* (2007) in marigold and Verma *et al.* (2018) in China aster.

Iron and boron are being essential elements for plant growth, as they enhance carbohydrate accumulation by stimulating photosynthesis, which may contribute to floret elongation. These findings are in line with the research of Chopde *et al.* (2015) in tuberose. Additionally, research findings demonstrate that rachis length (43.60 cm) and internodal length (3.65 cm) were significantly influenced by different iron treatments.

Table 1: Effects of different concentrations of iron and boron on growth and flowering attributes in gladiolus cv. Malaviya Shatabdi.

Treatment	Number of leaves/hills	Leaf width (cm)	Scape width (cm)	Floret diameter (cm)		Floret length (cm)	
				3 rd floret	5 th floret	3 rd floret	5 th floret
Control (Distilled water)	12.55	2.07	1.90	6.96	6.70	8.70	8.40
0.2 % B	13.16	2.24	2.14	7.23	6.90	9.23	8.53
0.4 % B	13.54	2.27	2.19	7.16	7.13	9.33	9.00
0.2 % Fe	14.50	2.45	2.28	7.26	7.03	9.90	9.03
0.4 % Fe	14.97	2.60	2.37	8.03	7.70	9.93	9.23
0.2 % B + 0.2 % Fe	15.81	2.66	2.48	8.13	8.19	10.00	9.80
0.2 % B + 0.4 % Fe	16.04	2.73	2.54	8.23	8.20	10.13	10.00
0.4 % B + 0.4 % Fe	15.78	2.62	2.40	8.03	7.73	9.96	9.46
C.D. at 5%	1.59	0.23	0.20	0.30	0.46	0.44	0.42

Similar observations have been reported by Chopde *et al.* (2015); Somkuwar *et al.* (2023) in gladiolus.

A significant variation in opened florets per spike can be attributed to the essential role of iron in promoting the growth and development of gladiolus plants, along with its stimulatory and catalytic influence on flower yield and metabolic processes. These results align with the findings of Mishra *et al.* (2018); Kashyap and Tikey (2020) in gladiolus, Hajizadeh *et al.* (2019) in gerbera.

Corm and cormel parameters: The study demonstrated a significant impact on number and

weight of cormels per hill. The highest cormel count (9.60) and weight (46.75 g) were recorded in plants treated with 0.2 % B + 0.4 % Fe. This increase could be attributed to the application of iron and boron, which stimulated vegetative and reproductive growth, leading to higher cormel production. These results align with the findings of Kashyap and Tikey (2020); Yashawanth *et al.* (2016); Pratap *et al.* (2005); Somkuwar *et al.* (2023); Singh *et al.* (2024) in gladiolus.

The treatment also had a significant effect on corm weight per hill, that can be attributed to iron's role in

promoting cell enlargement and boosting vegetative growth, which facilitates nutrient transfer to the corms, ultimately increasing their weight. Similar results were

reported by Chen *et al.* (1982); Yashawanth *et al.* (2016) in gladiolus.




Fig. 1. Effect of iron and boron on colour show and opening of 3rd and 5th floret in gladiolus cv. Malaviya Shatabdi.

Table 2: Effects of different iron and boron on flowering and corm parameters in gladiolus cv. Malaviya Shatabdi.

Treatment	No. of open florets/spike	Rachis length (cm)	Inter nodal length (cm)	Number of cormels/hill	Weight of corms/hill (g)
Control (Distilled water)	7.88	38.00	2.66	4.30	39.50
0.2 % B	6.96	40.53	3.21	4.82	41.31
0.4 % B	8.24	41.23	3.38	5.53	42.28
0.2 % Fe	8.36	41.76	3.41	6.31	43.18
0.4 % Fe	8.40	42.60	3.25	7.48	44.26
0.2 % B + 0.2 % Fe	8.87	43.60	3.56	9.03	45.40
0.2 % B + 0.4 % Fe	9.21	42.16	3.65	9.60	46.75
0.4 % B + 0.4 % Fe	8.71	1.99	3.55	8.35	44.92
C.D. at 5%	1.11	37.90	0.38	2.54	2.46

CONCLUSIONS

This study highlights that the foliar application of 0.2% B + 0.4% Fe significantly enhances plant growth parameters, flowering characters and as well as corm and cormel yield in gladiolus cv. Malaviya Shatabdi. The results confirm that this treatment is the most effective for improving vegetative development and reproductive yield. Thus, it can be concluded that applying 0.2% B + 0.4% Fe is the optimal strategy for maximizing growth and yield in gladiolus cultivation.

FUTURE SCOPE

Evaluation of the long-term effects of repeated foliar treatments of Fe and B on soil health and gladiolus productivity can be the main focus of this research. Micronutrient recommendations could be standardized with the aid of comparative studies across various gladiolus cultivars and agroclimatic areas.

Acknowledgment. The authors express their sincere gratitude to the Institute of Agricultural Sciences, Banaras Hindu University (BHU), Varanasi, for providing the necessary facilities and resources for this study. Special thanks are also extended to the Department of Horticulture (Floriculture and Landscaping), BHU, for their valuable guidance and support throughout the research.

REFERENCES

- Ahmad, I., Aslam, K. M., Qasim, M., Ahmad, R. and Randhawa, A. M. (2010). Growth, yield and quality of *Rosa hybrida* as influenced by various micronutrients. *Pakistan Journal of Agriculture Science*, 47(1), 5-12.
- Balkrishnan, V. M., Jawaharlal, T., Kumar B. and Ganga, M. (2007). Response of micro nutrients on flowering, yield and xanthophyll content in African marigold. *Journal of Ornamental Horticulture*, 10(3), 153-156.
- Bhandari, N. S., Srivastava, R. K., Tarakeshwari, K. R. and Chand, S. (2022). Effect of nano and macro iron sprays on growth, flowering, seed and oil yielding attributes in calendula (*Calendula officinalis* L.). *Journal of Horticultural Sciences*, 17(2), 353-362.
- Chopde, N., Nehare, N., Maske, S. R., Lokhande, S. and Bhute, P. N. (2015). Effect of foliar application of zinc and iron growth, yield and quality of gladiolus. *Plant Archives*, 15(1), 417-419.
- Chopde, N., Borse, G. H., Kuchanwar, O. and Ghodke, A. T. (2016). Effect of zinc sulphate and ferrous sulphate on growth and flowering of annual chrysanthemum. *Plant Archives*, 16(2), 594-596.
- Gowthami, L., Bhaskar, V. V. and Padmaja, V. V. (2022). Effect of essential heavy metals on chlorophyll and carotenoid content in tuberose (*Polianthus tuberosa*). *Current Horticulture*, 10(2), 44-47.

- Hajizadeh, S., Jabbarzadeh, Z. and Rasouli-Sadaghiani, M. H. (2019). Effect of fulvic acid and iron nano chelate application on flowering and vase life of *Gerbera jamesonii* cv. Dune. *Journal of Horticulture Science*, 33(4), 711-725.
- Kashyap, N. and Tikey, T. (2020). Effect of micronutrients on plant growth, flowering and corm production of gladiolus cv. Summer Sunshine. *The Pharma Innovation Journal*, 11(9), 2503-2506.
- Kumar, P., Kumar, J., Umrao, V. K. and Rajbeer, R. (2010). Effect of nitrogen and iron on growth and flowering parameters in African marigold (*Tagetes erecta* L.) cv. Pusa Narangi Gainda. *Annual Horticulture*, 3(1), 118-119.
- Memon, S. A., Abdul, R. A., Muhammad, A. B. and Mahmooda, B. (2013). Effect of zinc sulphate and iron sulphate on the growth and flower production of gladiolus (*Gladiolus hortulanus*). *Journal of Agricultural Technology*, 9(6), 1621-1630.
- Mishra, A., Singh, A. and Kumar, A. (2018). Effect of foliar feeding of zinc and iron on flowering and yield attributes of gladiolus (*Gladiolus grandiflorus* L.) cv. Nova Lux *Plant Archives*, 18, 1355-1358.
- Pratap, M., Reddy, S. A. and Reddy, Y. N. (2005). Response of pre-harvest micronutrient foliar spray on leaf nutrients and corm production in gladiolus. *Journal of Ornamental Horticulture*, 8(1), 18-22.
- Shireen, F., Nawaz, M. A., Chen, C., Zhang, Q., Zheng, Z., Sohail, H. and Bie, Z. (2018). Boron: functions and approaches to enhance its availability in plants for sustainable agriculture. *International Journal of Molecular Sciences*, 19(7), 1856.
- Singh, A. K., Hamidi M., Sisodia A. and Barman, K. (2024). Effect of iron on growth, flowering and corm parameters in gladiolus cv. Malaviya Shatabdi. *International Journal of Advanced Biochemistry Research*, 8(7), 574-577.
- Singh, A. K., Hembrom, R., Sisodia, A., Pal, A. K. and Asmita, A. (2016). Effect of iron and zinc on flowering and postharvest life in gladiolus (*Gladiolus* spp.). *The Indian Journal of Agricultural Sciences*, 86(10), 1316-1319.
- Singh, A. K., Sisodia, A., Sisodia, V. and Kirti, A. (2018). Effect of foliar application of boron and zinc on growth and flowering characters in African marigold cv. Pusa Narangi Gainda. *Journal of Ornamental Horticulture*, 21(1), 1-6.
- Singh, A. K., Sisodia, A., Sisodia, V. and Ray, P. (2017). Performance of Indian and exotic varieties of gladiolus under Eastern UP conditions. *Journal of Ornamental Horticulture*, 20(3), 153-157.
- Singh, J. P., Kumar, K., Katiyar, P. N. and Kumar, V. (2012). Influence of zinc, iron and copper on growth and flowering attributes in gladiolus cv. Sapna. *Progressive Agriculture*, 12(1), 138-143.
- Somkuwar, A. R., Singh, A. K., Sisodia, A., Lamsal, A. and Giri, S. (2023). Effect of boron on growth and flowering in gladiolus (*Gladiolus* spp.). *Current Horticulture*, 11(2), 56-59.
- Soni, S. S. and Godara, A. K. (2015). Effect of foliar application of borax, FeSO₄ and MnSO₄ on vegetative growth and flower production in gerbera. *Research in Environment and Life Sciences*, 8(4), 581-584.
- Swaroop, K., Singh, K. P. and Kumar, P. (2019). Evaluation of gladiolus (*Gladiolus grandiflora*) genotypes for morphological diversity and corm yield. *Current Horticulture*, 7(2), 48-51.
- Verma, V. K., Verma, J. P., Verma, H. K. and Meena, R. K. (2018). Efficacy of micronutrients on growth and flower production of China aster [*Callistephus chinensis* (L) NEES] cv. Princess. *International Journal of Agricultural Sciences*, 14(1), 160-164.
- Yashawanth, D., Chandrashekar, S., Naik, B. H. and Ganapathi, M. (2016). Influence of micronutrients on corm and cormel production in gladiolus (*Gladiolus grandiflorus* L.) cv. Summer Sunshine. *Advances in Life Sciences*, 5(16), 6014-6017.

How to cite this article: Anjana Sisodia, Devendra Kumar, Anil K. Singh, Anurag Srivastava, Sahil and Arun Kumar Maurya (2025). Effect of Different Level of Micronutrients on Growth, Flowering and Corm Yield of Gladiolus cv. Malaviya Shatabdi. *Biological Forum*, 17(5): 37-40.