



Impact of Organic and Inorganic Nutrient Sources on Soil Physico-chemical properties and Productivity of (*Stevia rebaudiana* Bertoni)

Ekamjot Kaur¹, Saurabh Sharma^{1*}, Manpreet Singh¹, Arshdeep Singh¹, Manisha Negi² and Navneet Kaur¹

¹PG Department of Agriculture, Khalsa College Amritsar (Punjab), India.

²Department of Soil Science and Water Management,

Dr. YS Parmar University of Horticulture and Forestry, Nauni, Solan (Himachal Pradesh), India.

(Corresponding author: Saurabh Sharma*)

(Received: 20 March 2025; Revised: 26 April 2025; Accepted: 23 May 2025; Published online: 14 June 2025)

(Published by Research Trend)

DOI: <https://doi.org/10.65041/BiologicalForum.2025.17.6.9>

ABSTRACT: The present study evaluated the effects of integrated nutrient management (INM) strategies on soil physico-chemical properties and yield performance of *Stevia rebaudiana* Bertoni under field conditions at Khalsa College, Amritsar, Punjab, India, during the 2022–2023 cropping season. A total of twelve treatment combinations integrating varying proportions of inorganic fertilizers (RDF) with farmyard manure (FYM) and vermicompost (VC) were assessed in a randomized block design with three replications. Results revealed that the combined application of 100% RDF with 2 t ha⁻¹ VC (T₃) significantly enhanced fresh and dry leaf yield (18.65 q ha⁻¹ and 5.16 q ha⁻¹, respectively) compared to control and sole fertilizer treatments. Soil health indicators also improved markedly under INM, with T₃ and T₂ (100% RDF + 5 t ha⁻¹ FYM) treatments showing the highest gains in water holding capacity (41.29% and 41.65%, respectively), organic carbon content (0.44% and 0.47%), and available NPK. Integrated treatments notably reduced soil bulk density while improving porosity, pH balance, and macronutrient availability. These results suggest that INM approaches incorporating VC or FYM can sustainably improve soil quality and stevia productivity, offering a viable alternative to sole chemical fertilizer use in medicinal crop cultivation.

Keywords: Stevia, Organic, Inorganic, Soil Health, Integrated Nutrient Management.

INTRODUCTION

The rising demand for medicinal plants highlights the importance of sustainable cultivation methods, as relying solely on wild sources cannot fulfill this rising need. By cultivating these plants, we can ensure a consistent supply while also helping to preserve biodiversity (Kumar *et al.*, 2024). *Stevia rebaudiana* Bertoni, commonly known as "sweet herb of Paraguay," is a perennial herbaceous plant belonging to the Asteraceae family. Native to the semi-arid mountainous regions of northeastern Paraguay, it also occurs in adjacent areas of Brazil and Argentina. It is a perennial plant (Peteliuk *et al.*, 2021) and is now commonly used as a natural preference to synthetic sweeteners like aspartame, saccharin and acesulfame-K. Stevia is prized for its possible health advantages in addition to its ability to sweeten food. It has been used to treat various conditions, including cancer, obesity, hypertension, fatigue, and depression. Additionally, stevia has applications in cosmetics and dental care. Many researchers have confirmed that it is safe for use by children (Carrera-Lanestosa *et al.*, 2017). Stevia is now successfully cultivated in several Indian states, including Rajasthan, Maharashtra, Punjab, and Orissa

(Goyal *et al.*, 2010). It is believed that stevia can serve as a profitable diversification option in Punjab because of its agronomic and economic advantages over the current cropping cycle of paddy and wheat (Minhas *et al.*, 2023).

Stevia grows well in a semi-humid subtropical climate and is best suited to well-drained soils, especially red and sandy loam types, with a preferred pH range of 6.5 to 7.5 (Kumar and Simon 2024). Stevia's growth is influenced by both environmental factors and agricultural practices, particularly nutrient management, which can improve both growth and yield (Kumar *et al.*, 2024). While chemical fertilizers enhance plant growth and active compounds, their excessive and unbalanced use - without assessing soil health - can harm groundwater, disrupt soil biodiversity and reduce organic matter, threatening long-term sustainability (Calapardo and Manigo 2024). Additionally, the excessive use of inorganic fertilizers in modern farming-especially for medicinal crops-can result in harmful residues in medicinal products, posing a potential threat to human health and the environment. Therefore, it is crucial to transition towards safe and sustainable organic farming practices (Umesha *et al.*, 2011). Integrated nutrient management (INM), which

combines inorganic fertilizer with organic manure, is an approach designed to maximize crop yield while maintaining soil health (Kumar *et al.*, 2024). Research has demonstrated that optimal nutrient management during the cultivation of *Stevia rebaudiana* can significantly influence the plant's overall productivity (Sniegowska *et al.*, 2024).

INM combining fertilizers and organic manures, offers an effective approach to preserving soil health and rectifying crop productivity (Bajpai *et al.*, 2006). In addition to lowering the requirement for inorganic fertilizers, using crop residues, vermicompost (VC), farmyard manure (FYM), organic manures, and green manures combined with inorganic fertilizers enhances nutrient efficiency. This improvement occurs due to the positive impact of organic materials on the physical, chemical, and biological properties of the soil (Prasad *et al.*, 1992). VC contains vital nutrients such as N (in the form of nitrate or ammonium), P and K, along with a rich diversity of beneficial microorganisms. These elements help support plant health and enhance recovery from pesticide exposure and various environmental stresses (Calapardo and Manigo 2024). An integrated approach that combines organic manures with chemical fertilizers can significantly enhance soil fertility on a long-term basis. This system provides nutrients in a balanced manner, leading to improved nutrient uptake by crops (Hussainy *et al.*, 2019). This study examines how INM techniques alter the characteristics of the soil and *Stevia rebaudiana* productivity.

MATERIAL AND METHODS

A field trial was subjugated during the 2022–2023 growing season at the Student's Research Farm, Khalsa College, Amritsar, Punjab, India. The study employed a randomized block design (RBD) with three replications, encompassing twelve treatment combinations that integrated varying proportions of chemical fertilizers and organic manures *viz.*, T₁: 100 % RDF alone, T₂: 100 % RDF + 5 t ha⁻¹ FYM, T₃: 100 % RDF + 2 t ha⁻¹ VC, T₄: 50 % RDF + 5 t ha⁻¹ FYM, T₅: 50 % RDF + 2 t ha⁻¹ VC, T₆: 50 % RDF alone, T₇: 25 % RDF + 5 t ha⁻¹ FYM, T₈: 25 % RDF + 2 t ha⁻¹ VC, T₉: 25 % RDF alone, T₁₀: 5 t ha⁻¹ FYM alone, T₁₁: 2 t ha⁻¹ VC alone and T₁₂: absolute control. The experimental field soil had an alkaline pH of 8.3, moderate amounts of accessible P (14.06 kg ha⁻¹) and K (247.11 kg ha⁻¹), and low amounts of available N (177.27 kg ha⁻¹) and organic carbon (0.39 %). The soil texture was sandy loam. Farmyard manure was incorporated into the field one week prior to transplanting at the recommended rate of 25 tonnes per hectare. As per the recommendation of Pal *et al.* (2017), the fertilizer dose applied was 110:45:55 kg ha⁻¹ of N:P:K. At the time of transplantation, a half dose of N was administered as a basal dose together with the full amounts of K and P. Throughout the crop growth period, maximum temperatures ranged between 18.97°C and 38.44°C, while minimum temperatures fluctuated from 2.4°C to 28.05°C. Relative humidity during the season varied

between 50.42% and 95.57%. Post-harvest, observations were recorded to assess changes in available NPK, organic carbon, bulk and particle density, porosity, water holding capacity, electrical conductivity (EC), soil pH, micronutrient levels, and stevia crop yield. Soil samples from each plot were analyzed to assess various parameters indicative of soil fertility and characteristics. Available N was determined using the alkaline permanganate method, as described by Subbiah and Asija (1956). P content was estimated through the colorimetric technique outlined by Olsen *et al.* (1954). K levels were measured using flame photometry, following the procedure established by Merwin and Peech (1951). EC and soil pH were measured using a digital pH meter and conductivity meter respectively (Jackson, 1973), and organic carbon content was estimated via the Walkley and Black rapid titration method (Walkley and Black 1934). Bulk density was determined using the core method (Prihar and Hundal 1971), while particle density was measured with the PAU moisture gauge (Prihar and Sandhu 1968). Water holding capacity was assessed through the Keen box method (Keen and Raczkowski 1921). The data collected during this study were subjected to statistical analysis using Microsoft Excel, incorporating techniques from Exploratory Data Analysis (EDA) to examine and interpret the dataset's underlying patterns and relationships. Treatment effects on soil fertility parameters, physical properties, and yield were evaluated at a 5% significance level using the Randomized Block Design (RBD) for assessing differences among treatment means.

RESULTS AND DISCUSSION

A. Physical properties of soil

The data conferred in Table 1 demonstrate that the application of various organic and inorganic nutrient sources significantly influenced soil physical properties, including bulk density, porosity, and water holding capacity, at the time of stevia harvest. Among the different treatments, the highest bulk density (1.45 g cm⁻³) was observed in the control treatment (T₁₂), which was statistically comparable to T₁ (1.43 g cm⁻³), T₆ (1.43 g cm⁻³), and T₉ (1.44 g cm⁻³). Conversely, the lowest bulk density (1.35 g cm⁻³) was recorded in T₂, with values in T₃ (1.37 g cm⁻³), T₄ (1.36 g cm⁻³), T₇ (1.36 g cm⁻³), and T₁₀ (1.37 g cm⁻³) not differing significantly. The reduction in bulk density under integrated treatments, compared to the control and sole application of inorganic fertilizers, may be accredited to increased organic matter content resulting from the addition of organic manures. This likely promoted better soil aggregation and increased pore space, thereby improving soil physical properties. Similar results were also documented by Yadav *et al.* (2024). Soil porosity varied between 45.28 and 49.18 per cent across treatments. The highest porosity (49.18%) was recorded in treatment T₂ (100% RDF + 5 t ha⁻¹ FYM), which was statistically comparable to T₄ and T₇ (both 48.68%). The lowest porosity was observed in the unfertilized control (T₁₂) at 45.28%, which was

statistically similar to T₉ (45.79%). The improvement in porosity under integrated treatments may be attributed to the enhanced formation of soil aggregates resulting from the inclusion of organic amendments such as farmyard manure and VC. These organic inputs contribute to improved soil structure by reducing bulk density and enhancing aggregate stability, ultimately increasing pore space. Furthermore, FYM and vermicompost enhance soil porosity by improving structure and aggregation. This increased porosity boosts soil aeration, facilitating better root respiration and microbial activity, ultimately contributing to improved soil productivity (Yadav *et al.*, 2024).

Water holding capacity (WHC) of the soil also responded positively to integrated nutrient management practices. The highest WHC (41.65%) was observed in T₂ (100% RDF + 5 t ha⁻¹ FYM), which was statistically comparable to T₃ (41.29%) and T₄ (40.72%). The lowest WHC was recorded in T₁₂ (unfertilized control) at 32.35%, followed by T₉ (35.36%). The observed increase in WHC under INM treatments may be linked to improved root proliferation, greater porosity, enhanced aggregate formation, and reduced bulk density due to the incorporation of organic manures. Also, the higher water holding capacity observed under INM practices can be attributed to the application of organic manures, which enhance soil fertility, boost microbial biomass, support beneficial organisms, and significantly improve the soil's ability to retain moisture (Devi *et al.*, 2021).

B. Chemical properties of soil

The data presented in Table 2 showed the effect of INM on pH, EC and organic carbon of soil. The highest pH (8.30) was recorded under treatment T₉ (25% RDF alone) whereas lowest pH (8.23) recorded under T₂ (100% RDF + 5 t ha⁻¹ FYM). The degradation of organic manures likely leads to the formation of organic acids, contributing to the slight decrease in soil pH. The reduction in soil pH may be attributed to the release of organic acids such as humic and carbonic acids during the mineralization of organic manures. Similar findings were reported by Roy and Kashem (2014).

The highest EC value (0.39 dS m⁻¹) was observed in treatment T₁ where only recommended dose of fertilizers was applied and the lowest EC (0.35 dS m⁻¹) was observed in T₅ where 50% RDF + 2 t ha⁻¹ VC, T₁₀ where 5 t ha⁻¹ FYM and T₁₂ where no fertilizer was applied. Similar results were shown by Thite *et al.* (2023).

The soil organic carbon content (0.47%) was observed highest in treatment T₂, which received 100% RDF along with 5 t ha⁻¹ FYM, and it was statistically similar to treatments T₄ (0.46%), T₅ (0.45%), T₇ (0.46%), and T₁₀ (0.45%). A significantly lower organic carbon value of 0.36 per cent was found in the control treatment (T₁₂), which was statistically on par with T₆ (0.38%). The enhanced SOC content noticed with manure treatments may result from the direct addition and gradual decomposition of organic inputs within the soil. Also, the increase in soil organic carbon is largely due

to the direct input of organic matter from applied organic manures (Thite *et al.*, 2023).

C. Macro nutrient availability in soil

During the investigation period, treatments exhibited significant variations in the available NPK content of the soil, as detailed in Table 3. Post-harvest analysis of stevia revealed that soil available N ranged from 174.40 to 226.15 kg ha⁻¹. Among the integrated nutrient management practices, treatment T₃ (100% RDF + 2 t ha⁻¹ VC) recorded the highest available N level (226.15 kg ha⁻¹), which was statistically comparable to T₂ (223.06 kg ha⁻¹). Similarly, treatments T₁ (209.05 kg ha⁻¹), T₄ (210.05 kg ha⁻¹), and T₅ (211.81 kg ha⁻¹) showed no significant differences among them. Treatments T₆ (206.47 kg ha⁻¹), T₇ (205.98 kg ha⁻¹), T₈ (207.21 kg ha⁻¹), and T₁₁ (204.47 kg ha⁻¹) were also statistically at par. The lowest N availability was observed in T₁₂ (absolute control) at 174.40 kg ha⁻¹, followed by T₉ with 199.84 kg ha⁻¹. The increased N availability in treatments that received a combination of organic manures and chemical fertilizers may be attributed to both the mineralization of nutrients from the organic sources and the mobilization of native soil N. Similar results were reported by Bhanwaria and Yadav (2016).

The data indicated that an increase in soil available P levels at the time of stevia harvest compared to the initial soil status. Post-harvest available P concentrations ranged from 13.47 to 21.53 kg ha⁻¹ across different treatments. The highest available P content (21.53 kg ha⁻¹) was observed in treatment T₃, which was statistically comparable to treatments T₂ (20.65 kg ha⁻¹), T₄ (19.47 kg ha⁻¹), and T₅ (20.21 kg ha⁻¹). Conversely, the lowest available P (13.47 kg ha⁻¹) was ascertained in the control treatment (T₁₂), which was statistically similar to T₉ (15.61 kg ha⁻¹). The current findings suggest that the slower mineralization of FYM has led to a reduced availability of P in the soil. In contrast, plots treated with VC exhibited comparatively higher P availability, likely due to the accelerated mineralization process associated with VC application. The beheld rise in soil P content can be imputed to the combined effects of organic manures and inorganic fertilizers. Organic manures, such as farmyard manure, contribute P directly to the soil and release organic acids during decomposition. The solubility and availability of P in the soil can be increased by these organic acids' ability to chelate cations that fix P, such as calcium, iron, and aluminium. Additionally, it has been demonstrated that applying organic manures to inorganic fertilizers compliment the amount of P available in the soil. This improvement is ascribed to the increasing microbial activity and altering the pH of the soil, both of which promote the solubilization of inorganic P sources and the mineralization of organic P complexes. Our findings are in line with the findings of Thite *et al.* (2023).

The highest available K (293.78 kg ha⁻¹) was recorded in treatment T₃ (100% RDF + 2 t ha⁻¹ VC), which was statistically comparable to T₂ (291.63 kg ha⁻¹). The lowest available K content (245.07 kg ha⁻¹) was

observed in the control treatment (T₁₂). The increased availability of K under integrated nutrient management treatments may be associated to the interaction between organic matter and soil clay, which reduces K fixation and enhances K release. Additionally, the direct contribution of K from organic manures to the soil's available pool further explains the observed increase. Application of organic amendments thus contributed to higher available K content in the soil (Bhanwaria and Yadav 2016).

D. Yield parameters

Data on fresh leaf yield as influenced by the combined application of organic and inorganic fertilizers are presented in Table 4. The maximum fresh leaf yield (18.65 q ha⁻¹) was recorded under treatment T₃ (100 % RDF + 2 t ha⁻¹ VC), which was statistically at par with T₂ (100 % RDF + 5 t ha⁻¹ FYM), yielding 18.43 q ha⁻¹. The lowest fresh leaf yield (9.01 q ha⁻¹) was observed in the control treatment (T₁₂), which did not receive any organic or inorganic inputs and was statistically similar to T₁₀ and T₁₁. The increased leaf yield observed under integrated nutrient management treatments can be

attributed to the continuous and balanced supply of both macronutrients and micronutrients. This consistent nutrient availability supports robust plant growth, leading to enhanced leaf and branch development, and ultimately, a higher fresh yield per unit area.

The scrutiny of data transpires that application of 100 % RDF + 2 t ha⁻¹ VC (T₃) produced significantly highest dry yield of leaf (5.16 q ha⁻¹) which was statistically at par with T₁ (4.64 q ha⁻¹) and T₂ (4.96 q ha⁻¹) but significantly better than all other treatments. Minimum dry yield of leaf (1.84 q ha⁻¹) was observed in control treatment T₁₂ which was at par with T₁₀ and T₁₁ i.e. 2.30 and 2.55 q ha⁻¹ respectively. This can be attributed to the fact that VC and inorganic fertilizers increased the availability of nutrients in soil and hence we can see the better absorption and uptake by plant. Optimal maintenance of essential elements within the shoot system enhances canopy photosynthesis by increasing both the size of the functional leaf area and the assimilation rate per unit area. These findings are consistent with those reported by Negi *et al.* (2022); Sharma *et al.* (2022).

Table 1: Effect of integrated nutrient management on bulk density, porosity and water holding capacity of soil after harvest of stevia crop.

Symbol	Treatments	Bulk density (g cm ⁻³)	Soil Porosity (%)	Water holding capacity (%)
T ₁	100% RDF alone	1.43	46.16	37.37
T ₂	100% RDF + 5 t ha ⁻¹ FYM	1.35	49.18	41.65
T ₃	100% RDF + 2 t ha ⁻¹ VC	1.37	48.30	41.29
T ₄	50% RDF + 5 t ha ⁻¹ FYM	1.36	48.68	40.72
T ₅	50% RDF + 2 t ha ⁻¹ VC	1.38	47.92	40.45
T ₆	50% RDF alone	1.43	46.04	36.56
T ₇	25% RDF + 5 t ha ⁻¹ FYM	1.36	48.68	39.64
T ₈	25% RDF + 2 t ha ⁻¹ VC	1.39	47.55	39.38
T ₉	25% RDF alone	1.44	45.79	35.36
T ₁₀	5 t ha ⁻¹ FYM alone	1.37	48.43	38.56
T ₁₁	2 t ha ⁻¹ VC alone	1.39	47.67	38.35
T ₁₂	Control	1.45	45.28	32.35
LSD (p<0.05)		0.02	0.63	1.05

Table 2: Effect of integrated nutrient management on pH, EC and organic carbon of soil after harvest of stevia crop.

Symbol	Treatments	Soil pH	Soil EC (dS m ⁻¹)	Organic carbon (%)
T ₁	100% RDF alone	8.29	0.39	0.39
T ₂	100% RDF + 5 t ha ⁻¹ FYM	8.23	0.36	0.47
T ₃	100% RDF + 2 t ha ⁻¹ VC	8.25	0.37	0.44
T ₄	50% RDF + 5 t ha ⁻¹ FYM	8.26	0.36	0.46
T ₅	50% RDF + 2 t ha ⁻¹ VC	8.26	0.35	0.45
T ₆	50% RDF alone	8.28	0.38	0.38
T ₇	25% RDF + 5 t ha ⁻¹ FYM	8.25	0.36	0.46
T ₈	25% RDF + 2 t ha ⁻¹ VC	8.26	0.38	0.44
T ₉	25% RDF alone	8.30	0.38	0.39
T ₁₀	5 t ha ⁻¹ FYM alone	8.27	0.35	0.45
T ₁₁	2 t ha ⁻¹ VC alone	8.26	0.36	0.44
T ₁₂	Control	8.29	0.35	0.36
LSD (p<0.05)		NS	NS	0.02

Table 3: Effect of integrated nutrient management on macronutrient availability in soil after harvest of stevia crop.

Symbol	Treatments	Available macronutrient (kg ha ⁻¹)		
		N	P	K
T ₁	100% RDF alone	209.05	18.34	271.37
T ₂	100% RDF + 5 t ha ⁻¹ FYM	223.06	20.65	291.63
T ₃	100% RDF + 2 t ha ⁻¹ VC	226.15	21.53	293.78
T ₄	50% RDF + 5 t ha ⁻¹ FYM	210.05	19.47	280.64
T ₅	50% RDF + 2 t ha ⁻¹ VC	211.81	20.21	283.54
T ₆	50% RDF alone	206.47	17.73	262.87
T ₇	25% RDF + 5 t ha ⁻¹ FYM	205.98	17.57	269.88
T ₈	25% RDF + 2 t ha ⁻¹ VC	207.21	18.32	272.36
T ₉	25% RDF alone	199.84	15.61	258.74
T ₁₀	5 t ha ⁻¹ FYM alone	202.91	16.17	259.59
T ₁₁	2 t ha ⁻¹ VC alone	204.47	16.81	260.67
T ₁₂	Control	174.40	13.47	245.07
LSD (p≤0.05)		4.08	2.21	3.28
Initial status		177.27	14.06	247.11

Table 4: Effect of integrated nutrient management application on average leaf yield of stevia crop.

Symbol	Treatments	Average leaf yield (q ha ⁻¹)	
		Fresh	Dry
T ₁	100% RDF alone	16.04	4.64
T ₂	100% RDF + 5 t ha ⁻¹ FYM	18.43	4.96
T ₃	100% RDF + 2 t ha ⁻¹ VC	18.65	5.16
T ₄	50% RDF + 5 t ha ⁻¹ FYM	14.71	4.05
T ₅	50% RDF + 2 t ha ⁻¹ VC	14.99	4.24
T ₆	50% RDF alone	13.52	3.83
T ₇	25% RDF + 5 t ha ⁻¹ FYM	11.97	3.23
T ₈	25% RDF + 2 t ha ⁻¹ VC	12.70	3.75
T ₉	25% RDF alone	11.64	2.99
T ₁₀	5 t ha ⁻¹ FYM alone	10.54	2.30
T ₁₁	2 t ha ⁻¹ VC alone	10.66	2.55
T ₁₂	Control	9.01	1.84
LSD (p≤0.05)		2.51	0.76

CONCLUSIONS

The present study, titled “**Effect of Integrated Nutrient Management (INM) Approach on Soil Properties and Yield of Stevia (*Stevia rebaudiana* Bertoni)**,” underscores the efficacy of combining organic and inorganic fertilizers in improving soil physico-chemical properties and stevia crop performance. The integration of organic manures with chemical fertilizers not only ameliorated soil health but also significantly boosted the growth, yield, and quality of stevia. Among the various treatments, the application of 100% RDF + 5 t ha⁻¹ FYM notably enhanced soil physico-chemical attributes compared to both the control and sole RDF application. Furthermore, the consolidation of 100% RDF with 2 t ha⁻¹ of VC (T₃) markedly improved vegetative growth, yield, and quality parameters of stevia over RDF alone and the control. Based on overall performance metrics, treatment T₃ (100% RDF + 2 t ha⁻¹ VC) emerged as the most effective strategy for augmenting soil health, yield, and quality of stevia. These findings suggest that the continued application of VC in conjunction with recommended fertilizer doses could offer sustainable improvements in stevia cultivation under long-term field conditions.

FUTURE SCOPE

The present findings demonstrate that integrating organic and inorganic nutrient sources can significantly improve soil physico-chemical properties and enhance the productivity of *Stevia rebaudiana* Bertoni. Future research could explore long-term impacts of INM strategies on soil microbial dynamics, enzyme activity, and carbon sequestration under stevia-based cropping systems. Evaluating the economic feasibility and environmental sustainability of INM practices across diverse agro-ecological zones will also be crucial. Moreover, integrating biofertilizers and precision nutrient management tools may further enhance nutrient use efficiency and crop quality. This approach could be extended to other medicinal and cash crops, supporting holistic and sustainable agricultural practices that align with climate-resilient farming goals.

Conflict of Interest. None.

REFERENCES

- Calapardo, M. M. and Manigo, B. I. (2024) Enhancing herbage growth, yield and quality of stevia (*Stevia rebaudiana* Bertoni) using bio-organic nutrients in varied soil media. *Sains Malaysiana*, 53(3), 533-547.

- Bhanwaria, R. and Yadav, B. L. (2016). Effect of organic manure and soil moisture regimes on residual soil fertility and biological properties under mustard crop with saline water irrigation. *Research in Environmental and Life Sciences*, 9(9), 1135-1139.
- Carrera-Lanestosa, A., Moguel-Ordóñez, Y. and Segura-Campos, M. (2017). *Stevia rebaudiana* Bertoni: a natural alternative for treating diseases associated with metabolic syndrome. *Journal of Medicinal Food*, 20(10), 933-943.
- Devi, R. B., Singh, V., Tiwari, D., Sudhakar, M. and Rakesh, M. (2021). Influence of organic manures and molybdenum on growth and yield of groundnut (*Arachis hypogaea* L.). *Biological Forum – An International Journal*, 13(3), 402-406.
- Goyal, S. K., Samsher, N. and Goyal, R. (2010). *Stevia rebaudiana* a bio-sweetener: a review. *International journal of food sciences and nutrition* 61(1), 1-10.
- Hussainy, S. A. H., Sathiya, K. and Durairaj, N. (2019). Integration of different organic manures and nitrogenous fertilizer and its effect on the growth and yield of rice. *Journal of Pharmacognosy and Phytochemistry*, 8(2), 415-418.
- Jackson, M. L. (1973) *Soil Chemical Analysis*. Prentice Hall of India (Pvt.) Ltd, New Delhi.
- Keen, B. A. and Raczkowski, H. (1921). The relation between the clay content and certain physical properties of a soil. *Journal of Agriculture Science*, 11, 441-449.
- Kumar, P. M., Sindhu, P. V. and Beena, C (2024). Assessing integrated nutrient management strategies for improved growth rates and nutrient uptake in stevia (*Stevia rebaudiana* Bertoni). *International Journal of Plant and Soil Science*, 36(11), 632-640.
- Kumar, P. M., Sindhu, P. V., Prameela, P. and Savitha, A. (2024). growth rate and nutrient uptake of stevia rebaudiana bertoni as influenced by organic manures in laterite soils of Kerala, India. *Journal of Experimental Agriculture International*, 46(10), 442-449.
- Kumar, P. V. and Simon, S. (2024). Efficacy of soil amendments with organic manures, *Lantana camara* and microalgae against *Alternaria alternata* of *Stevia rebaudiana*. *Biological Forum – An International Journal*, 16(2), 42-46.
- Merwin, H. D and Peech, M. (1951). Exchange ability of soil potassium in the sand, silt and clay fraction as influenced by the nature and complementary exchangeable cations. *Soil Science American Proceedings*, 15, 125-128.
- Minhas, T. S., Kataria, P. and Kaur, A. (2023). *Stevia* cultivation in Punjab: Some crucial narratives. *Journal of Agricultural Development and Policy*, 33(1), 78-84.
- Negi, M., Sharma, J. C. and Shukla, Y. R. (2022). Effect of organic manures and inorganic fertilizers on dry matter yield, soil and plant nutrient content in onion (*Allium cepa*). *Indian Journal of Agricultural Sciences*, 92(6), 680-683.
- Olsen, S., Cole, C., Watanabe, F. and Dean, L. (1954) Estimation of available phosphorus in soils by extraction with sodium bicarbonate. *United States Department of Agriculture Circular No. 939*.
- Pal, P. K., Kumar, A., Sanatasujat, S., Kumar, S., Shamra, M. and Sud, R. K. (2017). Agrotechnology of stevia (*Stevia rebaudiana* B.). Technical folder Published by CSIR-Institute of Himalayan Bioresource Technology, Palampur (Himachal Pradesh), India.
- Peteliuk, V., Rybchuk, L., Bayliak, M., Storey, K. B. and Lushchak, O. (2021). Natural sweetener *Stevia rebaudiana*: Functionalities, health benefits and potential risks. *EXCLI journal*, 20, 1412.
- Prasad, R., Sharma, S. N., Singh, S. and Lakshmanan, R. (1992). Agronomic practices for increasing nitrogen use efficiency and sustained crop production. In: *National Symposium for Resource Management for Sustained Production, Indian society of Agronomy, Rajasthan Agricultural University*, 1(1), 25-28.
- Prihar, S. S. and Hundal, S. S. (1971). Determination of bulk density of soil clods by saturation. *Geoderma*, 5(4), 283-286.
- Prihar, S. S and Sandhu, B. S. (1968). A rapid method for soil moisture determination. *Soil Science*, 105, 142-144.
- Roy, S. and Kashem, M. A. (2014). Effect of organic manures in changes of some soil properties at different incubation periods. *Open Journal of Soil Science*, 4, 81-86.
- Sharma, S., Sharma, J. C., Shukla, Y. R., Negi, M. and Sharma, K. (2022). Assessment of integrated application of organic and inorganic nutrient sources for soil quality and economics of cucumber (*Cucumis sativus* L.) production. *Communications in Soil and Plant Analysis*, 53(18), 2476-2483.
- Sniegowska, J., Biesiada, A. and Gasinski, A. (2024). Influence of the nitrogen fertilization on the yield, biometric characteristics and chemical composition of *Stevia rebaudiana* Bertoni grown in Poland. *Molecules*, 29(8), 1865-1870.
- Subbiah, B. V. and Asija, G. L. (1956). A rapid procedure for the determination of available nitrogen in soils. *Current Science*, 25, 259-260.
- Thite, M. D., Ranshur, N. J. and Amolik, S. M. (2023). Effect of Organics on Physico-chemical properties of Inceptisol, *Biological Forum – An International Journal*, 15(7), 307-313.
- Umesha, K., Smitha, G. R., Sreeramu, B. S. and Waman, A. A. (2011). Organic manures and bio-fertilizers effectively improve yield and quality of stevia (*Stevia rebaudiana*). *Journal of Applied Horticulture*, 13(2), 157-162.
- Walkley, A. and Black, I. A. (1934). An examination of the different method for determining soil organic matter, and a proposed modification of the chromic acid titration method. *Soil Science*, 37, 29-38.
- Yadav, O., Singh, S. S., Patel, V. K., Singh, V. K., Mishra, S. and Jaiswal, R. (2024). Impact of chemical fertilizers, organic manure and bio-inoculants on sustainable production of onion (*Allium cepa* L.). *Biological Forum – An International Journal*, 16(1), 31-36.

How to cite this article: Ekamjot Kaur, Saurabh Sharma, Manpreet Singh, Arshdeep Singh, Manisha Negi and Navneet Kaur (2025). Impact of Organic and Inorganic Nutrient Sources on Soil Physico-chemical properties and Productivity of (*Stevia rebaudiana* Bertoni). *Biological Forum*, 17(6): 53-58.