

## Effect of Weather Parameters on Seasonal Activities of *Plutella xylostella* (L.) on Cabbage

Vinod Kumar\*, Akhter Hussain, S.L. Sharma, D.R. Bajya, Manisha Sharma,  
J.K. Bana and Laxman Singh Saini

Department of Entomology,  
Sri Karan Narendra Agriculture University, Jobner (Rajasthan), India.

(Corresponding author: Vinod Kumar\*)

(Received: 03 October 2024; Revised: 04 November 2024; Accepted: 14 November 2024; Published online: 20 January 2025)  
(Published by Research Trend)

**ABSTRACT:** The field experiment was conducted at Horticulture Farm, S.K.N. College of Agriculture, Jobner, Rajasthan during two consecutive seasons i.e. Rabi, 2022-23 and Rabi, 2023-24. During investigation diamondback moth was reported as a major insect pest of cabbage crop. The infestation of diamondback moth was initiated from third week of December (51<sup>st</sup> SMW) and peaked in first week of February (6<sup>th</sup> SMW) and second week of February (7<sup>th</sup> SMW) in Rabi, 2022-23 and 2023-24 respectively. The correlation studies revealed that the *P. xylostella* population showed a significant positive correlation with maximum temperature ( $r = 0.57$ ) and sunshine hours ( $r = 0.56$ ) during Rabi, 2022-23. Whereas during 2023-24, it had a significant positive correlation with minimum temperature ( $r = 0.73$ ).

**Keywords:** Diamondback moth, seasonal incidence, correlation, abiotic factors.

### INTRODUCTION

Cabbage [*Brassica oleracea* var. *capitata* (Linn.): Cruciferae] is one of most abundantly consumed cruciferous vegetable all over the world (Matharu and Mehta 2017). It is an important cruciferous vegetable crop which growing mainly in Rabi season in India. It is cultivated extensively in the tropical and temperate regions of the world including India. It is one of the most popular and widely grown vegetable and has occupied second position in production after potato. It is a rich source of vitamin A, C and mineral including potassium, calcium, sodium and iron. It is good sources of protein (1.3 per cent) which contains all essential amino acids, particularly sulphur containing amino acids. Cabbage is commonly used as cooked vegetable. It also makes into a good salad mixed in tomato, green chillies, beet root, etc. Cabbage is used as boiled vegetable, salad, curries and pickles. The total area under cabbage cultivation in India is 432 thousand ha with an annual production of 10048.6 thousand metric tonnes, while in Rajasthan, the total area and annual production is 0.75 thousand ha and 7.48 thousand metric tonnes, respectively, (Anonymous, 2023). Cabbage crop is attacked by a number of insects viz., tobacco caterpillar, *Spodoptera litura* (F.); diamondback moth, *Plutella xylostella* (L.); cabbage leaf webber, *Crociodolomia bionotalis* (Zell); aphid, *Lipaphis erysimi* (Kalt); painted bug, *Bagrada cruciferarum* (Kirk.) and flea beetle, (Goeze) Rao and Lal (2005) from sowing to harvest. Among the pest complex of cabbage, diamond back moth (*Plutella xylostella* Linn.) is the most destructive of cultivated

cruciferous crops worldwide, Talekar and Shelton (1993). Kumar *et al.* (1983) reported 52 percent losses in marketable yield of cabbage due to the infestation of *Plutella xylostella*. Diamondback moth (DBM), *Plutella xylostella* (Lepidoptera: Plutellidae) is a serious pest and has a great economic importance worldwide. It was estimated that at least 53–80% loss in marketable yield is due to the DBM attack alone and loss could be more if the attack is severe. It has national importance on cabbage because it causes 50-80 per cent annual loss in the marketable yield. Krishnamoorthy (2004) reported 52 per cent losses in yield due to the attack of only diamondback moth. The knowledge of seasonal incidence of insect pests at different growth stages of cabbage crop will be helpful in evolving proper management schedule. The knowledge of seasonal incidence of insect pests at different growth stages of cabbage crop will be helpful in evolving proper management schedule. The information on seasonal incidence was however, generated by many workers (Sachan and Srivastava 1972; Shukla and Kumar 2004; Wagle *et al.*, 2005) from different regions of India. Hence, investigations on seasonal incidence of major pests of cabbage and their natural enemies in relation to weather parameters was undertaken and the results presented herein

### MATERIALS AND METHODS

Cabbage cv. Golden Acre was transplanting in 60 × 45 cm spacing, divided into five plots each measuring 3 × 2.25 m<sup>2</sup>, at Horticulture Farm “S.K.N. College of Agriculture, Jobner (SKNAU, Jobner) Rajasthan, India during two consecutive years i.e., Rabi, 2022-23 and

*Rabi*, 2023-24. The seed were sown on the seed beds on the marked lines and they were covered with the fine soil. The seed rate utilized was 600 g/ha (*i.e.*, seedlings required for transplanting one hectare area field. No insecticidal treatment was applied at any stage of the crop growth. Observations on the larval population DBM was conducted on 25 randomly selected cabbage plants in an experimental field. These observations were carried out weekly, starting from the appearance of pests and continuing until the harvest of the crops. To estimate the larval population of diamondback moth, direct visual counting method was used (Lal, 1998). The observations were recorded at weekly interval throughout the crop growth on five randomly selected plants from each plot. The metrological data regarding weather parameters were collected from Metrological Observatory, Department of Agronomy (SKNAU, Jobner). The observations taken on the incidence of the pest of diamondback moth and there was correlated with weather parameters, *viz.*, maximum and minimum temperature, relative humidity (morning and evening) and sunshine hours.

## RESULTS AND DISCUSSION

**Seasonal incidence of diamondback moth at different growth stages of the crop.** The data presented in Table 1 and 2 indicate that the incidence of Diamond back moth, *P. xylostella* commenced in 3<sup>rd</sup> week of December (51<sup>th</sup> SMW) during both the years, *i.e.*, *Rabi*, 2022-23 and 2023-24 with 1.24 and 1.20 larvae per plant, respectively. The incidence continued thereafter for a long period and reached to peak first week of February (6<sup>th</sup> SMW) with 4.44 and second week of February (7<sup>th</sup> SMW) with 4.84 larvae per plant during 2022-23 and 2023-24 respectively. The weather parameters indicated that the infestation of Diamond back moth on cabbage crop was started at maximum temperatures (25.5°C and 25 °C), minimum temperatures (4.6°C and 5.5°C), morning relative humidity (79 % and 68 %) evening relative humidity (37 % and 34 %) and sunshine hours (8.1 and 7 hrs. per day) during *Rabi*, 2022-23 and 2023-24, respectively. The maximum larval population of diamond back moth was recorded in first (5<sup>th</sup> SMW) and second (6<sup>th</sup> SMW) week of February (4.44 and 4.84 larvae per plant) at maximum temperature (26.9°C and 26.6°C) minimum temperatures (6.90°C and 7.2°C), morning relative humidity (82 % and 72 %), evening relative humidity (31 % and 29 %) and sunshine hours (9 and 7.2 hrs/day) during *Rabi*, 2023-24, respectively. After reaching the peak, the population of diamond back moth started to decline and reached to low level in the 3<sup>rd</sup> week of March during both the years. The present results are in agreement with those of Jat *et al.* (2017a); Pallavi *et al.* (2023) who reported that the infestation of diamond back moth, *P. xylostella* was started in 2<sup>nd</sup> week of December during both the years and reached to its peak in the 5<sup>th</sup> and 6<sup>th</sup> SMW during both the years,

respectively. The results are in agreement with those of Jat *et al.* (2017b), also reported that the population of diamond back moth peak during second week of February. Misra *et al.* (2017) ; Anitha & Kalasariya (2024) observed the incidence of *P. xylostella* was started in the 4<sup>th</sup> week of December (52<sup>nd</sup> SMW) and reached its peak in 3<sup>rd</sup> to 4<sup>th</sup> week of February (7<sup>th</sup> SMW) are in conformity of present finding. Similarly, Dhaked and Kumar (2019) reported that the diamond back moth was observed 3<sup>rd</sup> week of December (51<sup>th</sup> SMW) and 4<sup>th</sup> week of December (52<sup>nd</sup> SMW) during both the years, respectively and it's reached to peak level in 2<sup>nd</sup> week of February (7<sup>th</sup> SMW) and 3<sup>rd</sup> week of February (8<sup>th</sup> SMW) during both the years, respectively. The present findings are in similar relation with the findings of Patra *et al.* (2013); Saranya *et al.* (2017); Bhagat *et al.* (2018); Lal *et al.* (2020), reported that DBM initially appeared in the December and attained its peak in the different week of February.

### Correlation study between population of diamondback moth and abiotic factors on cabbage.

The larval population of *P. xylostella* showed positive significant correlation with maximum temperature ( $r=0.57$ ) and sunshine hours ( $r=0.56$ ) and non-significant correlation with minimum temperature ( $r=0.46$ ) whereas, non-significant negative correlation morning and evening relative humidity ( $r=-0.33$ ,  $-0.41$ ) during *Rabi*, 2022-23, whereas in 2023-24, it had significant positive correlation with minimum temperature ( $r=0.73$ ) while, non-significant positive correlation with maximum temperature ( $r=0.17$ ) and morning relative humidity ( $r=0.04$ ) and sunshine hours ( $r=0.20$ ) while, evening relative humidity showed non-significant negative correlation ( $r=-0.15$ ). These findings are in close agreement with those of Venkateswarlu *et al.* (2011); Mahendran *et al.* (2017) who, observed a significant positive correlation of diamondback moth population with maximum temperature, minimum temperature and bright sunshine hours. Similarly, Jat *et al.* (2017b). The diamond back moth exhibited a positive significant correlation with temperature and negative non-significant correlation with humidity. The present findings are conformity with those of Sultana *et al.* (2019) reported that positive correlations between maximum temperature. While Gopika *et al.* (2022) reported that the diamondback moth had highly significant positive correlation with minimum temperature. Bhagat *et al.* (2018) revealed that there was no statistically significant correlation between DBM populations and various weather parameters, including maximum and minimum temperatures, relative humidity and sunshine hours. The present findings are also supported by Jat *et al.* (2017c); Anitha & Kalasariya (2024); Sharma *et al.* (2024), who reported a significant positive correlation between the diamondback moth population and bright sunshine hours.

**Table 1: Seasonal incidence of diamondback moth, *Plutella xylostella* (L.) in relation to abiotic factors on cabbage Rabi, 2022-23.**


Sr. No.	SMW*	Mean larval population of DBM/ plant	Temperature (°C)		Relative Humidity (%)		Bright Sunshine (hrs/ day)
			Max	Min.	Mor.	Eve.	
1.	51	1.24	25.5	4.6	79	37	8.1
2.	52	1.40	23.3	2.2	79	30	1.9
3.	1	1.48	20.7	-1	82	40	7.5
4.	2	1.96	24.5	4.9	75	35	7.3
5.	3	2.12	20.1	-0.5	77	33	8.3
6.	4	2.24	21.3	3.6	80	42	4.1
7.	5	3.20	22	5.2	83	54	7.3
8.	6	4.44	26.9	6.9	82	31	9
9.	7	4.24	28.4	5.8	76	26	9.4
10.	8	4.00	31.9	9.8	73	25	8.6
11.	9	3.80	32.1	12	70	28	8.8
12.	10	3.40	30.7	11.3	77	28	6.9
13.	11	2.80	32.8	13.8	79	32	6.7
14.	12	1.88	28.5	13.1	81	42	6.3
Correlation coefficient (r)			0.57*	0.46(NS)	-0.33(NS)	-0.41(NS)	0.56*

SMW- Standard metrological week; \*Significant at 5% level and NS- Non significant


**Table 2: Seasonal incidence of diamondback moth, *Plutella xylostella* (L.) in relation to abiotic factors on cabbage Rabi, 2023-24.**

Sr. No.	SMW	Mean larval population of DBM/ plant	Temperature (°C)		Relative Humidity (%)		Bright Sunshine (hrs/ day)
			Max	Min.	Mor.	Eve.	
1.	51	1.20	25	5.5	68	34	7
2.	52	1.60	25	4.9	79	42	6.4
3.	1	1.68	17	3.5	79	58	2.7
4.	2	1.88	21.9	3.3	76	37	5.3
5.	3	2.20	23.2	3.5	75	35	6.5
6.	4	2.44	24	4.3	73	32	7.1
7.	5	4.60	26	11.1	79	35	5.5
8.	6	4.72	22.9	6.6	73	33	7.8
9.	7	4.84	26.6	7.2	72	29	7.2
10.	8	4.44	28	10.4	73	34	8
11.	9	3.80	26.4	11.6	70	42	6.26
12.	10	3.60	28.3	8.9	68	52	8.21
13.	11	2.60	30.6	7.6	74	29	8.38
14.	12	1.80	34.07	13.0	61.57	24.85	9.68
Correlation coefficient (r)			0.17 (NS)	0.73*	0.04 (NS)	-0.15(NS)	0.20

SMW- Standard metrological week; \*Significant at 5% level and NS- Non significant



**Fig. 1.** Seasonal incidence of diamondback moth, *Plutella xylostella* (L.) in relation to abiotic factors on cabbage Rabi, 2022-23.



**Fig. 2.** Seasonal incidence of diamondback moth, *Plutella xylostella* (L.) in relation to abiotic factors on cabbage Rabi, 2023-24.

## CONCLUSIONS

The present investigation made on seasonal incidence of diamondback moth in relation to abiotic factors on cabbage. The infestation of diamondback moth 3<sup>rd</sup> week of December (51<sup>th</sup> SMW) during both years. The population of diamondback moth gradually increased and reached to its peak in the first week of February (6<sup>th</sup> SMW) and second week of February (7<sup>th</sup> SMW) during 2022-23 and 2023-24 respectively. The population of diamondback moth was positively correlated with maximum temperature and sunshine in 2022-23 and it had also showed positively correlated with minimum temperature during 2023-24.

## FUTURE SCOPE

According to the present findings of the investigation, there is a plenty of scope to build the studies that can enhance and implement proper effective management practices for the control of insect pests. This research would aid in the implementation of natural enemies and pest IPM tactics.

## REFERENCES

- Anitha, V. and Kalasariya, R. (2024). Population Dynamics of Major Lepidopteran Pests on Cauliflower. *Indian Journal of Entomology*, 86(2), 430-433.
- Anonymous (2023). Department of Agriculture, Ministry of Agriculture and Farmers Welfare, Government of India.
- Bhagat, P., Yadu, Y. K. and Sharma, G. L. (2018). Seasonal incidence and effect of abiotic factors on population dynamics of diamondback moth (*Plutella xylostella* L.) on cabbage (*Brassica oleracea* var. *Capitata* L.) crop. *Journal of Entomology and Zoology Studies*, 6(2), 2001-2003.
- Dhaked, N. S. and Kumar, A. (2019). Population dynamics of diamond back moth (*Plutella xylostella* L.) on cabbage under agro-climatic conditions of Allahabad (UP), India. *Journal of Experimental Zoology*, 22(2) 801-804.
- Gopika, R., Sonkamble, M. M. and Lad, A. G. (2022). Seasonal incidence of major insect pests of cabbage and their correlation with weather parameters. *Journal of Entomological Research*, 46(1), 154-157.
- Jat, M. K., Rana, B. S. and Jain, A. K. (2017a). Seasonal incidence of major insect pests of cabbage with their natural enemies. *Indian Journal of Entomology*, 79(3), 257-260.
- Jat, M. K., Rana, B. S. and Jain, A. K. (2017b). Seasonal incidence of major insect pest of cabbage with their natural enemies. *Indian Journal of Entomology*, 79(3), 257-260.
- Jat, G. C., Swaminathan, R., Yadav, P. C. and Lekha (2017c). Diversity of pest complex and their associated natural enemies in cabbage. *Plant Archives*, 17(1), 655-658.
- Krishnamoorthy, A. (2004). Biological control of diamondback moth *Plutella xylostella* (L.), an Indian scenario with reference to past and future strategies. In *Proceedings of the International Symposium* (Kirk, A. A and Bordat, D. eds.), Montpellier, France, CIRAD (pp. 204-211).
- Kumar, K. H. K., Srinivasan, K. and Suman, C. L. (1983). Optimum control strategy of cabbage pests from a chemical control trial. *Progressive Horticulture*, 18, 104-110.
- Lal, J., Swaminathan, R., Meena, A. K. and Nagar, R. (2020). Seasonal incidence of major insect pests of cabbage, *Brassica oleracea* var. *capitata* L., *Journal of Entomology and Zoology Studies*, 8(3), 387-391.
- Patra, S., Dhote, V. W., Alam, S. K. F., Das, B. C., Chatterjee, M. L. and Samanta, A. (2013). Population dynamics of major insect pests and their natural enemies on cabbage under new alluvial zone of West Bengal. *Journal of Plant Protection Sciences*, 5(1), 42-49.
- Mahendran, B., Sharma, R. K. and Sinha, S. R. (2017). Population dynamics of natural enemies of *Plutella xylostella* and *Pieris brassicae* on cauliflower. *Indian Journal of Entomology*, 79(4), 411-416.
- Lal, O. P. (1998). Notes summer school on "Advance Technologies in Important Vegetable Crops, including Cole Crops". May 4-24, I.A.R.I. New Delhi, pp. 63-66
- Matharu, K. S. and Mehta, P. K. (2017). Field efficacy of indigenous plant extracts against diamondback moth *Plutella xylostella* (Lepidoptera: Plutellidae). *Environmental Ecology*, 35(4), 3417- 3421.
- Misra, A., Sarkar, N., Gope, A., Samanta, S. and Banerjee, S. (2023). Seasonal activities of diamond back moth (*Plutella xylostella* L.) in cabbage (*Brassica oleracea* var. *capitata*) and their relation with the various environmental factors under the gangetic plains of

- West Bengal. *Environment and Ecology*, 41(2), 1199-1203.
- Pallavi, V., Dhurgude, S. S., Patait, D. D. and Kokkula A. (2023). Seasonal Incidence of Major Insect Pests of Cabbage (*Brassica oleracea* var. *capitata* L.) at Parbhani, Maharashtra. *Biological Forum – An International Journal*, 15(10), 1615-1622.
- Rao, S. R. K. and Lal, O. P. (2005). Seasonal incidence of mustard aphid, *Lipaphis erysimi* (Kalt.) and diamondback moth, *Plutella xylostella* Linn., on cabbage. *Journal of Insect Science*, 18(2), 106.
- Sachan, J. N. and Srivastava, B. P (1972). Studies of the seasonal incidence of insect pests of cabbage. *Indian Journal of Entomology*, 34(2), 123-129.
- Saranya, V. S. L., Rana, B. S. and Murdia, A. (2017). Seasonal incidence of major insect pests of cauliflower. *Indian Journal of Applied Entomology*, 31(2), 85–89.
- Sharma, K., Mahla, M., Ahir, K. and Upadhyay, B. (2024). Effect of weather parameters on the incidence of diamondback moth, *Plutella xylostella* (L.) in cabbage. *Pest Management in Horticultural Ecosystems*, 30(1), 49-55.
- Shukla, A. and Kumar, A. (2004). Seasonal incidence of diamondback moth, *Plutella xylostella* (Linn.) in relation to abiotic factors in cabbage. *Plant Protection Bulletin*, 56(3), 37- 38.
- Sultana, M. S., Khatun, M. F., Alam, S. N. and Miah, M. R. U. (2019). Population abundance of leaf eating caterpillars of cabbage. *Bangladesh Journal of Agricultural Research*, 44(1), 79-87.
- Talekar, N. S. and Shelton, A. M. (1993). Biology, ecology, and management of the diamondback moth. *Annual review of entomology*, 38(1), 275-301.
- Venkateswarlu, V., Sharma, R. K., Chander, S., and Singh, S. D. (2011). Population dynamics of major insect pests and their natural enemies in cabbage. *Annals of Plant Protection Sciences*, 19(2), 272-277.
- Wagle, B. K. S., Saravanan, L., Sudha, J. P. and Gupta, P. (2005). Seasonal incidence of cabbage aphid, *Brevicoryne brassicae* (L.) and its natural enemies in relation to weather parameters on cabbage. *National Conference on Applied Entomology*, RCA, Udaipur, 26-28 September, pp. 22-23.

**How to cite this article:** Vinod Kumar, Akhter Hussain, S.L. Sharma, D.R. Bajya, Manisha Sharma, J.K. Bana and Laxman Singh Saini (2025). Effect of Weather Parameters on Seasonal Activities of *Plutella xylostella* (L.) on Cabbage. *Biological Forum – An International Journal*, 17(1): 131-135.