



Genetic Variability Studies in Castor Germplasm (*Ricinus communis* L.)

K. Chandana^{1*}, A.V.S. Durga Prasad², N. Sabitha¹ and M. Raghavendra³

¹Department of Genetics & Plant Breeding, S.V. Agricultural College, Tirupati (Andhra Pradesh), India.

²Department of Genetics & Plant Breeding, SMGR Agricultural College, Udayagiri (Andhra Pradesh), India.

³Department of Biochemistry, S.V. Agricultural College, Tirupati (Andhra Pradesh), India.

(Corresponding author: K. Chandana*)

(Received: 08 May 2025; Revised: 19 June 2025; Accepted: 16 July 2025; Published online: 06 August 2025)

(Published by Research Trend)

ABSTRACT: Forty diverse genotypes of castor (*Ricinus communis* L.) were evaluated during kharif, 2024 to estimate genetic variability, broad sense heritability, and genetic advance for 12 metric traits. The analysis of variance revealed significant amount of variation among the genotypes for all the traits studied. It was noted that the phenotypic coefficient of variation (PCV) was always higher than the genotypic coefficient of variation (GCV), indicating role of environment in governing the traits. Notably, traits like number of capsules on primary spike and total seed yield have high PCV and GCV implying the wide spectrum of variability for these traits. High heritability coupled with high genetic advance as a percentage of the mean was recorded for effective primary spike length, number of capsules on primary spike, number of effective spikes per plant, 100 seed weight and total seed yield. This indicates a predominant additive gene action in the expression of these traits, suggesting that direct selection for them would be rewarding towards genetic improvement of castor.

Keywords: PCV, GCV, Genetic advance, Heritability.

INTRODUCTION

Castor (*Ricinus communis* L., $2n=2x=20$), a premier non-edible commercial oilseed crop, belongs to spurge family, commonly known as Euphorbiaceae. India is the global supplier of castor oil and its by-products. India alone produces 18.79 LMT of castor bean from 10.30 lakh ha cultivated area with an average productivity of 1824 kg ha⁻¹ (INDIASTAT, 2023-24). In India, castor is chiefly grown in the states

Of Gujarat followed by Rajasthan, Andhra Pradesh, Telangana, Tamil Nadu, Karnataka and Madhya Pradesh. Presently castor crop is cultivated in 30 countries across the globe among which India is leading in production and export (Sadaiah *et al.*, 2021). India exports 83 per cent of world's castor oil to China, USA, Japan, Thailand and few European countries (Lokesh *et al.*, 2020). Castor is a sexually polymorphic species With

different sex forms viz., monoecious, pistillate, hermaphrodite and pistillate with interspersed staminate flowers (Delvadiya *et al.*, 2021; Reddy *et al.*, 2022). Castor seeds contain 45 to 55 per cent oil which is the major source of the unusual hydroxyl fatty acid 'ricinoleic acid' that has tremendous industrial significance (Senthilvel *et al.*, 2016). Castor oil contains substantial quantity of the unusual hydroxylated fatty acid "ricinolate" which enhances its consumption as a lubricant in power engines. It can

readily be dissolved in alcohol and transformed into biodiesel even at low temperatures. It is also used to manufacture soaps, printing inks, linoleum, varnishes and plasticizers. Castor oil is a potent laxative and is also a curing agent for skin problems viz., sunburn, crinkles and stretch symbols, etc. The by-product of castor oil expeller is the cake which contains vital organic nitrogen (6.4%), P₂O₅ (2.5%) and K₂O (1%) including micronutrients for organic farming. The plant stems are used as firewood and to prepare paper pulp by the paper mills. Beyond this, fresh castor leaves are used to rear eri-silkworms, while desiccated leaves are used as an insecticide in agriculture.

The success of plant breeding hinges on the existing genetic variability within the crop (Zheng *et al.*, 2010). Assessing and estimating genetic variability in the germplasm is crucial before initiating any crop improvement program or selecting appropriate breeding techniques. This analysis aids in developing high-yielding and high-quality cultivars, ultimately increasing production. Heritability plays a key role in determining the transmissibility of traits to future generations, which is vital for selecting component traits to improve yield. Heritability estimates, along with genetic advances, are more reliable in predicting the genetic gain achieved via selection compared to heritability estimates alone.

MATERIALS AND METHODS

The present investigation was carried out during *kharif*, 2024 at Agricultural Research Station, Ananthapuramu, Andhra Pradesh. The experimental material utilized for the present study comprised of 40 castor genotypes (37 inbred lines + three varietal checks) sown under rainfed alfisols in alpha lattice design replicated twice. Each genotype was planted in two rows of 6 m length adopting a spacing of 90 cm × 60 cm. All the recommended package of practices were adopted to raise a healthy crop. Observations on five randomly selected plants in each genotype for 12 metric traits *viz.*, days to 50% flowering of primary spike, days to maturity of primary spike, plant height up to primary spike (cm), number of nodes to primary raceme, effective primary spike length (cm), number of capsules per primary spike, number of effective spikes per plant, volume weight (g/100ml), hundred seed weight (g), total seed yield (kg/ha), final plant stand (no./plot) and oil content (%) were studied. Oil content in castor genotypes were estimated by Nuclear Magnetic Resonance Spectrometry (NMR) at Indian Institute of Oilseeds Research (IIOR), Hyderabad. The statistical analysis for various traits was computed using R statistical package.

RESULTS AND DISCUSSIONS

High magnitude of variability enhances the likelihood of developing desirable genotypes in crop plants. The primary objective of germplasm conservation is to collect and preserve the genetic diversity of native crop species, ensuring its availability for both present and future generations. In this study, significant differences among the castor genotypes were observed for all the traits examined, indicating the presence of considerable genetic variability. Such variability provides a valuable resource for crop improvement programs, enabling breeders to select and combine favourable traits effectively.

Traits with high heritability are generally easier to improve through selection, as their expression is less influenced by environmental factors and are more controlled by genetic makeup. Consequently, the selection process becomes more efficient, and the response to selection is greater. These findings underscore the importance of evaluating and conserving diverse germplasm to sustain long-term agricultural productivity and adaptability to changing environmental conditions.

The analysis of variance, along with estimates of genetic variability, heritability, and genetic advance as a percentage of the mean, are presented in Tables 1 and 2. In addition, the graphical representation of PCV, GCV, heritability, and genetic advance as a percentage of the mean for various traits is illustrated in Fig. 1. Highly significant differences were observed among the genotypes for all the traits under study, indicating the presence of substantial genetic variability within the population. This variation provides ample scope for selection and genetic improvement. The character days to 50% flowering of primary spike recorded a mean of 55.60 days, while days to maturity of primary spike

averaged 105.80 days. The final plant stand had a mean of 17.40 plants per plot, and plant height up to the primary spike averaged 47.04 cm. The number of nodes to the primary raceme showed a mean of 13.27, whereas the effective primary spike length had a mean of 24.71 cm. The number of capsules per primary spike averaged 51.14, and the number of effective spikes per plant showed a mean of 3.42. Volume weight recorded a mean value of 64.53 g/100 ml, while hundred seed weight had an average of 29.09 g. The total seed yield had a mean of 807.32 kg/ha, and oil content averaged 47.69 %.

Range of variation for days to 50% flowering ranged from 49.00 to 65.00 days, and days to maturity varied between 98.00 and 116.00 days. Final plant stand ranged from 15.00 to 20.00 plants per plot, while plant height up to the primary spike varied from 34.30 to 55.30 cm. The number of nodes to the primary raceme fluctuated between 10.50 and 15.50, while effective primary spike length extended from 15.60 to 39.90 cm. The number of capsules per primary spike ranged from 40.40 to 83.10, and the number of effective spikes per plant varied between 2.10 and 5.10. Volume weight ranged from 58.45 to 71.55 g/100 ml, and hundred seed weight varied from 22.45 to 39.45 g. Total seed yield showed a wide range of variation from 495.50 to 1661.40 kg/ha, whereas oil content ranged between 44.00 and 51.85%.

High estimates of PCV and GCV were recorded for number of capsules per primary spike (20.10% and 21.34%) and total seed yield (20.10% and 21.34%). Moderate PCV and GCV were noticed for hundred seed weight (GCV 11.76%, PCV 13.12%). Moderate GCV and High PCV noticed for the traits effective primary spike length (GCV 19.92%, PCV 21.35%) and number of effective spikes per plant (GCV 18.48%, PCV 20.10%). Low GCV and moderate PCV were noticed for plant height up to primary spike (9.04%, PCV 13.48%). Low PCV and GCV were noted for number of nodes to primary raceme (GCV 7.16%, PCV 9.40%), and final plant stand (GCV 4.91%, PCV 7.68%) days to 50% flowering of primary spike (GCV 5.69%, PCV 6.48%), days to maturity of primary spike (GCV 3.47%, PCV 3.92%) and oil content (GCV 2.84%, PCV 4.49%).

The results are in consonance with the findings of Rukhsar *et al.* (2018) for days to 50% flowering of primary spike, Deepak *et al.* (2024) for Days to maturity of primary spike. Abimiku *et al.* (2012); Getinet *et al.* (2014); Dapke *et al.* (2016) for plant height up to primary spike. Getinet *et al.* (2014); Rukhsar *et al.* (2018) for number of nodes to primary spike. Dapke *et al.* (2016) for effective primary spike length. Deepak *et al.* (2024) for number of capsules per primary spike. Halilu *et al.* (2013); Rukhsar *et al.* (2018) for number of effective spikes per plant. Rajavardhan *et al.* (2023); Deepak *et al.* (2024) for total seed yield. Rajavardhan *et al.* (2023) for oil content.

High heritability accompanied by moderate genetic advance was observed for the days to 50% flowering of primary spike (77.01% and 10.28%). while high

heritability coupled with low genetic advance was recorded for days to maturity of primary spike (78.08% and 6.31%). These patterns suggest the involvement of non-additive gene action in the inheritance of these traits. Similarly, moderate heritability combined with moderate genetic advance was found for plant height up to primary spike (45.04% and 12.50%) and number of nodes to primary spike (58.02% and 11.24%), moderate heritability and low genetic advance (39.09% and 5.17%). This indicates limited scope for improvement of these traits.

High heritability associated with high genetic advance as a percentage of the mean was recorded for five traits viz., effective primary spike length (87.05% and 38.28%), number of capsules per primary spike (88.71% and 38.99%), number of effective spikes per plant (84.60% and 35.02%) and 100 seed weight (80.40% and 21.73%) and total seed yield (87.49% and 61.74%) This indicates a predominant influence of additive gene action in the expression of these traits, suggesting that direct selection would be rewarding towards genetic improvement.

Similar results were reported by Udaya *et al.* (2013); Dapke *et al.* (2016) for days to 50% flowering of primary spike. Rukhsar *et al.* (2018) for days to maturity of primary spike. Najan *et al.* (2010); Patel and Patel (2014) for plant height up to primary spike. Rukhsar *et al.* (2018) for number of nodes to primary spike, height up to primary spike (45.04% and 12.50%)

and number of nodes to primary spike (58.02% and 11.24%), moderate heritability and low genetic advance (39.09% and 5.17%). This indicates limited scope for improvement of these traits.

High heritability associated with high genetic advance as a percentage of the mean was recorded for five traits viz., effective primary spike length (87.05% and 38.28%), number of capsules per primary spike (88.71% and 38.99%), number of effective spikes per plant (84.60% and 35.02%) and 100 seed weight (80.40% and 21.73%) and total seed yield (87.49% and 61.74%) This indicates a predominant influence of additive gene action in the expression of these traits, suggesting that direct selection would be rewarding towards genetic improvement.


Similar results were reported by Udaya *et al.* (2013); Dapke *et al.* (2016) for days to 50% flowering of primary spike. Rukhsar *et al.* (2018) for days to maturity of primary spike. Najan *et al.* (2010); Patel and Patel (2014) for plant height up to primary spike. Rukhsar *et al.* (2018) for number of nodes to primary spike. Rukhsar *et al.* (2018); Deepak *et al.* (2024) for effective primary spike length and number of capsules on primary spike. Udaya *et al.* (2013) for number of effective spikes per plant. Sadaiah *et al.* (2021); Deepak *et al.* (2024) for hundred seed weight. Rajavardhan *et al.* (2023); Deepak *et al.* (2024) for total seed yield and Nalini and Yadavendrakumar (2013) for oil content.

Table 1: ANOVA for seed yield and its attributes in 40 castor genotypes.

Sr. No.	Traits	Mean squares			
		Replication (df:1)	Treatment (df:39)	Blocks (df:6)	Error (df:33)
1.	Days to 50% flowering of primary spike	5.51	23.04**	3.87	2.99
2.	Days to maturity of primary spike	6.05	30.66**	5.40	3.77
3.	Final plant stand	0.001	2.51**	1.35	1.06
4.	Plant height up to primary spike	9.11	58.27**	9.64	22.08
5.	Number of nodes to primary raceme	2.20	2.46**	0.36	0.65
6.	Effective primary spike length	1.98	52.06**	1.41	3.61
7.	Number of capsules per primary spike	3.36	224.65**	16.35	13.44
8.	Number of effective spikes per plant	0.02	0.872**	0.06	0.07
9.	Volume weight	1.13	23.84**	14.68	10.44
10.	Hundred seed weight	10.80	26.27**	1.84	2.85
11.	Total seed yield	984.91	143416.12**	9813.12	9569.11
12.	Oil content	3.87	6.42**	0.77	2.74

Table 2: Mean, variability, heritability (*bs*) and genetic advance as per cent of mean for 12 characters in 40 Castor genotypes.

Sr. No.	Trait	Mean	GCV (%)	PCV (%)	h^2b (%)	GAM (%)
1.	Days to 50% flowering of primary spike	55.66	5.69	6.48	77.01	10.28
2.	Days to maturity of primary spike	105.80	3.47	3.92	78.08	6.31
3.	Final plant stand	17.40	4.91	7.68	40.84	6.47
4.	Plant height up to primary spike	47.04	9.04	13.48	45.04	12.50
5.	Number of Nodes to primary raceme	13.27	7.16	9.40	58.02	11.24
6.	Effective primary spike length	24.71	19.92	21.35	87.05	38.28
7.	Number of capsules per primary spike	51.14	20.10	21.34	88.71	38.99
8.	Number of effective spikes per plant	3.42	18.48	20.10	84.60	35.02
9.	Volume weight	64.53	4.01	6.42	39.09	5.17
10.	Hundred seed weight	29.09	11.76	13.12	80.40	21.73
11.	Total seed yield	807.32	32.04	34.26	87.49	61.74
12.	Oil content	47.69	2.84	4.49	40.17	3.71



DFF-Days to 50% flowering of primary spike; **NCS** -Number of capsules per primary spike; **DTM**-Days to maturity of primary spike; **NES** -Number of effective spikes per plan; **FPS** - Final plant stand; **VWT**- Volume weight; **PLH**-Plant height up to primary spike; **HSW**- Test weight; **NPS**-Number of nodes to primary raceme; **TSY**- Total Seed yield; **ESL**- Effective primary spike length; **OCT**- Oil content

Fig. 1. Genotypic and phenotypic coefficient of variability, Heritability and Genetic advance as per cent mean for 12 traits studied in castor genotypes.

CONCLUSIONS

Number of capsules on primary spike and total seed yield have high PCV and high GCV indicates the presence of high range of variability. High heritability coupled with high genetic advance as a percentage of the mean was recorded for effective primary spike length, number of capsules on primary spike, number of effective spikes per plant, 100 seed weight and total seed yield. This indicated a predominant influence of additive gene action in the expression of these traits, suggesting that direct selection would be rewarding towards their genetic improvement in castor.

FUTURE SCOPE

Future research should focus on selecting yield attributing traits like effective primary spike length, number of capsules on primary spike, number of effective spikes per plant, 100 seed weight and total seed yield aid in genetic improvement of castor.

REFERENCES

- Abimiku, O. E., Azagaku, E. D. and Ndor, E. (2012). Genetic variability and correlation studies in some quantitative characters in castor (*Ricinus communis* L.) accessions. *Asian Journal of Agricultural Sciences*, 4(6), 368-372.

- Dapke, J. S., Naik, M. R., Vaidya, G.B., Vanve, P. B., Narwade, A. V. and Rajkumar, A. K. (2016). Genetic variability in castor (*Ricinus communis* L.). *European Journal of Biotechnology and Bioscience*, 4(4), 39-40.
- Deepak, K. A., Manjunatha, T., Hemalatha, V. and Chary, D. S. (2024). Variability, Correlation Patterns and Principal Component Analysis (PCA) for Seed Yield and Contributing Traits in Castor (*Ricinus communis* L.). *Journal of Advances in Biology & Biotechnology*, 27(8), 1217-1227.
- Delvadiya, I. R., Madariya, R. B. and Ginoya, A. V. (2021). Heterosis and Inbreeding Depression Study in Castor (*Ricinus communis* L.) Using 21 Generations. *Biological Forum – An International Journal*, 13(4), 26-32.
- Getinet, A., Beemnet, M. K., Girma, T and Chalachew, E. (2014). Phenotypic variability in Ethiopian castor (*Ricinus communis* L.) accessions. *International Journal of Advanced Biological and Biomedical Research*, 2(12), 2909-2914.
- Halilu, A. D., Aba, D. A. and Ogunwole, J. O. (2013). Genetic variability, genetic gain and relationships of yield and yield components in castor (*Ricinus communis* L.). *Research & Reviews in Biosciences*, 7(5), 181-186.
- INDIASTAT (2023-2024). Agriculture Area and Crops Growth Statistics 2023, 2024 [online] Available at: <<https://www.indiastat.com/data/agriculture>> [Accessed 18 June 2024].
- Lokesh, G. B., Dandoti, K. and Wali, V. B. (2020). Trends in the direction of trade of Castor oil exports from India. *Indian Journal of Economics and Development*, 16(3), 381-388.
- Najan, B. R., Kadam, J. R. and Kadlag, A. D. (2010). Genetic variability studies in castor (*Ricinus communis* L.). *Journal of Oilseeds Research*, 27, 77-79.
- Nalini, T. and Yadavendrakumar (2013). Genetic variability and character association in castor (*Ricinus communis* L.). *Current Advances in Agricultural Sciences*, 5(2), 256-258.
- Patel, J. K. and Patel, P. C. (2014). Genetic Variability, Heritability and Genetic Advance for yield and yield components in castor genotypes. *International Journal of Plant Science*, 9(2), 385-388.
- Rajavardhan, Y., Lal, J. J., Lakshamma, P., Kumar, C. V. and Kumar, A. A. (2023). Role of Genetic Variability and Their Associated Character on Yield Attributing Traits in Castor (*Ricinus communis* L.) Germplasm Lines. *International Journal of Environment and Climate Change*, 13(10), 265-271.
- Reddy, P. S., Proya, P. B., Sadaiah, K., Vanisri, S. and Kumar C. V. S. (2022). Genetic Variability and Character Association Studies among Yield Attributing Traits in Genepool Lines of Castor (*Ricinus communis* L.). *Biological Forum – An International Journal*, 14(2), 1277-1281.
- Rukhsar Patel, M. P., Parmar, D. J. and Kumar, S. (2018). Genetic variability, character association and genetic divergence studies in castor (*Ricinus communis* L.). *Annals of Agrarian Science*, 16, 143-148.
- Sadaiah, K., Neelima, G., Rani, C. V. D., Rani, V. D., Madhuri, G., Nalini, N., Sujatha, M., Shankar, V. G., Kumar, M. P., Lavanya, C. and Lal, J. J. (2021). Genetic parameters, diversity and character association studies in germplasm lines of Castor (*Ricinus communis* L.). *Electronic Journal of Plant Breeding*, 12(4), 1134-1141.
- Senthilvel, S., Shaik, M., Anjani, K., Shaw, R. K., Kumari, P., Sarada, C. and Ushakiran, B. (2016). Genetic variability and population structure in a collection of inbred lines derived from a core germplasm of Castor. *Journal of Plant Biochemistry and Biotechnology*, 26(1), 27-34.
- Udaya, B. K., Satyanarayana Rao, V. and Srinivasa Rao, M. (2013). Genetic variability in castor (*Ricinus communis* L.) for yield and its contributing traits. *International Journal of Food and Veterinary Sciences*, 3(3), 103-108.
- Zheng, L., Qi, J. M., Fang, P.P., Su, J. G., Xu, J. T. and Tao, A. F. (2010). Genetic diversity and phylogenetic relationship of castor as revealed by SRAP analysis. *Wuban Zhiwuxue*, 28, 1-6.

How to cite this article: K. Chandana, A.V.S. Durga Prasad, N. Sabitha and M. Raghavendra (2025). Genetic Variability Studies in Castor Germplasm (*Ricinus communis* L.). *Biological Forum*, 17(8): 75-79.