



## Impact of Aloe vera Juice as Supplementary Nutrient on Larval, Cocoon and Silk Parameters in *Bombyx mori* L.

Roja V.<sup>1</sup>, Yashaswini B.R.<sup>2</sup> and Sushmitha C.<sup>2\*</sup>

<sup>1</sup>Department of Studies in Sericulture Science,

University of Mysore, Manasagangotri, Mysore (Karnataka), India.

<sup>2</sup>Department of Sericulture, College of Agriculture, GKVK, Bengaluru (Karnataka), India.

(Corresponding author: Sushmitha C.\*)

(Received: 06 June 2025; Revised: 22 July 2025; Accepted: 17 August 2025; Published online: 09 September 2025)

(Published by Research Trend)

DOI: <https://doi.org/10.65041/BiologicalForum.2025.17.9.6>

**ABSTRACT:** The silkworm *Bombyx mori* L. is a lepidopteran insect used as laboratory tool for various experiments. Being a domesticated insect it has been reared mainly for the production of silk. The successful production of the silk not only governed by dynamic environmental factors but also affected by quality of mulberry leaf provided to silkworm. The feeding of nutritionally enriched leaves provide better growth and development of silkworm larvae, as well as directly influence the quality and quantity of silk production. Hence, dietary nutrients are the most important factor which determine quality and quantity of silk production. An experiment was carried out to know the impact of mulberry leaf fortified with aloe vera at varied concentrations viz., 0.2, 0.4, and 0.6% on larval, cocoon and silk parameters in the FC2 silkworm bivoltine hybrid. The study reveals that larvae feed on mulberry fortified with aloe vera at 0.6% treatment yielded the most pronounced improvements in economic parameters. Specifically, larval weight (4.96 g), cocoon weight (2.13 g) shell weight (0.50 g), pupal weight (1.62 g), shell ratio (23.64 %), filament length (1288m), filament weight (0.43 g), renditta (0.43), denier (3.04) and raw silk percentage (20.45 %) were all significantly higher compared to both control and lower-concentration treatments. The outcome of the study indicated that, there is profound increase due to aloe vera treatment in larval growth and cocoon characters than the control with enhanced quantity and quality silk production.

**Keywords:** Aloe vera, fortification, mulberry, silkworm, cocoon and silk parameter.

### INTRODUCTION

Silkworm *Bombyx mori* L. is a well-known lepidopteran (Family: Bombycidae), the larval instars of which feed on the leaves of mulberry (*Morus* sp). The growth, development of larva and subsequent cocoon production are greatly influenced by nutritional quality of mulberry leaves. Needless to say, it has to derive its nutritional requirements from mulberry leaves for its growth and development. Any variation in the nutritional components of mulberry leaves may have some influence on the growth and development of silkworm. Though the availability of silkworm nutrients in mulberry leaf is ensured on most occasions, some time they may not be available in adequate quantities for the larval growth. The quality of the leaf can have profound influence on the superiority of silk produced by *B. mori*. Therefore, the production of good cocoon crop is largely dependent on the quality of leaves. In other words, leaves of superior quality enhance the chances of realizing good cocoon crop (Ravikumar, 1988). It has also been demonstrated that the dietary nutrition management has a direct influence on quality and quantity of silk production in *B. mori*. The

nutritional parameters of mulberry leaves and silk production are related to each other. The nutritional composition of mulberry leaves varies depending on the nature of the genotype, soil and environmental conditions, precipitation (rainfall/irrigation), agronomical practices employed, plant protection measures adopted, etc. (Murugan *et al.*, 1998).

Cocoon production by *B. mori* depends on its rearing on mulberry leaves. For this reason, the quality of its food (mulberry leaf) needs to be high for realizing the full potential of the silkworm in terms of cocoon production. Nevertheless, the quantity of leaf fed by the larva also assumes great significance for accumulating considerable biomass that should culminate in the improvement of cocoon yield. Understandably, whenever the leaf quality is inferior, there arises a need for fortifying the leaf using supplementary nutrients such as vitamins, proteins, amino acids, carbohydrates, minerals, etc. so that the nutritional status of the leaf would be ultimately matching with the nutritional requirement of the silkworm. In addition, there is also a necessity to enhance the appetite in the silkworm so that it would grow better and to improve the disease

tolerance that should facilitate higher rate of survival of the larvae (Legay, 1958).

Like in other organisms, in the silkworm, *B. mori* too, nutrition plays an important role in improving the growth and development. The intake of nutrient by the silkworm larvae is influenced by the availability of feed. Therefore, silkworm nutrition is considered as a major area of research in sericulture (Legay, 1958). Nutrition study in silkworm is an essential pre requisite for its proper commercial exploitation. *B. mori* requires specific essential sugars, amino acids, proteins and vitamins for its normal growth, survival and also for the silk gland growth. Akhtar and Asghar (1972) found that vitamins and mineral salts played an important role in the nutrition of silkworm. Karakasy (1990) showed that silkworms fed on mulberry leaves enriched with ascorbic acid showed better growth and produced heavier cocoons. Similar observations were made by Madhu Babu (1992).

The feeding of nutritionally enriched leaves showed better growth and development of silkworm larvae and, in turn, influenced the quality and quantity of silk produced. Nearly 70% of the silk proteins produced by silkworm are directly derived from the protein of the mulberry leaves. In recent years a number of botanicals have been identified with insect growth regulation (IGR), juvenile hormone (JH) secretion, etc. Experimental evidence consistently shows that fortifying mulberry leaf with vitamin C, B-complex vitamins or minerals viz., Zn significantly improves larval growth, silk gland development and cocoon productivity in silkworm, *Bombyx mori* (Yashaswini *et al.*, 2025). Fortifying mulberry leaf with amino acids (L-Serine and glycine) or protein rich supplements (drone brood, protinex) significantly improves larval growth, silk gland development and cocoon productivity in silkworm, *Bombyx mori*. Optimal gains are observed at moderate supplementation levels during the fifth instar (Yashaswini *et al.*, 2025). However, silkworm, *B. mori* supplemented with plant extracts namely *Tribulus terrestris*, *Pssoralea coryleifolia*, *Phyllanthus niruri*, *Polypodium Sp.*, *Parthenium hysterophorus* and *Tridax procumbens* increases larval growth as well as cocoon parameters (Rajashekargouda, 1991; Murugan *et al.*, 1998). According to Mamadapur (1994); Santosh Kumar (1997), application of Lantana camara and Clerodendroninermæ at 5% once on 48h old fifth instar larvae increased all the economic traits. A good number of reports are available on fortifying agents, feed additives and botanical extractions which enhances economics parameters of the silk worm. However, study on impact of botanical extracts on larval, cocoon and silk parameters of silkworm is meager. In this context, the present study was under taken.

## MATERIAL AND METHODS

The experiment was carried out in the Department of studies in sericulture science, Manasagangaotri, University of Mysuru, which is which is located at 12° 18' 26" North Latitude and 76° 38' 59" East Longitude at an altitude of 770 m (2,530 ft) above mean sea level.

Roja *et al.*,

*Biological Forum*

**Maintenance of mulberry garden:** The experiment was conducted using mulberry garden (V-1 variety) maintained at Manasagangothri campus, Mysuru. The tender and medium mulberry leaves were used for young-age silkworm rearing. While, coarse leaves were used for late-age rearing. To maintain good quality leaf for feeding, proper methods were followed during harvesting and storage. Fresh leaves were collected during morning and evening hours of the day and stored in leaf preservation chamber covered with wet gunny cloth to reduce loss of moisture and degradation of nutrients (Sekharappa *et al.*, 1991).

**Disinfection of rearing house and appliances:** To ensure the rearing house and appliances pathogen free, they were thoroughly cleaned with water and 2% bleaching powder solution followed by 2.5% Sanitech solution @1.5 liter per square meter to achieve disinfection (Dandin *et al.*, 2003).

**Procurement of disease free layings (DFLs):** The disease free layings (eggs) of popular bivoltine silkworm hybrids namely FC2 (CSR6 × CSR26) was procured from the National Silkworm Seed Organization (NSSO), Central Silk Board (CSB), Mysuru. The egg cards were surface sterilized by dipping them in 2% formalin for 8-10 minutes and dried in shade before use.

**Incubation of DFLs:** The eggs of FC2 were incubated at room temperature (25°C) and 70-80% relative humidity by adopting standard incubation method and were subjected to black boxing for about 48 h at pin head stage (Benchamin and Nagaraj 1987).

**Rearing of FC2 silkworm:** After brushing the eggs, newly hatched silkworms were carefully shifted on to the rearing bed and fed fresh tender leaves and reared by employing stranded rearing techniques advocated by Dandin and Giridhar (2010).

Selection of plant extract

Aloe vera Classification

Kingdom-Plantae

Order - Asparagales

Family - Asphodelaceae

Sub family- Asphodeloideae

Genus - Aloe

Species - vera

Aloe vera is a stem less or very short-stemmed plant growing to 60-100cm tall, spreading by offsets. The leaves are thick and fleshy, green to grey-green with some varieties showing white flecks on their upper and lower stem surfaces. The margin of leaf is serrated and has small white teeth. The flowers are produced in summer on a spike up to 90cm tall. Aloe vera forms arbuscular mycorrhiza, a symbiosis that allows the plant better access to mineral nutrients in soil (Damodhara *et al.*, 2024).

**Preparation of plant extract.** The aloe vera (Dabur) procured from Charaka Ayurvedic Medicines and General Stores, Brindavan Extension Mysuru-20. The stock solution was diluted at the rate of 0.2, 0.4 and 0.6% by using distilled water. The prepared solution was preserved in refrigerator at 4°C until further use.

For the experimentation, the larvae is divided into five batches viz., batch I (T1), batch II (T2) and batch III

(T3) were reared with mulberry leaf fortified with amla juice at 0.2, 0.4 and 0.6%, respectively. Whereas batch IV (T4) larvae were reared on mulberry leaf sprayed with distilled water (control) and batch V (T5) larvae were reared on mulberry leaf alone (absolute control). For each treatment, three replications were maintained and each with 100 larvae along with controls.

**Silkworm larval and economic parameters:** The silkworms were carefully monitored every day and from the first day of V instar moulting, the larval weight was determined everyday till the worms started to spin. Then the cocoon weight, pupal weight, shell weight, shell ratio, silk productivity, filament length, filament weight and denier were recorded.

**Larval weight (g):** For recording mature larval weight (g), ten larvae were randomly selected from each treatment, replication-wise during fifth day of fifth instar.

**Cocoon weight (g):** Cocoon weight (g) was calculated by taking ten randomly selected cocoons from all groups and weighed using an electronic balance. Weight of each cocoon from each group was recorded separately.

**Pupal weight (g):** Pupal weight (g) was weighed using an electronic balance, after removing the floss. The cocoons were cut open and the pupae were taken out without causing any damage to them.

**Shell weight (g):** Shell weight (g) was calculated by taking randomly selected 10 cocoons and cut open with

the help of a blade and the shell weight was taken accurately.

The parameters such as shell ratio, silk filament length, denier, renditta and raw silk percentage were calculated by using following formulae:

$$\text{Shell ratio (\%)} = \frac{\text{Shell weight (g)}}{\text{Cocoon weight (g)}} \times 100$$

$$\text{Filament length (L)} = R \times 1.125$$

Where, L = total length of the silk filament

R = number of revolutions recorded

1.125 = circumference of epprouvette in meter

$$\text{Denier} = \frac{\text{Total weight of reeled silk (g)}}{\frac{\text{Total length of reeled silk (m)}}{\text{Weight of cocoon reeled (g)}}} \times 9000$$

$$\text{Renditta} = \frac{\text{Total weight of reeled silk (g)}}{\text{Weight of raw silk (g)}} \times 9000$$

$$\text{Raw silk percentage} = \frac{\text{Silk weight (g)}}{\text{Cocoon weight (g)}} \times 100$$

## RESULT AND DISCUSSIONS

**Larval and Cocoon weight:** Silkworm larvae fed on mulberry leaves fortified with aloe vera juice expressed variation in respect of larval weight with maximum being at 0.6% (4.965 g) followed by 0.4 % (4.692) and 0.2% (4.523g) respectively. While, it was minimum in absolute (4.487g) and distilled water controls (4.564g). The highest cocoon weight of 2.132g 0.6% followed by 0.4% (2.090g) and 0.2% (2.010g). As against to this, it was lowest in absolute (1.901g) and distilled water (1.950g) control batches (Table 1).

**Table 1: Effect of mulberry leaves fortified with aloe vera juice at different concentrations on larval and cocoon parameters of FC2 silkworm hybrid.**

Concentration (%)	Larval Weight (g)	Cocoon Weight (g)	Pupal Weight (g)	Shell Weight (g)	Shell ratio (%)
0.2	4.52±0.04	2.01±0.01	1.56±0.12	0.44±0.12	21.94±0.54
0.4	4.69±0.007	2.09±0.002	1.61±0.07	0.47±0.07	22.54±0.25
0.6	4.96±0.05	2.13±0.53	1.62±0.03	0.50±0.13	23.64±0.42
Control(Dil. water)	4.56±0.03	1.95±0.42	1.53±0.09	0.41±0.09	21.38±0.52
Absolute control	4.48±0.02	1.90±0.003	1.49±0.10	0.40±0.10	21.15±0.35

It is evident that aloe vera juice is a richest source of vitamin A, E and other nutrients and supplementation of aloe vera juice obviously boost the immune system of the larva which enhance larval weight in turn reflect on more cocoon weight. Perhaps, many vitamins and biomolecules are found in the diet are phagostimulatory in nature and it could be one of the reasons which enhance food intake by the silkworm resulting in increased larval and cocoon weight (Mane *et al.*, 1997). These results are on line with earlier observations of Pratheesh *et al.* (2007), who have reported that silkworm larvae supplemented with *Achyranthes aspera* (4.04g), *Achyranthes recimosus* (4.00g), *Parthenium hysterophorus* (3.920g), *Tribulus terrestris* (4.02g) and *Withania somnifera* (4.02g) at 5% registered highest larval weight over remaining concentrations (3 and 8%) as well as control batches (3.8g). Further, Rajashekaragouda (1991) opined that administration of *Psoralea corylifolia* L. and *Tribulus terrestris* L. plant extracts at 10µg per larvae increased the cocoon weight. Barge and Pardeshi 2018, also noticed that increase in larval weight due to *Sida acuta*

plant extract at 0.5, 1.0, 1.5 and 2.0%, (2.336, 2.401, 2.419 and 2.559g), respectively against control (2.319g), similarly, Anil Kumar and Prashanth (2018), have noticed that, larvae supplemented with soya bean flour at 2, 4 and 6% exerted higher larval and cocoon weight over control. As per Manjula *et al.* (2011) larvae fed on mulberry leaf enriched with *Dolichos lablab* and *Vigna unguiculata* at 7.5% concentration has resulted increased larval and cocoon weight as against control. Mamadapur (1994) who reported the administration of 4µl of *Latana camara* and 7µl of *Clerodendron inermae* leaf extracts increases the larval weight. Kiran Kumara *et al.* (2024) also found that worms fed on mulberry leaves fortified with amla pulp juice at 0.2% concentration recorded higher larval and cocoon weight of 4.665 and 2.032g, respectively over control. Further, Krishna Prasad *et al.* (2001) opined that increase in larval weight in the cross breed PMxNBAD, was due to supplementation of mulberry leaves with potato leaf extract. Similar trend was also noticed by Shivakumar *et al.* (1995); Rajashekaragouda (1991); Takhliq (2012) in other plant extracts.

**Pupal weight:** The pupa formed by the larvae reared on mulberry leaves treated with aloe vera juice at 0.6% registered maximum pupal weight of (1.628g) followed by 0.4% (1.619g) and 0.2% (1.569g) over distilled water (1.533g) and absolute (1.499g) controls (Table 1). The silkworms fed on mulberry leaves sprayed with aloe vera at 0.6% concentration exerted highest pupal weight (1.628 g). The increase in pupal weight may be due to the increase in feeding efficacy during Vth instar larvae by the additional supplementation of aloe vera juice. These results are also supported by the observations of Sundar Raj *et al.* (2001) who reported that supplementation of soyabean flour to the silkworm recorded highest pupal weight of 1.23g over control batch (0.85g). Similarly, Kiran Kumara *et al.* (2024) found that silkworms supplemented with amla at 2.0% concentration recorded higher pupal weight of 1.563g over control.

**Shell weight and Shell ratio:** The larvae reared on mulberry leaves fortified with aloe vera juice at varied concentrations exhibited notable differences in respect of shell weight with highest being at 0.6% (0.504g) followed by 0.4% (0.471g) and 0.2% (0.441g) respectively. Whereas, it was least in absolute and distilled water controls (0.417 and 0.402g) respectively (Table 1).

The shell ratio and filament length are the most important economic parameters (Chanda *et al.*, 2013) and supplementation aloe vera juice might have accelerated biosynthesis of sericin and fibroin in turn resulting in increased shell ratio and filament length. These results are more or less parallel with finding of Walaa *et al.* (2018), who have opined that larvae supplemented with ascorbic acid, lemon and sweet orange enhance shell percentage of 23.81, 22.07 and 21.49% over control (21.15%). This type of trend was observed for filament length at 4% concentration. Sisodia *et al.* (2019) have reported increased shell ratio and filament length with amla extract. Similar results were also observed with the supplementation of tender coconut (Amit *et al.*, 2015) and with Aloe vera (Vitthalrao and Anil 2012). These results are in agreement with the earlier observations of Rajashekaragouda (1991) who reported that increased shell weight in petroleum ether extract of *P. corylifolia* and *T. terrestris* and same was reported by Mamadapur (1994) with plant extract of *C. inermis* and *L. camara*. Further, significant higher shell ratio of 16.37% on potato leaf extract was also observed by Krishna Prasad *et al.* (2001). Kiran Kumara *et al.* (2024) also found that silkworms supplemented with amla at 2.0% concentration recorded higher shell weight and shell ratio of 0.469 and 23.08%, respectively over control.

**Table 2: Effect of mulberry leaves fortified with aloe vera juice at different concentrations on silk parameters of FC2 silkworm hybrid.**

Concentration (%)	Filament Length (m)	Filament Weight(g)	Denier	Renditta	Raw silk %
0.2	1197±18.52	0.37±0.07	2.80±0.10	5.38±0.29	18.56±0.63
0.4	1224±44.23	0.40±0.02	2.96±0.08	5.18±0.10	19.28±0.38
0.6	1288±84.25	0.43±0.08	3.04±0.78	4.89±0.30	20.45±0.72
Control(Dil. water)	1060±119.4	0.34±0.04	2.96±0.09	5.58±1.14	17.9±0.70
Absolute control	1034±62.42	0.33±0.02	2.90±0.12	5.69±0.24	17.57± 1.07

**Filament length:** It is one of the major contributing quantitative traits in silkworm. The silkworm reared on mulberry leaves supplemented with aloe vera recorded marked differences with regard to filament length with maximum being at 0.6% (1288 m) followed by 0.4% and 0.2% (1224 and 1197m). On the other hand, it was shortest in absolute (1034m) and distilled water controls (1060m) (Table 2).

The silkworms fed on mulberry leaves sprayed with aloe vera at 0.6% concentration exerted longer filament length of 1288m against control batches (1060m). The increase in filament length may be due to higher silk protein synthesis by the additional supplementation of aloe vera juice. These results are also supported by the administration of 2% aloe vera extract to silkworm increases filament length of 1003.0m over control batch (Vitthalrao and Anil 2012). Similar results were also noticed with *Clerodendron inermis* supplementation (Mamadapur, 1994). Further, Kiran Kumara *et al.* (2024) revealed that silkworms supplemented with amla at 0.2% concentration expressed higher filament length 1204m as against respective control.

**Filament weight and Denier:** The silkworm larvae supplemented with aloe vera at 0.6% exerted highest filament weight of (0.436g) followed by 0.4% (0.403 g) and 0.2% (0.373g) over absolute (0.349g) and distilled water (0.334g) controls. Denier denotes size of filament obtained from cocoons of silkworm breeds. Marginal variation was noticed in respect of denier among breeds provided with mulberry leaves fortified with aloe vera juice at different concentrations. In FC2 silkworm shows lowest denier at 0.2% (2.80) followed by 0.4% (2.96), absolute control (2.90). As opposed to this, highest denier was noticed with 0.6% (3.04) (Table 2). The larvae supplemented with aloe vera at 0.6% concentration recorded highest filament weight of 0.43g and on the other hand, lowest denier of 2.80 recorded at 0.2% concentration. However, not much more variations was registered for denier in distilled water control (2.96) and absolute control (2.90). The increase in filament weight and denier may be due to higher utilization of aloe vera juice for the synthesis silk proteins. The results corroborate the earlier finding of Pardeshi *et al.* (2014); Pardeshi *et al.* (2017) who have noticed that silkworms supplemented with *Amaranthus*

*hybridus* extract at 2% concentration enhance the filament weight (0.198g) and denier (2.004). Similar results were observed with *Alpermanthera sessilis* extract at 2per cent. According to Saritha Kumari *et al.* (2011), lowest denier of 2.31 and 2.36 were recorded by the silkworm supplemented with *Phyllanthus niruri* and *Adathoda vasica* extracts against control (2.41). Furthermore, Sridevi (2003), noticed increased filament 535 weight and denier with *Withania somnifera*, *Terminalia arjuna* and *Terminalia cordifolia* plant extracts. Similar result was also observed with supplementation of *Xanthium indicum* extract at 2.0% (Pardeshi *et al.*, 2014). Kiran Kumara *et al.* (2024) also found that silkworm supplemented with amla at 0.2% concentration exerted higher filament weight and denier of 0.401g and 2.998 respectively, over control.

**Renditta and Raw silk percentage:** Renditta trait indicates actual silk available from the cocoon. Notable variation was noticed with respect to renditta among breed provided with mulberry leaves fortified with alovera juice at varied concentration. In the FC2 silkworm lowest renditta was recorded at 0.6% (4.489) followed by 0.4% (5.186) and 0.2% (5.389). In contrast highest renditta was seen in absolute and distilled water control batches. The raw silk percentage indicates that the amount of silk obtained or produced from the cocoon. The mulberry leaves fortified with the aloe vera juice shows raise in the raw silk percentage. The 0.2% concentration shows the lowest raw silk percentage of 18.56% followed by 0.4% and 0.6% concentrations (19.28 and 20.45%), respectively. While, it was lowest in absolute and distilled water control (17.57 and 17.9%) (Table 2).

The silkworms supplemented with aloe vera at 0.6% concentration expressed lowest renditta and higher raw silk percentage (4.89 and 20.45%) respectively as against control batch (5.58 and 17.9%). It is presumed that the silkworm larvae absorbed and utilized the nutrients of aloe vera juice might have enhance these parameters. These results are in conformity with those of Sujatha *et al.* (2015) who reported that mulberry leaves fortified with phytochemicals reserpine and picrotoxin enhance raw silk percentage. Similarly, lowest renditta was recorded in drumstick leaf extract over other leaf extracts might be due to utilization of additional nutrients for the cocoon formation (Anil Kumar *et al.*, 2017). These results are in agreement with the earlier observation of Anil Kumar and Prashanth, (2018) who have opined that silkworm larvae CSR2 supplemented with 4% soybean flour registered higher raw silk percentage of 19.22% and lower renditta of 6.850 as against control 15.93% and 7.550, respectively. Similarly, Kiran Kumara *et al.* (2024) reported that silkworms supplemented with amla at 0.2% concentration registered lowest renditta (5.067) and highest raw silk percentage (19.73%) over respective control.

## CONCLUSIONS

The silkworm larvae nourished by mulberry larvae fortified with aloe vera revealed promising results. The larvae fed mulberry leaves supplemented with 0.6%

aloe vera, enhancing all the commercial parameters. Hence, farmers can use aloe vera (0.6%) for commercial rearing. Mulberry leaves are a traditional food for silkworms, but adding aloe vera to their diet seems to have improved their growth and other commercial parameters. Aloe vera is rich in vitamin A, C, E and other nutrients, which could be contributing to the positive effects observed in the silkworms. By supplementing mulberry leaves with 0.6% aloe vera, farmers can potentially enhance the quality and yield of silk produced by silkworms. This could be particularly beneficial for commercial silk production, as it may lead to higher yields or better-quality silk, ultimately increasing profitability for farmers. Overall, this finding holds promise for silk farmers looking to improve their yields and product quality of mulberry.

## FUTURE SCOPE

Systematic field trials (scaling up the best performing supplements e.g. 0.6 % aloe vera), implementation studies (evaluating cost, ease of use and farmer acceptance under real-world conditions), genotype-specific trials (assessing whether particular silkworm strains respond better to certain supplements) and interdisciplinary collaborations integrating entomology, molecular biology, biotech and sustainable agriculture.

## REFERENCES

- Akhtar, M. and Asghar, M. (1972). Role of vitamins and mineral salts in silkworm nutrition. *Journal of Sericulture Research*, 10, 123-130.
- Amit, S., Kalyankar Hugar, Govind and Murkute (2015). Influence of tender coconut water on the economic parameters of silkworm, *Bombyx mori* L. *Journal of Global Bioscience*, 4, 3325-3329.
- Anil Kumar, M. N., Jagadish, M. C. and Mahadevaswamy, M. (2017). Impact of aqueous plant extract on the economic traits of the silkworm breeds CSR2. *International Journal of Academic Research*, 2, 94-100.
- Anil Kumar, M. N. and Prashanth, J. (2018). Influence of mulberry leaf with soya bean flour supplementation on the economic traits and amino transferase activity in *Bombyx mori* L. *International Journal of Zoology Studies* 3, 59-64.
- Barge, S. B. and Pardeshi, A. B. (2018). Influence of dietary supplementation of *Sida acuta* plant extract on the mulberry silkworm, *Bombyx mori* L. *International Journal of Zoology Studies* 3, 199-202.
- Benchamin, K.V. and Nagararaj, C.V. (1987). Silkworm rearing techniques in appropriate sericulture techniques (Ed. Jolly, M.S.). CSR&TI, Mysore, pp.63-106.
- Chanda, S., Saha, L. M., Das Kar, N. B. and Bindroo, B. B. (2013). Correlation among shell percent, cocoon yield and reeling parameters of Multi × Bi cocoons under different agro-climatic conditions of west Bengal, India. *Int. Jr. Indust. Entomology*, 26, 74-80.
- Damodhara, G. N., Shalini, K. R., Karthik, R., Manjunatha, Gunashekhar, H., Mahendra, K. R., Joopaka Ramu, Suresh Kumar Mahala, Yashasvini Goswamy and Roja, V. (2024). Study of the Influence of *Aloe vera* Juice at varied Concentrations on Biomolecules in the Silkworm Hybrid FC2. *Biological Forum – An International Journal*, 16, 192-197.
- Dandin, S. B., Giridhar, K. and Rao, A. P. (2003). Disinfection protocols for rearing houses and appliances.

- Sericulture Hygiene Practices, 3rd ed., Central Silk Board, Bangalore, India.
- Dandin, S. B. and Giridhar, K. (2010). Handbook of Sericulture Technologies. Central Silk Board, Bangalore, p. 427.
- Karaksy, M. (1990). Effect of ascorbic acid-enriched mulberry leaves on silkworm growth and cocoon production. *Sericulture Science*, 21, 87-92.
- Kiran Kumara D, M. N. Anil Kumar, Sandya B. Raju and Chethan B. K. (2024) Application of Amla Juice as Supplementary Nutrient on Biomolecules and Commercial Parameters in *Bombyx mori* L. *Centre for Advanced Research in Agricultural Sciences*, 15(02) 531-538.
- Krishna Prasad, N. K, Sannappa, B., Reshma, T. K. and Keshava Prasad, P. K. (2001). Effect of solanaceous leaf extracts on growth, development and cocoon traits of PM X NB4D2 silkworm *Bombyx mori* L. *Bull. Ind. Acad. Seri*, 5, 10-15.
- Legay, J. M. (1958). Nutrition study as a major area of research in sericulture. *Proceedings of the International Congress of Sericulture*, 9, 150-155.
- Mane, J., Rayer, S. G., Patil, G. M. and Vage, M. K. (1997). Mul-berry- A Hobson's choice. *Indian Silk*, 36, 38-41.
- Madhu Babu, P. (1992). Effects of feeding nutritionally enriched leaves on silkworm larvae and silk production. *Journal of Applied Sericulture*, 31, 185-190.
- Mamadapur, G. (1994). Application of *Lantana camara* and *Clerodendron inermae* in silkworm rearing for enhanced cocoon yield. *Journal of Sericulture Research*, 2, 64-69.
- Manjula, S., Sabhanayakam, S., Mathivanan, V. and Saravanan, N. (2011). Modulations in the haemolymph of silkworm *Bombyx mori* (L). (Lepidoptera: Bombycidae)] fed with mulberry leaves augmented with cowpeas (*Vigna unguiculata*). *Int. Jr. Nut. Phar. Neuro. Diseases*, 1, 64-68.
- Murugan, K., Jeyabalan, D., Senthikumar, N., Senthilnathan, S. and Sivaprakasam, N. (1998). Growth Promoting effect of plant products on silkworm. A Biotechnological Approach. *Journal of Scientific and Industrial Research*, 57, 740-745.
- Pardeshi, A. B. and Bajad, P. N. (2014). The effect of nutritional supplementation with *Amaranthus hybridus* Linn. extract of economic performance of mulberry silkworm, *Bombyx mori* L. *Sch. Acad. Jr. Bioscience*, 2, 272-276.
- Pardeshi, A. B. and Barge, S. B. (2017). Impact of oral supplementation of *Alternanthera sessilis* plant extract on the economic parameters of silkworm, *Bombyx mori* L. *Bioscience Discovery*, 8, 596-601.
- Pratheesh Kumar, P. M., bhargava, S. K., Prabhakar, C. J. and Kamble, C. K. (2007). Botanicals mediated reproductive enhancement in mulberry silkworm (*Bombyx mori* L.) *Int. Jr. Indust. Entomology*, 14, 45-50.
- Rajashekharagouda, R. (1991). Studies on methods to increase silk yield on *B. mori* L., Ph.D. Thesis, Tamil Nadu Agricultural University, Coimbatore, pp. 122-138.
- Ravikumar, M. (1988). Influence of mulberry leaf quality on cocoon production. *Sericulture Review*, 15, 99-105.
- Santosh Kumar, G. (1997). Application of *Lantana camara* and *Clerodendron inermae* in silkworm rearing for improved economic traits. *Sericulture Bulletin*, 7, 45-50.
- Saritha Kumari, S., Bhaskar, R. N. and Latha, S. (2011). *In vivo* effect of medicinal botanicals on rearing parameters of the silkworm, PM × CSR2. *Current Biotica*, 5, 40-45.
- Sekharappa, B. M., Muniraju, E., Gururaj, C. S. and Chaluvachari (1991). Studies on the preservation of tender leaves for young age silk worm rearing. Paper presented in 10<sup>th</sup> National Symposium on life Science, Mysore, Abstract No. 4.
- Shivkumar, G. R. and Magadhun, S. B. (1994). Effect of phytoestrogens on the larval maturation and other economic character of the silkworm, *Bombyx mori* L. Annual report of the CSR&TI. pp 76-80.
- Sisodia, N. S. and Gaherwal, S. (2019). Effects of amla extract (*Phyllanthus emblica*) on biological and economical characteristics of the silkworm, *Bombyx mori*. *Pramana Research Journal*, 9, 1022-1028.
- Sridevi, G. (2003). Effect of mulberry leaves fortified with medicinal botanicals on the performance of mulberry silkworm, *Bombyx mori* L. M. Sc (Seri) Thesis, Univ. Agri. Science, Bangalore. pp 116.
- Sujatha, K., Satish, J. and Anitha, J. (2015). Effect of medicinal botanical (*Ocimum sanctum*), Family, Labiateae on commercial parameters of the silkworm, *Bombyx mori*, L. *International Journal of Multidisciplinary and Current Research* 3, 76-78.
- Sundar Raj, S., Chinnaswamy, K. P. and Neelu Nangia (2001). Soyabean to boost cocoon production. *Indian Silk*, 39, 11-13.
- Takhlique, M. D. (2012). Medicinal plant extracts and their valuable effect on various parameters of silkworm hybrid PM × CSR2. *International Journal of Green and Herbal Chemistry*, 1, 391-394.
- Vitthalrao, B. K. and Anil, N. S. (2012). Influence of aloe vera (L.) herbal formulation on the larval characters and economic parameters of silk worm *Bombyx mori* L. (Race: PM × CSR2). *The Ecoscan*, 1, 321-326.
- Walaa Sallam, El-Sharkawy F, El-sharkawy, Saad (2018). Technological effects of lemon, sweet orange juice and ascorbic acid as supplements additive on mulberry silkworm larvae, *Bombyx mori* L. *Jr. Product and Development* 23, 135-145.
- Yashaswini, B. R., Sushmitha, C. and Veenita, M. K. (2025). A review on fortification of mulberry leaf with vitamins and mineral supplements: Impact on larval development and cocoon quality of *Bombyx mori*. *Journal of Basic and Applied Research International*, 31(4) 97-109.

**How to cite this article:** Roja V., Yashaswini B.R. and Sushmitha C. (2025). Impact of Aloe vera Juice as Supplementary Nutrient on Larval, Cocoon and Silk Parameters in *Bombyx mori* L. *Biological Forum*, 17(9): 39-44.