

## Influence of Physical Characters of Different Grafted Scion on the Rootstock of Mango (*Mangifera indica* L.) under Shade Net Condition of Prayagraj Region

Vinay Joseph Silas<sup>1\*</sup>, V.M. Prasad<sup>2</sup>, Vijay Rajwade<sup>3</sup> and Saket Mishra<sup>4</sup>

<sup>1</sup>Ph.D. Scholar, Department of Horticulture, NAI, SHUATS, Prayagraj (Uttar Pradesh), India.

<sup>2</sup>Professor, Department of Horticulture, NAI, SHUATS, Prayagraj (Uttar Pradesh), India.

<sup>3</sup>Associate Professor, Department of Horticulture, NAI, SHUATS, Prayagraj (Uttar Pradesh), India.

<sup>4</sup>Assistant Professor, Department of Horticulture, NAI, SHUATS, Prayagraj (Uttar Pradesh), India.

(Corresponding author: Vinay Joseph Silas\*)

(Received: 01 January 2023; Revised: 13 February 2023; Accepted: 17 February 2023; Published: 22 February 2023)

(Published by Research Trend)

**ABSTRACT:** The experiment was conducted during the year 2019-20 & 2021-22, at Department of Horticulture, Naini Agricultural Institute, SHUATS, Prayagraj with the object to study the success and initial growth performance of grafts of some important varieties of mango under net house condition. The experiment consists of ten different varieties of mango Amrapali, Kishan Bhog, Gulabkhas, Kesar, Totapuri, Dasher, Malgoa, Himsagar, Nisar Pasand & Bombay Green as treatments and replicated thrice in Completely Randomized Design (CRD). The results of present investigation clearly showed that, grafts of different mango varieties studied had a significant influence on the maximum values of growth parameters like Graft Diameter (23.80 mm), scion diameter (17.35 mm), rootstock diameter (20.06 mm), length of grafted plant (88.71 cm), Days to sprout (14.25), Number of leaves (21.50), survival percentage at 150 DAG (90.00%) and rootstock length (65.72cm ) were recorded maximum in the graft of variety Dasher while Stionic ratio was found best in the graft of variety Gulabkhas (0.90), Scion length (25.14cm) was found best in the graft of variety Malgoa and maximum leaf area (307.43 cm<sup>2</sup>) was found in kesar.

**Keywords:** Mango, Rootstock, Scion, Grafting, Monoembryonic.

### INTRODUCTION

Mango is very well-adapted to tropical and subtropical climates and it thrives well in almost all regions from the sea level to an altitude of 1400 meter. The mango tree is hardy in nature and it can endure even the temperature as high as 48°C rainfall and high humidity during flowering and fruit development reduces fruit yield. It is grown widely, and it prefer a warm frost free climate with a well-drained (soil pH = 5.5-7.5), winter dry season and temperature ranging from 24-27 °C and the annual rainfall 400-3600mm (16-142 inch). It does not do well beyond a pH of 7.5.

Mango grafts are typically raised on unidentified seedling rootstocks, which causes variance among the grafts. To have uniform development, high yield, good quality fruits, and dwarf stature of plants for high density planting, it is crucial to standardise the rootstocks for various mango cultivars in various agroclimatic areas. This is conceivable if the rootstocks are grown asexually or using polyembryonic rootstocks, which are true types since they originated from nuclear tissues. It is also well known that rootstocks had a significant impact on most fruit crops' growth, subsequent fruit-bearing habits, and fruit quality. Therefore, it is necessary to choose the

appropriate rootstocks from the available varieties locally. Since these are used as rootstock for grafting and budding, raising of rootstocks and proper use of rootstocks is equally important.

For commercial plantations, mango is mostly grown vegetatively through side grafting and wedge grafting, which are more effective and affordable techniques than other forms of vegetative multiplication. Mango plants propagated via wedge grafting are ready for planting in around six months as opposed to a year with side grafting. The performance of a compound horticultural tree is determined by both rootstock and the scion (Bose *et al.*, 1991). The majority of mango trees are grafted trees that are typically cultivated in commercial orchards or in kitchen gardens. Each tree is made up of two parts: the rootstock, which provides the root system, and the scion, which forms the tree canopy. These two components are equally important to the survival of a tree. Grafting is a common and preferred vegetative propagation method for mango trees Bally (2006). Furthermore, proper alignment of scion and rootstock cambium tissues could determine the graft success Pina and Errea (2005).

Softwood grafting, a form of vegetative propagation that is effective, economical, quick, and can produce grafts in as little as a year, offers distinct advantages over other approaches. Therefore, softwood grafting results in better and more uniform orchard installations, initial success, and a reduced risk of death.

For grafted plants to survive, graft healing and acclimation during the grafting process are crucial. Therefore, the development of callus tissue followed by the union of vascular tissues is necessary for the successful union of the scion and rootstock. Successful grafting take and subsequent growth of scion shoot and development depend on a number of factors including scion variety, rootstock materials, and prevailing environmental conditions (Hartmann *et al.*, 2002).

Rootstocks, are always seedlings at their core. Monoembryonic non-descriptive seedlings are frequently utilised in India, in particular. Depending on where they are replicated and the environment, monoembryonic seedlings exhibit enormous variations in germination and vigour. Mango stones are only available in dry and semi-arid areas during the drier months of the year (April–June), which has a negative impact on the vigour and germination rates in these areas. According to reports, these regions have high graft-take during the month of September. The rootstocks raised during September months will not attain desirable girth and growth of grafting, thereby reducing the success of graft take and further survivability of grafted plants (Kumar *et al.*, 2017).

## MATERIALS AND METHODS

The investigation was conducted during 2019-2020 and 2021-2022 at the Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj.

**Materials.** The Scion was arranged from Central Institute of Subtropical Horticulture (CISH), Lucknow and the rootstock from Patel Nursery (in front of Chaka Block, Prayagraj). There was 10 varieties of Scion (each 30 in number) and will be grafted on the polyembryony rootstock of Mango.

**Selection of scion material.** The mother plant used were selected which was disease free and 6 months to 12 months old. The scion shoots of 25-30 cm length with 3 – 4 healthy buds were used for grafting. Selected scion shoots were defoliated on the mother plant about 5 – 7 days prior to detachment and at the same time, the apical growing portion of selected shoots were also beheaded which helped in forcing the dormant buds to swell.

**Raising of rootstock.** To raise the rootstock nursery, fully riped desi mango fruits/stones were collected from disease free, healthy, well managed and actively growing mango trees. One of the area's well-known model farmers provided the native rootstocks, and he managed them according to suggested cultural practises. Local types are typically favoured as rootstocks due to their

disease resistance and environmental adaptation. Stones were collected, cleaned with fresh water, and treated with 10 grammes of carbendazim in 10 litres of water. In the month of August, stones were sowed in elevated beds with a 10 cm 5 cm spacing, vertically. The stones germinated in two to three weeks, and following germination, the seedlings and stones were moved to the nursery beds after the leaves had turned brown and had grown to a quarter of their original size.

At the time of transplanting, the tap roots of these seedlings were pruned by retaining most of the fibrous roots. These transplanted seedlings were regularly watered and protected from the frost during winter season. Seedlings resumed growth in the end of February or beginning of March and became graftable from March onwards.

**Softwood grafting technique.** The top fresh growth (Bronze coloured shoots) developed on the rootstock was decapitated with sharp knife. After this, a longitudinal cut of 5 cm in length was given on the terminal trimmed shoot. The top of the rootstock appeared like the letter 'V'. The leaves below the 'V' cut were kept intact. A scion that was around the same thickness as the rootstock's severed shoot was chosen. The utilised scion was about 12 cm long. By shaving off the bark and a small amount of wood from the two opposing sides, the bottom end of the scion was transformed into a gently sloping wedge measuring about 5 cm. On the remaining two sides, care was taken to preserve the bark. A 1.5 cm wide by 45 cm long, 200 gauge thickness white translucent polythene tape was used to tightly fasten the wedge-shaped scion that had been manufactured in the manner described above. The scion on newly made graft were covered with small transparent polythene covers to avoid the desiccation of scions, by creating humidity near and above the graft union region.

**After care of the graft.** The polythene bags holding the grafted mango plants were kept in some shade. Regular watering of the plants was done with care made to ensure that water did not reach the graft union area. When sprouts began to emerge on the scion, the translucent polythene covers that had been placed on it after grafting were removed. When observed, sprouts that appeared on the rootstock beneath the graft union were routinely removed. Regular prophylactic pesticide spraying was also applied to control pest attacks.

## RESULTS AND DISCUSSION

The results of the investigation regarding the influence of scion varieties and rootstock on growth of mango have been presented in Table 1. The findings have been divided into the following sub-headings:

**Graft diameter.** The maximum graft diameter was recorded in Dasherri (23.80 mm) followed by Amarpali, Totapuri, Malgoa, Nisar Pasand and Himsagar, whereas the minimum graft diameter was found in Kesar (15.80

mm). Similar trend of increase in girth of plant was recorded by Mukherjee and Mujumdar (1964); Mane *et al.* (2020). The performance of scion depends upon stionic compatibility with rootstock and the perfectness of union Waghmare (1990).

**Scion diameter.** The maximum scion diameter was recorded in Dasherri (17.35 mm) followed by Amarpali, Gulabkhas, Totapuri, Malgoa, Himsagar and Nisar Pasand, whereas the minimum scion diameter was found in Kesar (11.98 mm.). These results are also similar with Nalage *et al.* (2010); Shivaram *et al.* (2018); Shinde *et al.* (2011). The increase in scion diameter was due to the main effect of grafting time was highly significant

**Rootstock Diameter.** The maximum rootstock diameter (20.06 mm) was recorded in Dasherri followed by Amarpali, Gulabkhas, Totapuri, Malgoa, Himsagar and Nisar Pasand, whereas the minimum scion diameter (14.00 mm) was found in Kesar. These results are also similar with Nalage *et al.* (2010); Shivaram *et al.* (2018).

**Stionic Ratio.** The maximum stionic ratio (0.90) was recorded in Gulabkash followed by Kishanbhog, Kesar, Nisar Pasand and Bombay Green. The maximum stock: scion ratio might be due to better compatibility of stock and scion, whereas the minimum stionic ratio (0.73) was found in Totapuri. The highest stock to scion ratio may be a result of stock and scion's improved compatibility. Better food material movement from leaves to other plant components is made possible by the stock-scion compatibility Bobade *et al.* (2018).

**Length of grafted plant.** At 150 DAG, the maximum total length grafted plant (88.71 cm) was recorded in Dasherri followed by Malgoa, Himsagar, Nisar Pasand and Gulabkhas. The maximum length of grafted plants of variety Dasherri might be due to higher length of tap root and more shoot length. whereas the minimum total length grafted plant (69.24 cm) was found in Totapuri. These results are also similar with Patel and Amin (1976); Praveena *et al.* (2018) in polyhouse, Sivudu *et al.* (2014) in shade net; Alam *et al.* (2000).

**Days to Sprout.** The days to sprout on scion of different varieties graft on the rootstock of mango were recorded

after wedge grafting. The maximum days to sprout (14.25) was recorded in Dasherri followed by Nisar Pasand, Totapuri, Bombay Green and Malgoa, whereas the minimum days to sprout (27.15) was found in Kishanbhog.

**Height of Sprouts.** The maximum height of sprout (14.88cm) was recorded in Dasherri followed by Malgoa, Amarpali, Nisar Pasand and Himsagar, whereas the minimum height of sprout (10.30 cm) was found in kesar. These results are in association with the result of Mulla *et al.* (2007) who recorded maximum length of sprout in the month of May with softwood grafting in mango

**Number of leaves.** The maximum number of leaves per grafted plant (21.50) was recorded in Dasherri followed by Gulabkhas, Nisar Pasand, Amarpali and Himsagar, whereas the minimum number of leaves per grafted plant (15.37) was found in kesar. The favourable weather during the grafting period led to increased cell activity and early sprouting, which are the causes of the increased number of leaves. Similar result was obtained by Panicker (1986) in Alphonso mango where highest number of leaves was produced in the month of May in softwood grafting.

**Survival % at 150 DAG.** The maximum survival percent of grafted plant at 150 DAG (90.00) was recorded in Dasherri followed by Totapuri, Amarpali, Nisar Pasand, Bombay Green and Malgoa, whereas the minimum survival percent of grafted plant at 150 DAG (20.00) was found in Kishanbhog. The maximum survival in variety Dasherri could be attributed to early healing of graft union, more number of leaves, and strong vegetative growth on account of better root development in the grafts of this variety as different root parameters like length of tap root, secondary roots in the grafts of this variety. Better root growth might have supplied required quantum of water and nutrients needed for strong vegetative growth of the grafts, which might have resulted in the higher survival percentage in grafts of this variety Ghojage *et al.* (2011).

**Table 1: Different physical characters of grafted scion on the Rootstock of Mango (*Mangifera indica* L.) under Shade Net condition of Prayagraj region.**

Varieties	Varieties name	Graft Diameter	Scion diameter	Rootstock Diameter	Stionic Ratio	Length of grafted plant	Days to Sprouts	Height of Sprout	Number of leaves	Survival % at 150 DAG	Scion Length	Rootstock length	Leaf area
V <sub>1</sub>	Amarpali	20.79	15.05	18.75	0.80	76.94	25.13	13.12	15.93	58.33	16.94	59.83	307.43
V <sub>2</sub>	Kishanbhog	17.04	13.02	15.16	0.86	69.24	27.15	11.10	14.01	31.67	19.01	50.85	267.06
V <sub>3</sub>	Gulabkhas	19.00	15.11	16.86	0.90	78.51	24.16	13.05	16.44	55.00	23.03	56.01	257.47
V <sub>4</sub>	Kesar	15.80	11.98	14.00	0.86	73.42	28.45	10.30	15.37	20.00	21.42	52.75	198.10
V <sub>5</sub>	Totapuri	18.03	14.00	19.31	0.73	77.26	20.14	11.32	16.80	61.67	18.94	58.36	265.77
V <sub>6</sub>	Dasherri	23.80	17.35	20.06	0.87	88.71	14.25	14.88	21.50	90.00	23.27	65.72	316.49
V <sub>7</sub>	Malgoa	20.83	13.90	18.37	0.76	86.25	21.72	11.07	15.88	63.33	25.14	61.19	289.59
V <sub>8</sub>	Himsagar	19.93	13.41	17.89	0.75	82.34	24.65	11.09	15.25	58.34	24.67	57.62	261.72
V <sub>9</sub>	NisarPasand	18.87	13.60	17.57	0.78	76.83	22.99	12.78	14.08	63.33	20.88	55.74	254.46
V <sub>10</sub>	Bombay Green	19.99	13.23	17.30	0.76	80.22	22.65	12.58	15.15	58.33	21.16	59.58	271.88
	F-Test	S	S	S	S	S	S	S	S	S	S	S	S
	C.D. 0.5%	0.16	0.32	0.44	0.02	1.51	2.59	0.52	1.40	28.34	0.41	1.19	19.46
	S.Ed (+)	0.07	0.15	0.21	0.01	0.71	1.23	0.25	0.66	13.49	0.19	1.63	9.26

**Scion Length.** The maximum scion length (25.14 cm) was recorded in Malgoa followed by Gulabkhas, Kesar, Himsagar and Bombay Green, where as the minimum scion length (16.94 cm) was found in Amrapali. These results are also similar with Patel and Amin (1976); Praveena *et al.* (2018) in polyhouse, Sivudu *et al.* (2014) in shade net.

**Rootstock length.** The maximum rootstock length (65.72 cm) was recorded in Dasherri followed by Amarpali, Gulabkhas, Malgoa and Totapuri, where as the minimum rootstock length (50.85 cm) was found in Kishabhog. These results are in association with the result of Mulla *et al.* (2007); Singh *et al.* (2014) who recorded maximum length of rootstock with softwood grafting.

**Leaf area.** Different lengths of scion shoots significantly affected leaf area per plant. The maximum length of tap root grafted plant (307.43 cm<sup>2</sup>) was recorded in Amrapali followed by Himsagar, Nisar Pasand, Bombay Green and Dasherri, whereas the minimum length of tap root grafted plant (198.10cm<sup>2</sup>) was found in kesar Waghmare (1990).

## CONCLUSIONS

The present investigation, can be concluded that the performance of grafts of Dasherri, Malgoa, Nisar Pasand & Amrapali varieties was found superior at nursery stage as grafts of these varieties have required comparatively well survival percentage. The growth parameters like graft diameter and shoot: root ratio were found maximum in these varieties. The root and biomass characters were also significantly superior in grafts of said varieties. Hence, it can be suggested that, for large scale multiplication of mango plants through wedge grafting, the scion wood of Dasherri, Malgoa, Amrapali & Nisar Pasand varieties may be used, as grafts of these varieties have recorded maximum success and higher values of growth parameters.

## FUTURE SCOPE

Grafting can reduce the incidence of plant disease, thus increasing potential yields. With reduced plant disease pressures, farmers can reduce their use of pesticides and may be more likely to be able to grow organically. Entrepreneurs can specialize in the production of grafted vegetable seedlings, earning income while selling improved seedlings to farmers.

**Acknowledgement.** The authors wish to thanks Department of Horticulture, Sam Higginbottom University of Agriculture, Technology and Science, Prayagraj for providing necessary facility in conducting the experiment.

**Conflict of Interest.** None.

## REFERENCES

Alam, M. N., Islam, M. S., Uddin, M. Z., Barman, J. C. and Quamruzzaman (2000). Effect of age of seedling and

variety of scion in stone grafting of mango. *Indian Journal of Sustainable Crop Production*, 1(2), 27-32.

- Bally, I. S. E. (2006). *Mangifera indica* (mango) Anacardiaceae (cashew family). Species Profiles for Pacific Island Agroforestry [www.traditionaltree.org](http://www.traditionaltree.org). Accessed on 20 April, 2009.
- Bobade, D. H., Ingole, R. H. and Kadam, A. S. (2018). Effect of Different Scion Varieties of Mango on Growth and Biomass Production performance of Stone Grafts (*Mangifera indica* L.). *International Journal of Current Microbiology and Applied Science*, (2018) *Special Issue-6*, 1642-1648.
- Bose, T. K., Mitra, S. K. and Sandhu, M. K. (1991). Propagation of tropical and subtropical horticultural crops. *Naya Prakash, Kolkata*, pp. 1-62.
- Ghojage, A. H., Swamy, G. S. K., Kanamadi, V. C., Jagdeesh, R. C., Kumar, P., Patil, C. P., & Reddy, B. S. (2011). Effect of season on softwood grafting in jamun (*Syzygium cumini* Skeels.). *Acta Horticulturae*, 890, 123-127.
- Hartman, H. T., Kester, D., Davies, F. T. and Geneve, R. L. (2002). *Plant Propagation Principles and Practices*, 7th Ed. Newfessy, Pricetice Hall, pp. 421-422.
- Kumar Ajay Karan., Varu, D. K., Archana, Puja Panda and Hota, Debashish (2017). Standardization of grafting time and height on success of softwood grafting in mango (*Mangifera indica* L.). *Journal of Pharmacognosy and Phytochemistry*, 6(5), 2803-2807.
- Mane, H. M., Dalvi, N. V., Khandekar, R. G., Salvi, V. G., Salvi, B. R., Bhingarde, R. T. and Bhagwat, S. R. (2020). Impact of Season on Softwood Grafting in Champaca (*Michelia champaca* L.) cv. Soundarya under Konkan Agroclimatic conditions. *International Journal of Current Microbiology and Applied Science*, (2020) *Special Issue-11*, 2021-2027.
- Mukherjee, S. K. and Majumdar, P. K. (1964). Effect of different factors on the success of veneer grafting in Mango (*Mangifera indica* L.). *Indian Journal of Horticulture*, 21, 4650.
- Mulla, B. R., Angadi, S. G., Mathad, J. C., Patil, V. S. and Mummigatti, U. V. (2007). Studies on softwood grafting in Jamun (*Syzygium cumini* Skeels.). *Karnataka Journal of Agricultural Science*, 24(3), 366-368.
- Nalage, N. A., Magar, S. D., Bhosale, S. S. and Mhetre, D. A. (2010). Effect of height of rootstock on success of epicotyl grafting in mango (*Mangifera indica* L.) cv. KESAR. *International Journal of Agricultural Science*, 6(1), Jan.-June, 2010.
- Patel, M. H. and Amin, R. S. (1976). Possibilities of bench grafting on young seedling of mango (*Mangifera indica* L.) under Anand condition. *Indian Journal of Horticulture*, 33(2), 156-161.
- Panicker, P. (1986). Studies on softwood grafting in Mango (*Mangifera indica* L.) var. Alphonso M.Sc. (Agri.) thesis submitted to D.B.S.K.K.V., Dapoli.
- Pina, A. and Errea, P. (2005). A review of new advances in mechanism of graft compatible-incompatible in Prunus spp. *Sci Hort.*, 106, 1-11.
- Praveena, Kumar R., Chandfre, Gowda M., Mounashree, S. and Vidya, A. (2018). Impact of season, growing condition and growth parameter of softwood grafting in

- Jamun (*Syzygium cumini* Skeel). *Journal of Pharmacognosy and Phytochemistry*, (1S), 1173-1176.
- Shinde, S. B., Saiyad, M. Y., Jadhav, R. G. and Chavda, J. C. (2011). Effect of time on softwood grafting success and survival of Jamun (*Syzygium cumini* Skeels) grafts. *International Journal of Agricultural, Science*, 7(1), 8-85.
- Shivaram, B., Virendra, S., Mahawer, L. N. and Arjun Lal Regar (2018). Effect of media and age of rootstock on epicotyl grafting of mango (*Mangifera indica* L.) cv. Amrapali. *International Journal of Chemical Studies*, 2018, 6(3), 3239-3243.
- Singh, Niranjana, Tripathy, S. M. and Vikas Ghumare (2014). Studies on growth and survival of stone grafts as influenced by age of seedling rootstock in mango (*Mangifera indica* L.) cv. Amrapali. *Journal of Applied and Natural Science*, 6(2), 716-719.
- Sivudu, B. V., Reddy, M. L. N., Baburatan, P. and Dorajeerao, Adda (2014). Effect of structural conditions on veneer grafting success and survival of mango grafts. *Plant Archives*, 2014, 14(1), 71-75.
- Waghmare, G. M. (1990). Studies on standardization of softwood grafting in Sapota (*Achras zapota* L.) var. Kalipatti. M.Sc. (Ag.) thesis submitted to D.B.S.K.K.V., Dapoli.

**How to cite this article:** Vinay Joseph Silas, V.M. Prasad, Vijay Rajwade and Saket Mishra (2023). Influence of Physical Characters of Different Grafted Scion on the Rootstock of Mango (*Mangifera indica* L.) under Shade Net Condition of Prayagraj Region. *Biological Forum – An International Journal*, 15(2): 961-964.