

17(4): 58-59(2025)

ISSN No. (Print): 0975-1130 ISSN No. (Online): 2249-3239

Medicinal Management of Urolithiasis in a Six-year-old Male Dachshund Dog

Jayaprasad A.^{1*}, Bante N.² and Sinha Y.K.³ ¹Teaching Associate, Department of Veterinary Medicine, Dau Shri Vasudev Chandrakar Kamdhenu University, Durg (Chhattisgarh), India. ²Teaching Associate, Teaching Veterinary Clinical Complex, Dau Shri Vasudev Chandrakar Kamdhenu University, Durg (Chhattisgarh), India. ³School of Wildlife Forensic and Health, Masters Student, Nanaji Deshmukh Veterinary Science University, Jabalpur (Madhya Pradesh), India.

(Corresponding author: Jayaprasad A.*) (Received: 29 January 2025; Revised: 07 March 2025; Accepted: 22 March 2025; Published online: 12 April 2025) (Published by Research Trend)

ABSTRACT: A 6-year-old male dachshund dog was presented to the Teaching veterinary clinical complex, durg, with a history of vomition, hematuria, anorexia and dysuria. All the physiological parameters were within range except serum creatinine and blood urea nitrogen. Abdominal ultrasonography showcased hyperechoic area within the bladder suggestive of cystolith and radiography finding revealed an opaque mass.

Keywords: Dachshund, Dysuria, Creatinine, Ultrasonography, Cystolith.

INTRODUCTION

Amongst canines, urolithiasis is a common problem that needs to be dealt with utmost care and urgency. Because of high recurrence rates, it's a point of concern. These stones may be made up of urates, phosphates, oxalates, silica and carbonates (Mulyani et al., 2024). Males are more prone than females for urolithiasis as the anatomical arrangement of penis fits suitably for urolith lodgment. Urolithiasis is highly lifethreatening condition that needs to be dealt with emergency (Formsa and Saini 2019). Certain infectious agents or trauma can predispose to urolith formation. According to Singh and Sailo (2013), certain factors leading to urolith formation are inorganic proteins, lack of crystal inhibitors along with crystallization in urine. Also age, breed, gender or any abnormality of the urinary tract needs to be considered (Rath et al., 2023). Difficulty in urination or strangurea along with hematuria are common clinical manifestations with stones in bladder or urinary tract (Hoxha and Rapti 2018). It is utmost important to have ultrasonography and X-ray findings to manage urolithiasis (Ucajev et al., 2024). There is severe bacterial infiltration seen in urolith suspected cases. Urinary tract diseases causes high morbidity amongst canines and felines and hence antimicrobial therapies are a must (Weese et al., 2019).

CASE HISTORY AND OBSERVATIONS

A 6-year-old male dachshund dog weighing 15 kg was brought to Teaching Veterinary Clinical Complex, Padmanabhpur, Durg with clinical signs of vomition, anorexia, hematuria and dysuria. Temperature, heart rate and respiration rates were within the normal range.

Jayaprasad et al.,

Biological Forum

Skin fold test examination revealed moderate dehydration and conjunctival mucous membrane was congested. Spike in serum biochemical parameters like creatinine (3 mg/dl) and blood urea nitrogen (65 mg/dl) were suggestive of urinary tract disorder.



Fig. 1. Hyperechoic area with acoustic shadows.



Fig. 2. Radiopaque mass. **17(4): 58-59(2025)**

Ultrasonography signified hyperechoic area along with acoustic shadows. In lateral plain radiograph, opaque mass was detected. The condition was diagnosed as urolithiasis.

Treatment. Treatment protocol included fluid therapy with Dextrose 5% and Ringers lactate through intravenous route to maintain the hydration status of the body. Antibiotic Amoxicillin and potassium clavulanate @ 12.5 mg/ kg P.O. twice daily along with urinary alkalizer disodium hydrogen citrate @ 7ml. P.O. once daily and allopurinol @ 20 mg/kg/day P.O. was administered for 15 days.

CONCLUSIONS

Urolithiasis is considered as one of the most common disorder in dachshunds. Their dissolutions are important to prevent any further complications leading to anuria. Proper antibiotic therapy for atleast 2 weeks in uroliths or associated urinary tract infections lead to good prognosis. Allopurinol is important to prevent ammonium crystal formation in the urine. Once the main triggering factor (stones) gets dissolved, blood urea nitrogen (40 mg/dl) and creatinine (1.2 mg/dl) values return to their normal values post treatment.

FUTURE SCOPE

For efficient management of reoccurrence of urolithiasis, surgical interventions are must.

REFERENCES

- Formsa, A. and Saini, N. S. (2019). Canine urolithiasis and concurrent urinary bladder abnormalities: symptoms, haematology, urinalysis and comparative radiographic and ultrasonography diagnosis. Veterinary Medicine Open Journal, 4(1), 18-26.
- Hoxha, Z. and Rapti, D. (2018). Breed predisposition differences of dogs with urolithiasis in tirana district. Anglisticum Journal (IJLLIS), 7(4), 1857-8187.
- Mulyani, G. T. Pramono, A. B. and Pangestiningsih, T. W. (2024). Diagnosis and Treatment of Urolithiasis in a Toy Poodle Dog. *Open Veterinary Journal*, *14*(3), 937-940.
- Rath, P., Sarangi, S., Mukherjee, P., Roy, S., Nandi, S. K. and Ghosh, D. (2023). Urolithiasis in Canines: A Comprehensive Review. *Journal of The Andaman Science Association*, 28(1), 1-12.
- Singh, K. B. and Sailo, S. (2013). Understanding Epidemiology and Etiologic Factors of Urolithiasis: An Overview. Science Vission, 13(4), 169-174.
- Ucajev, I., UCAJEV, B. and Pavlovic, I. (2024). Urolithiasis Vesicae Urinariae Canis, Case Report. Scientific Papers Journal Veterinary Series, 67(3), 127-129.
- Weese, J. S., Blondeau, J., Boothe, D., Guardabassi, L. G., Gumley, N., Papich, M., Jessen, L. R., Lappin, M., Rankin, S., Westropp, J. L. and Sykes, J. (2019). International Society for Companion Animal Infectious Diseases (ISCAID) Guidelines for the Diagnosis and Management of Bacterial Urinary Tract Infections in Dogs and Cats. *The Veterinary Journal*, 247, 8-25.

How to cite this article: Jayaprasad A., Bante N. and Sinha Y.K. (2025). Medicinal Management of Urolithiasis in a Six-yearold Male Dachshund Dog. *Biological Forum*, 17(4): 58-59.