

Studies on the Impact of Growing Transgenic Cotton on Soil Health in Major Bt cotton Growing areas of Tamil Nadu, India

T. Sherene¹, R. Kavimani² and Bharathi Kumar³

¹Assistant Professor, Agricultural College & Research Institute, TNAU, Trichy, (Tamil Nadu), INDIA

²Principal, Institute of Agriculture, TNAU, Kumulur, Trichy (Tamil Nadu), INDIA

³Assistant Professor, Cotton Research Station, TNAU, Veppanthattai, Perambalur (Tamil Nadu), INDIA

(Corresponding author: T. Sherene)

(Received 03 October 2018, Accepted 06 January, 2018)

(Published by Research Trend, Website: www.researchtrend.net)

ABSTRACT: There is a persistent environmental concern that transgenic Bt-crops have indirect undesirable effect to natural and agroecosystem function. We investigated the effect of Bt-cotton (with *Cry I Ac* gene) on soil biology in Bt cotton growing soils of Perambalur district, Tamil Nadu under rainfed scenario. Soil samples randomly from ten Bt cotton growing fields were selected in each of the taluks of Perambalur district of Tamil Nadu region, India, where Bt-cotton has been growing at least for ten continuous years and side by side non-Bt cotton grown soils were also collected to compare the extent of adverse effect of Bt toxin, if any. Samples were analyzed for various soil biological indicators like microbial population, microbial respiration, Microbial Biomass Carbon (MBC), Microbial Biomass Nitrogen (MBN), and soil Dehydrogenase (DHA) activities. The soil biological indicators like microbial population, soil respiration, DHA, MBC and MBN were found to be comparatively higher in Bt-grown soils than their non Bt counter parts over a period of 10 years.

Keywords: Soil health, Transgenic cotton, soil biological indices

How to cite this article: T. Sherene, R. Kavimani and Bharathi Kumar (2019). Studies on the impact of growing Transgenic cotton on Soil Health in major Bt cotton growing areas of Tamil Nadu, India. *Biological Forum – An International Journal*, 11(1): 18-23.

INTRODUCTION

There is a growing concern about cultivating transgenic cotton and its effects on general soil health. Most of the studies on impact of transgenic crops on soil properties carried out were restricted to contained conditions (Liu *et al.* 2005). Although some research has examined the environmental impacts of the 'aboveground' portion of transgenic crops, relatively fewer research effort has focused on the effects of these crops on soil microbes (Bruinsma *et al.* 2003) although no risk of growing transgenic Bt cotton on soil health is reported (Sun *et al.* 2007, Sarkar *et al.* 2009). Biological indicators of soil quality that are commonly measured include soil organic matter, respiration, microbial biomass (total bacteria and fungi,) and mineralizable nitrogen. The Bt-toxin has the potential to enter the soil system throughout the Bt-cotton-growing season, through root release and root turn over processes (Motavalli *et al.* 2004). While Bt occurs naturally in soil, growth of transgenic Bt-crop causes a large increase in the amount of Cry endotoxin present in agricultural systems, e.g. roughly 0.25 g ha⁻¹ produced naturally (calculated from approximately 1000 *Bacillus thuringiensis* spores g⁻¹ soil (Blackwood and Buyer 2004).

Genetically modified cotton genotypes incorporating a crystal (*Cry*) toxin producing *cryIAC* gene derived from *Bacillus thuringiensis* (Bt) were introduced in India for

commercial cultivation in the year 2002 (Morse *et al.* 2005). The transgenic crop, now popularly called Bt cotton, represents about 90% of cotton cultivated area in Tamil Nadu, India. In India, no comprehensive field study is available on the effects of growing transgenic cotton on soil biology. We evaluated the effects of growing transgenic Bt cottons and their counterpart (non-transgenic cotton) on selected soil biological attributes under rainfed conditions of Perambalur district in deep Vertisol.

MATERIALS AND METHODS

A. Soil sampling

Rhizosphere soil samples were collected 10 days before the harvest of crop at 30-45 cm depth from transgenic cotton growing fields of various taluks viz., Perambalur, Veppanthattai, Alathur and Veppur of Perambalur district and were labeled and transported back to the laboratory in polyethylene bags and stored at 4°C before analysis (Fig. 1). Soil sampling was also done in the non Bt cropped areas to assess the soil quality changes if any. As both cultivars of cotton were alike, except for the presence of the Bt-gene, it was assumed that any differences in soil ecological functions were attributable to the Bt-gene introduction in the cotton genotypes.

Normally, Bt cotton will be raised under rainfed conditions during the rainy season (October–December) with 90 × 45 cm spacing every year under rainfed scenario. Normal agronomic practices were followed for raising the crop.



Fig. 1. District Map of Perambalur, Tamil Nadu, India.

B. Soil biological indices

Soil microbial population. Samples (10 g fresh weight) were serially diluted in 90 mL Ringers solution up to 10^{-3} dilution and an aliquot of 1 mL of the aliquot was pour plated into selective media (nutrient agar for bacteria), Martin’s Rose Bengal Agar for fungi, Ken-Knight and Munaier’s Agar for actinomycetes and Buffered yeast agar for yeast. The plates were incubated at optimum temperature ($28 \pm 1^\circ\text{C}$ for bacteria and yeast; $30 \pm 1^\circ\text{C}$ for fungi and actinomycetes) in triplicates. The functional groups of microbes were enumerated by following standard microbiological methods (Wollum 1982). The microbial colonies appearing after the stipulated time period of incubation (3 days for bacteria and yeast; 5 days for fungi; 7 days for actinomycetes) were counted as colony forming units and expressed as cfu/g.

Soil respiration. Soil respiration was measured as the CO_2 evolved from moist soil, adjusted to 55% water holding capacity and pre-incubated for seven days at $22\text{--}25^\circ\text{C}$ with 10 mL of 1 mol/L NaOH. The CO_2 production was then measured by back titrating unreacted alkali in the NaOH traps with 1 mol/L HCl to determine $\text{CO}_2\text{-C}$ (Anderson 1982).

Soil microbial biomass carbon (MBC). Soil microbial biomass carbon was determined using the CHCl_3 fumigation-extraction method (Vance *et al.* 1987). Samples of moist soil (10 g) were used, and K_2SO_4 -extractable C was determined using dichromate digestion. Microbial biomass carbon was calculated using the equation:

$$\text{Biomass C} = 2.64 \text{ EC,}$$

Where, EC – (organic C in K_2SO_4 from fumigated soil) – (organic C in K_2SO_4 from non-fumigated soil).

Soil Microbial biomass Nitrogen (MBN). Soil microbial biomass nitrogen was estimated as $\text{MBN} = \text{EN}/0.54$ (Brookes *et al.*, 1985) where EN (Extractable Nitrogen) is the difference between N extracted from fumigated and non-fumigated samples

Dehydrogenase activity (DHA). Dehydrogenase activity (DHA) in soils was determined following the method of Casida *et al.* (1964) by the colorimetric measurement of reduction of 2, 3, 5-triphenyl tetrazolium chloride (TTC). Each soil sample (10 g) was treated with 0.1 g CaCO_3 and incubated for 24 h at 37°C . The triphenylformazan formed was extracted from the reaction mixture with methanol and assayed at 485 nm. FDA was measured following the method of Schnürer and Rosswall (1982) using 3, 6-diacetyl fluorescein as substrate and measuring the fluorescence at 490 nm.

C. Statistical analysis

Significant ($P < 0.01$ and $P < 0.05$) differences between Bt and non-Bt cotton on soil biological attributes were analyzed in the SAS programme (version 9.1). Tukey’s multiple comparison tests were done to determine the differences between Bt and non-Bt cotton crops.

RESULTS AND DISCUSSION

A. Impact of Bt cotton on soil microbial population

Bacterial and fungal population was significantly higher in Bt cotton grown soil compare with non-Bt soil at 0–15 cm depth. Soil bacterial population ranged from $30\text{--}58 \times 10^6$ CFU /g, Fungal population ranged from $14.3\text{--}16.5 \times 10^3$ CFU /g and actinomycetes ranged from $4.0\text{--}5.7 \times 10^3$ CFU /g in Bt cotton grown soils. Where as, in non Bt soils, bacterial, fungal and actinomycetes population were in the range of $25\text{--}33 \times 10^6$ CFU /g, $12.0\text{--}14.7 \times 10^3$ CFU /g and $2.8\text{--}3.8 \times 10^3$ CFU /g respectively. The increase in microbial population indicates no adverse effects of growing Bt cotton on soil microbial activity. The differences in the microbial population of Bt and non-Bt cotton hybrids may be attributed to variations in root exudates quantity, composition and root characteristics bring about by the genetic makeup of the cotton rather than expression of *cry* gene. Previous studies (Yan *et al.* 2007) have shown that the qualitative and quantitative differences in root exudation of Bt cotton could strongly influence the structure of microbial communities in the rhizosphere. Higher microbial populations in transgenic cotton grown soil were also reported by several workers (Shen *et al.* 2006, Kapur *et al.* 2010). Hu *et al.* (2009) based on their multiple-year cultivation showed that transgenic Bt cotton was not found to affect the rhizosphere functional bacterial population.

B. Impact of Bt cotton on soil respiration

The soil respiration was in the range of 224 -308µg of CO₂/ g / h in Bt cotton grown soils compared to non Bt cotton soils (168 -202µg of CO₂/ g / h) of various taluks of Perambalur district. Soil respiration rate was significantly (*P* < 0.01) highest in the Bt cotton grown soil followed by non-Bt grown soil. The increased soil respiration rate with Bt cotton in our study is the outcome of higher root volume in Bt cotton compare to non-Bt cotton that have stimulated the microbial growth and activity by enhanced resource availability.

C. Impact of Bt cotton on soil microbial biomass carbon.

Soils under Bt cotton hybrids had an average significantly (*P* < 0.01) higher amounts of MBC in the range of 175-191µg/g compared with the non-Bt 162 - 170 µg/g. The increased MBC in the soil grown with Bt cotton is attributed to higher root volume compared with non-Bt cotton. Possibly readily metabolizable carbon and nutrient availability at Bt cotton rhizosphere and differences in root exudates are perhaps the most influential factors contributing to increased microbial colonization and subsequent higher MBC in soils under Bt cotton. Earlier, Sarkar *et al.* (2009) reported a significant correlation between root volume of Bt cotton and soil MBC that supports the findings of Lynch and Panting (1980) that soil MBC increased with root growth and rooting density of the crop.

D. Impact of Bt cotton on soil microbial bio mass nitrogen.

The soil Microbial Biomass Nitrogen was in the range 0.43-1.48 per cent in Bt cotton grown soils whereas it was 0.073-0.092 per cent in non Bt counter parts. The increased MBN in the soil grown with Bt cotton is

attributed to higher root volume compared with non-Bt cotton. This might be due to comparatively higher root volume and associated biomass of Bt cotton that serve as a substrate for microbes to act and react with the soil when compared to its non Bt.

E. Impact of Bt cotton on soil dehydrogenase activities

Soil enzymes were suggested as one of the potential biological indicators of soil quality because of their relationship to soil biology, ease of measurement, and rapid response to changes in soil management. In our present study, the soils under Bt cotton had higher dehydrogenase activities (0.174 -0.228 µg TPF/g /h) than under non-Bt(0.068-0.079 µg TPF/ g / h) crop. DHA is considered as an indicator of the oxidative metabolism in soils and thus of the microbiological activity (Garcia *et al.* 1997) because it is linked to viable cells. Soil DHA reflects the total range of oxidative activity of soil microflora and, consequently it may be a good indicator of microbiological activity in the soil (Skujins 1976). Positive correlations between dehydrogenase activity and Bt cotton cultivation are also reported (Singh *et al.* 2013). DHA in soil depends on the content of soluble organic carbon (Zaman *et al.* 2002) and the increased organic matter in the surface soil horizon enhanced the soil enzyme activities. Studies by Furczak and Joniec (2007) showed that stimulation of DHA was accompanied by an increase in the number of the microbial groups and improvement in other living conditions (aeration and moisture). The low dehydrogenase activity indicates the low biological activity mainly due to the low soil organic carbon and the calcareous nature of the soil and poor soil fertility status in rainfed condition (James, 2002a, b; Benedict and Ring, 2004).

Table 1: Effect of Bt and non Bt cotton on soil microbial population in Perambalur district(Mean values of ten villages in eachtaluks).

Sl. No	Taluku	General microflora in Bt cotton grown soils (CFU/g)			General microflora in non Bt cotton grown soils (CFU/g)		
		Bacteria × 10 ⁶	Fungi × 10 ³	Actinomycetes × 10 ³	Bacteria × 10 ⁶	Fungi × 10 ³	Actinomycetes × 10 ³
1.	Veppanthattai	42	15.0	4.8	29	14.7	3.8
2.	Perambalur	58	14.3	4.0	33	13.8	2.8
3.	Alathur	30	14.8	5.2	25	12.0	2.9
4.	Veppur	35	16.5	5.7	25	14.3	3.1
	Range values	30-58	14.3-16.5	4.0-5.7	25-33	12.0-14.7	2.8-3.8
	SD	8.034	1.491	0.56	4.877	1.913	0.814

Table 2: Effect of Bt and non Bt cotton on soil microbial respiration and Dehydrogenase activity in soils of Perambalur district (Mean values of ten villages in eachtaluks).

S.No.	Taluku	Bt cotton grown soils		Non Bt cotton grown soils	
		DHA (µg TPF/g / h)	Soilrespiration µg of CO ₂ / g / h	DHA (µg TPF/g / h)	Soilrespiration µg of CO ₂ / g / h
1.	Veppanthattai	0.2137	224	0.071	164
2.	Perambalur	0.2281	264	0.068	181
3.	Alathur	0.1983	308	0.075	202
4.	Veppur	0.1739	286	0.079	201
	Rangevalues	0.174 -0.228	224-308	0.068-0.079	168-202
	SD	0.024	26.464	0.006	16.494

Table 3: Effect of Bt and non Bt cotton on soil Microbial Biomass Carbon (MBC) and Microbial Biomass Nitrogen (MBN) in soils of Perambalur district (Mean values of ten villages in each taluks).

S. No.	Taluku	Bt cotton grown soils		Non Bt cotton grown soils	
		MBC ($\mu\text{g/g}$)	MBN (%)	MBC ($\mu\text{g/g}$)	MBN (%)
1.	Veppanthattai	191	1.481	170	0.0813
2.	Perambalur	185	0.784	165	0.0732
3.	Alathur	175	0.427	162	0.0835
4.	Veppur	181	0.691	169	0.0918
	Rangevalues	175-191	0.43-1.48	162-170	0.073-0.092
	SD	4.671	0.310	3.273	0.007




Fig. 2. Dehydrogenase activity.




Fig. 3. Soil Respiration.




Fig. 4. Microbial Boimass Carbon.




Fig. 5. Microbial Biomass Nitrogen.

CONCLUSIONS

This study has demonstrated that cultivation of transgenic Bt cotton expressing *cryIAc* gene had no adverse effects on soil biological activities such as microbial population, soil respiration, dehydrogenase activity, microbial biomass carbon, and microbial biomass nitrogen. Based on the overall observations, growing Bt cotton was found to have a positive impact on soil biological activities. Our results suggest that cultivation of Bt cotton expressing *cryIAc* gene may

not pose ecological or environmental risk. Thus, the transgenic plants, either through the products of introduced genes and modified rhizosphere chemistry or through altered crop residue quality, have the potential to significantly change the essential ecosystem functions such as nutrient mineralization, carbon turnover and plant growth under long run. It needs continuous monitoring of Bt cotton grown soil environment for their biological indicators.

REFERENCES

- Anderson J.P.E. (1982). Soil respiration. In: Page A.L., Miller R.H., Keeney D.R. (eds.): *Methods of Soil Analysis. Part 2 – Chemical and Microbiological Properties*. 2nd Edition. ASA and SSSA, Madison, 837–871.
- Benedict, J.H. and D.R. Ring. (2004). Transgenic crops expressing Bt proteins: current status, challenges and outlook. In: Koul, O., Dhaliwal, G.S. (eds.), *Transgenic Crop Protection: Concepts and Strategies*. Science Publishers, Inc., Enfield, NH, USA, pp. 15–83.
- Blackwood, C.B. and J.S. Buyer. 2004. Soil microbial communities associated with Bt and non-Bt corn in three soils. *Journal of Environmental Quality*, **33**: 832–836.
- Brookes, P.C., A. Landham, G. Pruden and D.S. Jenkinson (1985). Chloroform fumigation and the release of soil nitrogen: A rapid direct extraction method to measure microbial biomass nitrogen in soil. *Soil Biology and Biochemistry*, **17**: 837–842.
- Bruinsma M., G.A. Kowalchuk, J.A. van Veen (2003). Effects of genetically modified plants on microbial communities and processes in soil. *Biology and Fertility of Soils*, **37**: 329–337.
- Brusetti L., P. Francia, C. Bertolini, A. Pagliuca, S. Borin, C. Sorlini, A. Abruzzese, G. Sacchi, C. Viti, L. Giovannetti, E. Giuntini, M. Bazzicalupo, D. Daffonchio. (2005). Bacterial communities associated with the rhizosphere of transgenic Bt 176 maize (*Zea mays*) and its non transgenic counterpart. *Plant and Soil*, **266**: 11–21.
- Casida L.E., D.A. Klein, T. Santoro (1964). Soil dehydrogenase activity. *Soil Science*, **98**: 371–376.
- Furczak J. and J. Joniec. (2007). Changes in biochemical activity of podzolic soil under willow culture in the second year of treatment with municipal-industrial sewage sludge. *International Agrophysics*, **21**: 145–152.
- Garcia C., T. Hernandez, F. Costa. (1997). Potential use of dehydrogenase activity as an index of microbial activity in degraded soils. *Communications in Soil Science and Plant Analysis*, **28**: 123–134.
- Hu H.Y., X.X. Liu., Z.W. Zhao, J.G. Sun, Q.W. Zhang Q.W., X.Z. Liu, Y. Yu. (2009). Effects of repeated cultivation of transgenic Bt cotton on functional bacterial populations in rhizosphere soil. *World Journal of Microbiology and Biotechnology*, **25**: 357–366.
- James, C. (2002). Preview: Global status of commercialized transgenic crops: 2002. ISAAA Briefs No. 27, Ithaca, NY. Retrieved from <http://www.isaaa.org/S>
- Kapur M., R. Bhatia, G. Pandey, J. Pandey, G. Paul D, R.K. Jain. (2010). A case study for assessment of microbial community dynamics in genetically modified Bt cotton crop fields. *Current Microbiology*, **61**: 118–124.
- Liu B., Q. Zeng, F. Yan, H. Xu, C. Xu. (2005). Effects of transgenic plants on soil microorganisms. *Plant and Soil*, **271**: 1–13.
- Lynch J.M., L.M. Panting. (1980). Cultivation and the soil biomass. *Soil Biology and Biochemistry*, **12**: 29–33.
- Morse S., R.M. Bennett, Y. Ismael. (2005). Genetically modified insect resistance in cotton: Some farm level economic impacts in India. *Crop Protection*, **24**: 433–440.
- Motavalli, P.P., R.J. Kremer, M. Fang and N.E. Means. (2004). Impact of genetically modified crops and their management on soil microbially-mediated plant nutrient transformations. *Journal of Environmental Quality*, **33**: 816–824.
- Sarkar B., A.K. Patra, T. Purakayastha, M. Megharaj (2009). Assessment of biological and biochemical indicators in soil under transgenic Bt and non-Bt cotton crop in a sub-tropical environment. *Environmental Monitoring and Assessment*, **156**: 595–604.
- Schnürer J., T. Rosswall. (1982). Fluorescein diacetate hydrolysis as a measure of total microbial activity in soil and litter. *Applied and Environmental Microbiology*, **43**: 1256–1261.
- Shen R.F., H. Cai, W.H. Gong (2006). Transgenic Bt cotton has no apparent effect on enzymatic activities or functional diversity of microbial communities in rhizosphere soil. *Plant and Soil*, **285**: 149–159.
- Singh R.J., I.P. Ahlawat., S. Singh. (2013). Effects of transgenic Bt cotton on soil fertility and biology under field conditions in subtropical inceptisol. *Environmental Monitoring and Assessment*, **185**: 485–495.
- Skujins J. (1976). Enzymes in soil. In: McLaren A.D., Peterson G.H. (eds.): *Soil Biochemistry*. Marcel Dekker, Inc., New York, 371–414.
- Sun C.X., L.J. Chen., Z.J. Wu, L.K. Zhou, Shimizu H. (2007). Soil persistence of *Bacillus thuringiensis*(Bt) toxin from transgenic Bt cotton tissues and its effect on soil enzyme activities. *Biology and Fertility of Soils*, **43**: 617–620.
- Vance E.D., P.C. Brookes., D.S. Jenkinson. (1987). An extraction method for measuring soil microbial biomass C. *Soil Biology and Biochemistry*, **19**: 703–707.
- Wollum A.G. (1982). Cultural methods for soil microorganisms. In: Page A.L., Mille R.H., Keeney D.R. (eds.): *Methods of Soil Analysis. Part 2 – Chemical and Microbiological Properties*. ASA and SSSA, Madison.
- Yan W.D., W.M. Shi, B.H. Li, and M. Zhang. (2007). Over expression of a foreign Bt gene in cotton affects the low-molecular-weight components in root exudates. *Pedosphere*, **17**: 324–330.
- Zaman M., K.C. Cameron, H.J. Di, and K. Inubushi. (2002). Changes in mineral N, microbial biomass and enzyme activities in different soil depths after surface applications of dairy shed effluent and chemical fertilizer. *Nutrient Cycling in Agroecosystems*, **63**: 275–290.