

Hematological analysis of Molluscan species *Bellamyia bengalensis* and *Lamellidens marginalis*

N.A. Kambale and V.V. Potdar

Department of Zoology, Shivaji University, Kolhapur, (MS) INDIA

ABSTRACT : Two molluscan species, freshwater snail *Bellamyia bengalensis* and a bivalve *Lamellidens marginalis* were used for the hematological analysis. Hemolymph of both the species was collected from the body parts by using nonlethal technique in in-situ condition. Cells observed in the hemolymph were compared morphologically and quantitatively to discuss its role in the mechanism of circulation, transportation and also by means of defense mechanism in the body of both species. Cellular comparisons were made at the species level.

Keywords : Mollusca, Haemocyte, *B. bengalensis*, *L. marginalis*

INTRODUCTION

In spite of the constant progress of studies on molluscan blood cells, there is no uniform and satisfactory classification system. Therefore these cells classified on one hand, the role of blood cells in various functional and development stages, and to the other hand, either by morphological aspects in the body (Adamowicz and Bolaczek, 2003). Hemolymph, the circulatory fluid of molluscs, which is slightly bluish in colour due to presence of respiratory pigment hemocyanine. Hemolymph transports nutrients, respiratory gases, enzymes, metabolic wastes and also toxicants throughout the body. Hemolymph with plasma and corpuscles can provide information pertinent to health assessment of animals or population. A large number of colorless stellate amoebocytes or corpuscles also referred as leucocytes are found in plasma, which are collectively called as hemocytes (Gustafson and Stoskop, 2005). Although hematology has been explored considerably and studies have been made on blood of molluscs, there is still much opportunity for investigation of blood differentiation and its mechanism in various body processes of freshwater snails and bivalves by means of their crucial role in body process. In general molluscs play a role in the balance of nature, and also act as biological indicator, in determining the degree of pollution of water and terrestrial environment. By considering the important role of hemocytes in molluscs, we have decided to study hematological analysis of two freshwater species as *Bellamyia bengalensis* and *Lamellidens marginalis*. Comparison of two species quantitatively as well as qualitatively was done at cellular level.

MATERIAL AND METHODS

For the present investigation two molluscan freshwater species (*Bellamyia bengalensis* and *Lamellidens marginalis*) were used. The first Freshwater prosobranch snail *Bellamyia bengalensis* belongs to class gastropoda, were collected from 'Rajaram tank', near the campus of Shivaji University Kolhapur, Maharashtra, India. The snails were collected and brought to laboratory in polythene bags. After the cleaning,

snails were kept in large plastic trough having 50 lits. The snails were provided with proper ventilation and food, like *Hydrilla*, *Pistia*, etc. After acclimatization for a week, healthy, adult and same sized snails (2.8-3.5 gm. weight, and 23-26 mm shell height) were sorted out and used for experimental studies.

Similarly the second molluscan freshwater bivalve species *Lamellidens marginalis* belongs to class pelecypoda inhabiting along the bank of Krishna river was specifically collected from south bank of Haripur and north of Ankali villages, Dist. Sangli, Maharashtra, India. After collection, the shells of these bivalve molluscs were cleaned to remove the fouling algal mass and mud and were kept for acclimatization for a week. The bivalves were provided with proper ventilation and food in the form of zooplanktons. The healthy, adult and same sized bivalves (9.5cm to 10.5 cm length in shell size) were used for experiment.

NONLETHAL METHOD OF HEMOLYMPH COLLECTION

Collection of hemolymph from the gastropod snail *Bellamyia bengalensis* was carried out in following manner. Prior to bleeding snail were washed with cold water in order to remove the faeces and excess mucus. It was then held in the left hand and then the syringe with a needle size 26 G½ (0.45mm × 13mm) was inserted in foot, through the operculum. The pressure is applied by withdrawing the plunger of the syringe and needle was slowly moved deeper until drops of pale blue hemolymph were aspirated. After the collection of required hemolymph the snail was returned to its enclosure.

In case of bivalve molluscs, *Lamellidens marginalis* hemolymph was collected by gently prying the shell to open approximately 5 to 7 mm. with a thin knife. The shell was held open with tissue forceps. The foot was visible between slightly gaping of shell valves, as a highly muscular white surface, then it was gently penetrated with a 26 G½ (0.45 mm × 13mm) sized needle. Required quantity of

1989), parasite invasion (Drozdowski and Zbikowska, 1994), injuries (Sminia, 1981), water content in the tissues (Zbikowska, 1998), and general condition of the organism (Barracco *et al.*, 1993). Older individuals may have a twice higher number of blood cells compared to young animals.

The study reveals that, the use of nonlethal technique for hemolymph collection, from foot is effective and allows the repeated bleeding in molluscs. At a time, as the welfare of invertebrates is increasingly attracting attention, it is very important that, procedures such as bleeding noninvasive as possible. According to Idakieva and Chakarska, (2009) hemolymph of molluscs can be collected by making diagonal slits on foot. Berg, (1995) and Naimo *et al.*, (1998), developed some relatively noninvasive technique for hemolymph collection from mantle and foot respectively. Similarly Gustafson and Stoskopf (2005) recorded collection of hemolymph from anterior adductor muscle of molluscs *Ellipto complanata* and found no adverse effect of repeated hemolymph sampling on growth and servivality of animal.

By considering functional aspects hemocytes can be distinguished as stem cells, phagocytes, trophic cells, haemostatically active cells which are responsible for blood hemostasis, and (Glinski and jarosz, 1997). On the basis of morphology, two basic cell types are recognized among molluscs hemocytes (Carballal, *et al.*, 1997). Agranulocytes (hyalinocytes) and granulocytes. (Adamowicz and Bolaczec, 2003). Some time morphology based classification is not enough to classify molluscan hemocytes, therefore several authors have used other criteria such as the enzymes content (Moore and Lowe 1977, Bayne *et al.*, 1979, Pipe 1990 a) antigenic characterization (Yoshino and Granath 1983, 1985, Dikkeboom, *et al.*, 1985, Morvan 1991, Noel *et al.* 1994) or lectin related properties (Pipe 1990 b). Hematological analysis in our study found that number of agranulocytes was more in *L. marginalis* than *B. bengalensis*. Where as number of granulocytes was more in *B. bengalensis* than *L. marginalis* which indicates that, hemoral defense mechanism was stronger in freshwater snail *B. bengalensis* as compared to bivalve *L. marginalis*.

ACKNOWLEDGEMENT

Authors are thankful to the U.G.C. New Delhi, for providing financial support under Major Research Project, Head, Department of Zoology, Shivaji University, Kolhapur. For their kind encouragement in the progress of work.

REFERENCES

Adamowicz, A. & Bolaczec. (2003). Blood cells morphology of the snail *Helix Aspersa Maxima* (Helicidae). *Zoologica Poloniae.*, **48**(1-4): 93-101.

Adema, C.M., Harris, R.A. & Van Deutekom-Mulder, E.C. (1992). A comparative study of hemocytes from six different snails: morphology and functional aspects. *J. Inv. Path.*, **59**: 24-32.

Barracco, M.A., Steil, A.A. & Gargioni, R. (1993). Morphological characterization of the hemocytes of the pulmonate snail *Biomphalaria tenagophila*. *Mem. Inst. Oswaldo Cruz, Rio de Janeiro.*, **88**(1): 73-83.

Berg, D.J., Haag, W.R., Guttman, S.I. & Sickel, J.B. (1995). Mantle biopsy: a technique for nondestructive tissue-sampling of freshwater mussels. *J.N. Am. Benthol Soc.*, **14**: 577-581.

Bayne, C.J., Moore, M.N., Carefoot, T.H. & Thompson, R.J. (1979). Hemolymph functions in *Mytilus Californianus*: the cytochemistry of hemocytes and their responses to foreign implants and hemolymph factors in phagocytosis. *J. Invertebr Pathol.*, **34**: 1-20.

Carballal, M.J., Carmen, L.M. & Carlos, A. (1997). Hemolymph cell types of the mussel *Mytilus galloprovincialis*. Diseases of aquatic organisms., **29**: 127-135.

Dikkeboom, R., Tijnagel, JMGH, Van der Knaap WPW. (1988) Monoclonal antibody recognized haemocyte subpopulation in juvenile and adult *Lymnaea stagnalis*: functional characteristics and lectin binding. *Dev Comp Immunol.*, **12**: 17-32.

Dikkeboom, R. Van der Knaap WPW, Maaskant, J.J. & De Jong, A.J. (1985). Different subpopulations of hemocytes in juvenile, adult and *Trichobilharzia ocellata*-infected *Lymnaea stagnalis*: a characterization using monoclonal antibodies *ZParasitenkd.*, **71**: 815-819.

Drozdowski, A. & Zbikowska, E. (1994). Wlyw pasozytnicznych larw przywr na aktywnosc immunologiczna slimakow. *Prz. Zool.*, **38**(1-2): 19-23.

Glinski, Z. & Jarosz, J. (1997). Układ immunologiczny mieczakow. In: Zjawiska odpornosci przeciwwzakaznej u bezkregowcow Wyd. UMC. Lublin., 90-100

Idakieva, K. & Chakarska, I. (2009). Purification of hemocytine from marine gastropod *Rapana Thomasiana* using ammonium sulphate precipitation method. *Biotechnol. & Biotechnol.*

Lori, L. Gustafson. & Michael K. Stoskopf. (2005). Evaluation of a nonlethal technique for hemolymph collection in *Ellipto Complanata*, a freshwater bivalve (Mollusca: Uniondale). *Disease of aquatic animals.*, **65**: 159-165.

Moore, M.N. & Lowe, D.M. (1977). The cytology and the cytochemistry of the haemocyte of *Mytilus edulis* and their responses to experimentally injected particals. *J. Invertebr pathol.*, **29**: 18-30.

Morvan, A. (1991). Caracterization antigenique des hemocytes de *Ihuitre Crassostrea gigas* et analyse de leur activite anti-infectieuse oxygene- dependante. Rapport de D.E.A., Universite de Bordeaux II.

Naimo, T.J., Damschen, E.D., Rada, R.G., & Monroe, E.M. (1998) Nonlethal evaluation of the physiological health of unionid mussels: methods for biopsy and glycogen analysis. *J. N. Am. Benthol Soc.*, **17**: 121-128.

Noda, S. & Loker, E.S. (1989). Effect of infection with *Echinostoma paraensei* on the circulating haemocyte population of the host snail *Biomphalaria glabrata*. *Parasitol.*, **98**: 35-41.

Noel, d. Pipe, R. Bachere, E. & Mialhe, E. (1994). Antigenic characterization of haemocyte subpopulations in the mussel *Mytilus edulis* by means of monoclonal antibodies. *Mar Biol.*, **119**: 549-556.

Ottaviani, E. (1989). Selected decreased of haemocytes of the freshwater snail *Planorbarius corneus* (L.) (Gastropoda, pulmonata) after bacteria injection. *Expientia*, **45**: 368-369.

Pipe, R.K. (1990a). Hydrolytic enzymes associated with the granular hemocytes of the marine mussel *Mytilus edulis*. *Histochem J.*, **22**: 595-603.

Pipe, R.K. (1990b). Differential binding of lectins to hemocytes of the mussel *Mytilus edulis*. *Cell tissue Res.*, **261**: 261-268.

Sminia, T., (1981). Gastropods. In: *Invertebrate blood Cells*. Ratcliffe N.A. and Rowley A.F. (eds.) Academic Press, London- New York, **2**: 191-232.

Yoshino, T.P. & Granath, W.O. JR. (1985). Characterization of molluscan phagocyte subpopulations based on lysosomal enzyme markers. *Exp. Zool.*, **226**: 205-210.

Zbikowska, E. (1998). Comparative quantitative studies of hemocytes of the snail: *Helix pomatia* L. and *Lymnaea stagnalis* (L.) (Gastropoda: Pulmonata). *Biol. Bull. Poznan.*, **35**(1): 25-32.