



A Note on Range Extension of *Mazus surculosus* D. Don from Indian Desert and Addition for Flora of Rajasthan, India

C.S. Purohit*, Amit Kumar and S.L. Meena

Botanical Survey of India, Arid Zone Regional Centre (Rajasthan), India.

(Corresponding author: C.S. Purohit*)

(Received 23 May 2025, Revised 19 July 2025, Accepted 08 August 2025)

(Published by Research Trend, Website: www.researchtrend.net)

ABSTRACT: *Mazus surculosus* D. Don is a small sized herb distributed mainly in North and North Eastern India. The present study deals with range extension of *Mazus surculosus* D. Don to Rajasthan from Sikar district and addition for flora of Rajasthan. Photographs, herbarium sheet, description and habitat of the species are discussed here for its easy identification.

Keywords: *Mazus surculosus*, addition, flora, Rajasthan, Indian desert.

INTRODUCTION

The Indian desert lies in the western India and includes a vast area of around 170,000 km² (66,000 sq.mi.), expanding from Gujarat and Rajasthan upto Punjab and Haryana. It is bounded on the west by the irrigated Indus River plain, on the north and northeast by the Punjab Plain, on the southeast by the Aravalli Range, and on the south by the Rann of Kachchh. This region receives a little rainfall and remains dry throughout the year. The average temperature in the Indian desert ranges between 75–70 degree Celsius in summers and 39–50 degrees Celsius in winters having warm days and cooler nights. This desert experiences a subtropical desert climate with high pressure and strong winds. In spite of harsh climatic and xerophytic conditions, this region harbours several plant, animal and birds species.

Several publications have been made dealing the floristic account of Indian desert, among which the two most pioneer works were carried out by Singh and Singh (2006) (Biodiversity of Desert National Park) and Bhandari (1990) (Flora of Indian Desert). The other comprehensive work on the Indian desert floral wealth was done by Shetty and Singh (1991) in the form of “Flora of Rajasthan” in three volumes. Some other publications on desertic plants include (Ramachandra Rao, 1941; Nair and Joshi 1957; Puri and Jain 1960, 1962; Gupta and Bhandari 1965; Meena, 2000; Purohit and Sharma 2006; Sharma & Purohit 2013; Kumar & Purohit 2015; Purohit, 2019; Purohit *et al.*, 2019; Purohit, 2020a, 2020b, 2020c, 2020d; Purohit *et al.*, 2020a, 2020b; Purohit, 2021, 2024, 2025; Indliya and Kumawat 2024; Purohit *et al.*, 2023, 2024, 2025; Ritu

et al., 2024; Purohit & Meena 2025) etc. have contributed to the flora of Indian desert of Rajasthan.

The genus *Mazus* was first described by Loureiro, Joao de in Flora cochinchinensis (1790). It contains 41 accepted species worldwide (POWO, 2025) and 05 species in India (Dash & Mao 2020). *Mazus surculosus* D. Don, belonging to family Scrophulariaceae, has been reported from Arunachal Pradesh, Himachal Pradesh, Jammu and Kashmir, Manipur, Punjab, Sikkim, Uttarakhand and West Bengal (Das & Mao 2020). This is the first time that this species has been found from Indian desert, Rajasthan.

METHODOLOGY

During the floristic survey of Indian desert, the authors noticed a population of *Mazus* species, different from *Mazus pumilus* (Burm.f.) Steenis. Voucher specimens were collected and preserved in BSJO. After detailed scrutiny of available literatures (Hooker, 1890; Blatter and Halberg 1921; Pandey *et al.*, 1983; Bhandari, 1990; Shetty & Singh 1991; Wu & Raven 1998; Grierson and Long 2001; Pusalkar & Singh 2012) and major herbaria (CAL, RRLH, M, BSJO, BSA, RUBL, JAC, BLAT, DCH), the specimen was identified as *Mazus surculosus* D. Don.

RESULT AND DISCUSSION

Mazus surculosus D. Don, Prodr. in Fl. Nepal. 87. 1825.

Herbs, upto 9 cm tall, stoloniferous, sparsely white villous. Stems short; stolons slender, internodes long. Basal leaves rosulate; hairy, lamina spatulate to

obovate, 2–7 cm including petiole, tapering at base, margin crenate, obtuse at apex. Cauline leaves smaller, opposite, petiolate; leaf blade obovate. Flowers bluish white, in terminal racemes, bracteole smaller than pedicel. Pedicel 15–18 mm long. Calyx tube campanulate, 10–12 mm; 5-lobed, lobes $\frac{1}{3}$ as long as calyx, ciliate, broadly ovate. Corolla bilabiate, twice the calyx; upper lip 2 lobed, 4–5 mm; lower lip 3 lobed, 7–10 mm long, spreading, larger than upper lip; middle lobe ovate and longer than lateral lobes. Stamen 4.

Capsule ovoid, ca. 4–6 mm long. Seeds small, many (Fig. 1).

Flowering & Fruiting: Jun-Oct.

Specimens Examined: Rajasthan, Sikar, Kaledi Berry, 27°45'33.38"N, 75°14'16.76"E, Alt.- 411m, 24.10.2023, C.S. Purohit & Amit Kumar 38248 (BSJO) [Fig. 2 & 3].

Distribution: Bhutan, China, Nepal and Pakistan. In India: Arunachal Pradesh, Himachal Pradesh, Jammu & Kashmir, Manipur, Punjab, Sikkim, Uttarakhand, West Bengal (Dash & Mao 2020) and Rajasthan (Sikar, present study) [Fig. 4].



Fig. 1. Shows plant *Mazus surculosus* D. Don.



Fig. 2. Herbarium sheet of *Mazus surculosus* D. Don.



Fig. 3. Habitat location of *Mazus surculosus* D.Don.




Fig. 4. Shows distribution of *Mazus surculosus* D. Don in India.

Habitat: The plant prefers marshy places or water logged regions along the road side and fields. In the present study the plant was located in open grassland near along the road in Kaledi Berry in Sikar district at an altitude of about 812.5 m along with some common occurring species like *Cenchrus purpureus* (Schumach.) Morrone (Poaceae; 38243); *Chloris gayana* Kunth. (Poaceae; 38246); *Clerodendrum phlomidis* L.f. (Lamiaceae; 38242); *Eleusine indica* (L.) Gaertn. (Poaceae; 38258); *Eragrostis amabilis* (L.) Wight & Arn. (Poaceae; 38249); *Erigeron bonariensis* L. (Asteraceae; 38250); *Euphorbia jodhpurensis* Blatt. & Hallb. (Euphorbiaceae; 38254); *Moringa oleifera* Lam. (Moringaceae; 38245); *Nyctanthes arbor-tristis* L. (Oleaceae; 38253); *Portulaca oleracea* L. (Portulacaceae; 38257); *Solanum nigrum* L. (Solanaceae; 38244), etc.

Conservation Status: *Mazus surculosus* D. Don has not yet been assessed on the IUCN criteria.

CONCLUSIONS

Mazus surculosus D. Don has been discussed here with its photo plates, keys for identification and habitat. This is the first instance that this species has been reported its extended distribution from the Indian desert and addition for flora of Rajasthan. Further exploration surveys are needed to disclose its more new locations in the arid regions. Also, IUCN assessment of this species is essential to understand its present conservation status.

Acknowledgement. The authors are thankful to Director, BSI for his continuous guidance and support. Thanks are also extended to Sh. Ranjeet Meghwal, Field assistant, BSI, AZRC, Jodhpur, Rajasthan for his assistance during the field work. Thanks are also due to

officials of Forest Department, Rajasthan for provide help during field survey.

Conflict of Interest. None.

REFERENCES

- Bhandari, M. M. (1990). Flora of the Indian Desert. *Scientific Publishers, (Revised Edition)*. Jodhpur.
- Blatter, E. & Hallberg, F. (1921). Species novae indiaorientales, *The Journal of Indian Botany*, 2(1 & 2), 48–49.
- Grierson, A. J. C. & Long, D. G. (2001). Flora of Bhutan, *Royal Botanic Gardens*, Edinburgh, 2, 1-1675.
- Gupta, R. K. & Bhandari, M. M. (1965). Flora of Rajasthan- A review. *Annales of Arid Zone*, 4(2), 236–238.
- Hooker, J. D. (1890). The Flora of British India. London. 2, 253.
- Indliya, P. and Kumawat, A. (2024). Response of Barley (*Hordium vulgare* L.) to Different Sources of Phosphorus in Arid Region of Rajasthan. *Biological Forum- An International Journal*, 16(4), 212-215.
- Dash, S. S. & Mao A. A. (2020). Flowering plants of India: An annotated checklist (Dicotyledons), *Botanical Survey of India*, 2, 219.
- Kumar, S. & Purohit, C. S. (2015). Conservation of Threatened Desert Plants. pp-148, Scientific publishers, Jodhpur.
- Meena, V. P. (2000). Desert Ecology, *Pointer Publisher*, Jaipur.
- Nair, N. C. & Joshi, M. C. (1957). Sand-dune vegetation of Pilani and its neighbourhood, *Journal of Indian Botanical Society*, 36, 599–617.
- Pandey, R. P., Shetty, B. V. & Malhotra, S. K. (1983). A preliminary census of Rare and Threatened plants of Rajasthan. In An assessment of Threatened plant of India (eds: S.K. Jain and R.R. Rao). *Botanical Survey of India, Howrah*.
- POWO (2025). <https://powo.science.kew.org/taxon/urn:lsid:ipni.org:names:37809-1> (assessed on June 25, 2025).
- Puri, G. S. & Jain, S. K. (1960). Trees and grasslands in Rajasthan. *Indian Forester*, 86, 85–86. 280.
- Puri, G. S. & Jain, S. K. (1962). Botanical exploration in Western India. *Tropical Ecology*, 3, 49–69.
- Purohit, C. S. (2019). A note on some rare plants of Rajasthan reported from Todgarh-Raoli wildlife sanctuary. *Int. J. Usuf. Mngt.*, 20, 36-45.
- Purohit, C. S. (2020a). Biodiversity of Todgarh-Raoli wildlife sanctuary with GIS mapping of EET species. Botanical Survey of India, Kolkata. 581pp. (*Project Report*).
- Purohit, C. S. (2020b). Ethno-medicinal and Economic Plants of Todgarh-Raoli wildlife sanctuary, Rajasthan – A Review. In Souvenir of International Webinar on “Global Environmental Challenges, Biodiversity, Principles of Guru Jambheshwar Ji and Remedies” organized by JNVU, Jodhpur and Jambhani Sahitya Academy, 3–5 June, 2020, Bikaner, pp. 54 – 66.
- Purohit, C. S. (2020c). *Dalechampia* & *Micrococoa*– Two generic addition for flora of Aravalli range, India with Status of Family-Euphorbiaceae of Todgarh-Raoli wildlife sanctuary, Rajasthan, India. *Journal of New Biological Reports*, 9(2), 209–219.
- Purohit, C. S. (2020d). Seven families addition under class Polypetalae for flora of Todgahr-Raoli wildlife sanctuary, Rajasthan. *Indian Journal of Arid Horticulture*, 1(2), 91-95.
- Purohit, C. S. (2021). *Striga todgarhica* (Orobanchaceae)- A new parasitic species from Todgarh-Raoli Wildlife Sanctuary, Rajasthan, India. *Journal of New Biological Reports*, 10(2), 89–94.
- Purohit C. S. (2024). A note on Family Asteraceae of Todgarh-Raoli Wildlife Sanctuary, Rajasthan, India. *Species*, 25, 1–26.
- Purohit, C. S. (2025). Reinstatement of the Name *Striga todgarhica* (Orobanchaceae) and a New Synonym of the Name *Striga angustifolia* var. *masuria*. *International Journal of Theoretical & Applied Sciences*, 17(2), 01–08.
- Purohit, C. S. & Meena, S. L. (2025). Reinstatement of the name *Alysicarpus monilifer* var. *venosa* (Fabaceae) and their IUCN Red List Assessment. *Species*, 26, e18s3101.
- Purohit, C. S., Maina, V. & Kumar, R. (2020a). GIS mapping of Endemic, Endangered and Threatened plants of Rajasthan. Botanical Survey of India, Kolkata 385pp. (*Project Report*).
- Purohit, C. S., Jain, K. & Merushikha (2020b). A note on family Lamiaceae in Todgarh-Raoli wildlife sanctuary, with addition of two species for flora of Rajasthan, *Int. J. Sci. Res. Sci. Technol.*, 7(2), 438-444.
- Purohit, C. S., Kulloli, R. N. & Bharti, A. (2019). *Euphorbia jodhpurensis* Blatt. & Hallb. an endemic plant from the Indian desert and its range extension from Todgarh-Raoli wildlife sanctuary, Rajasthan. *Euphorbia World*, 15(3), 18–22.
- Purohit, C. S. & Sharma, S. C. (2006). A note on Food and Fodder Grasses in North-west Rajasthan. *Geobios*, 34, 103–104.
- Purohit, C. S., Vivek, C. P., Dey, B. C. & Singh, L. J. (2023). *Hygrophila auriculata* (Acanthaceae): A new addition for Flora of Andaman & Nicobar Islands. *Journal of New Biological Reports*, 12(1), 5–8.

- Purohit C. S., Kumar A. & Meena S. L. (2024). Discovery of New Population of an Endemic species *Euphorbia jodhpurensis* Blatt. & Hallb. from Indian Desert, Rajasthan, India. *Journal of New Biological Reports*, 13(1), 9–12.
- Purohit, C. S., Dey, B. C., Ekka, G. A. & Singh, L. J. (2025). A Note on Occurrence of *Alpinia zerumbet* (Zingiberaceae) from Campbell Bay, Andaman and Nicobar Islands, India. *International Journal on Emerging Technologies*, 16(1), 54–56.
- Pusalkar, P. K. & Singh, D. K. (2012). Flora of Gangotri National Park, Western Himalaya, India. *Botanical Survey of India*, 487.
- Ramachandra Rao, Y. (1941). A list of some of the common plants of the desert areas of Sind, Baluchistan, Rajputana, Kathiawar and southwest Punjab, with some local names. *Indian Council of Agricultural Research, Bulletin No. 43*, 1–45.
- Ritu, Jain, K., Mourya, O. N. & Purohit, C. S. (2024). A note on occurrence of *Phaulopsis dorsiflora* (Retz.) Sant. (Acanthaceae) from Chhattisgarh state, India. *Journal of New Biological Reports*, 13(1), 15–18.
- Sharma, S. C. & Purohit, C. S. (2013). Grasses of North-west Rajasthan. Madhu publication, Bikaner pp-234.
- Shetty, B. V. & Singh, V. (1991). Flora of Rajasthan Vol. II. *Botanical Survey of India*, Howrah.
- Singh, V. & Singh, M. (2006). Biodiversity of Desert National Park, Rajasthan, *Botanical Survey of India*, Howrah.
- Wu, Z. & Raven, P. H. (1998). Flora of China, *Science Press* (Beijing) & *Missouri Botanical Garden Press* (St. Louis), 18, 1–449.

How to cite this article: C.S. Purohit, Amit Kumar and S.L. Meena (2025). A Note on Range Extension of *Mazus surculosus* D. Don from Indian Desert and Addition for Flora of Rajasthan, India. *International Journal of Theoretical & Applied Sciences*, 17(2): 105–109.