



## Reinstatement of the Name *Striga todgarhica* (Orobanchaceae) and a New Synonym of the Name *Striga angustifolia* var. *masuria*

C.S. Purohit\*

Botanical Survey of India, Arid Zone Regional Centre (Rajasthan), India.

(Corresponding author: C.S. Purohit\* chandansinghpurohit@yahoo.com)

(Received 05 March 2025, Revised 26 April 2025, Accepted 20 May 2025)

(Published by Research Trend, Website: [www.researchtrend.net](http://www.researchtrend.net))

DOI: <https://doi.org/10.65041/IJTAS.2025.17.2.1>

**ABSTRACT:** *Striga todgarhica* C.S. Purohit is reinstated in this paper. It used to be treated as a synonym of *Striga angustifolia* (D.Don) C.J. Saldanha but morphological characters of Calyx and Corolla support the reinstatement. These two species, together with *S. asiatica* make up a small group that is distributed in sandy-gravel desert area and have similar morphological characters. A key is provided to distinguish between them, and there is discussed and also recently describe species *S. khordhae* synonymized under *S. angustifolia* var. *masuria*.

**Keywords:** Reinstatement, *Striga todgarhica*, Endemic, New species, Synonym.

### INTRODUCTION

The genus *Striga* was first recognized and validly published by Loureiro 1790 belonging to the family Orobanchaceae. The genus comprises about 42 species with the highest diversity in Tropical Africa (Jayanthi *et al.*, 2014) where 28 taxa have been recorded of which 22 are endemic (Mohamed *et al.*, 2001, Fischer *et al.* 2011). The genus is represented by approximately 11 species and 3 infra-species (Jeeva *et al.*, 2012; Omalsree *et al.* 2015; Mao & Dash, 2020; Purohit 2021; Devi & Kumar, 2024). Six taxa so far reported from Rajasthan (Blatter & Hallberg, 1918; Bhandari, 1978; Shetty & Singh 1991; Kumar & Purohit, 2015; Purohit *et al.*, 2020; Purohit, 2020a & b, 2021; Purohit *et al.*, 2023; Purohit & Meena 2025).).

#### Reinstating *Striga todgarhica* C.S. Purohit

*Striga todgarhica* C.S. Purohit is endemic species of Rajasthan was originally described by C.S. Purohit from gravel area of Uperli Bhabhan, Pali, Rajasthan (Purohit, 2021) and housed in BSJO (Herbarium, Botanical Survey of India, Arid Zone Regional Centre, Jodhpur, Rajasthan). Due to type specimens available for examined, Arigela *et al.*, (2024) reduce it to a synonym of *Striga angustifolia* (D.Don) Saldanha based on climatic and ecological factors and without any herbarium collection from type locality or any other locality.

During the field survey of different parts of Aravalli hills, Indian desert, Mount Abu, Katchh, various sanctuary of Rajasthan and collected the plant with flower & fruit. After carefully examined the type specimens and newly collected specimens of both

species, author found that calyx and corolla to be significantly different. According to Hepper (1963) characters of diagnostic value in separating species include number of ribs in the calyx, indumentum, relative sizes of calyx and of corolla tubes, size of the bract subtending each flower, and flower color. Number of ribs in calyx is important identification characters in genus *Striga* and on the basis of this character classified all species under genus *Striga* (Berner *et al.*, 1997). Temperature is an important factor affecting the distribution of *Striga*, as prolonged exposure to high temperatures and humid conditions is required to break seed dormancy in *Striga* (Ejeta and Gressel, 2007).

The calyx of *S. todgarhica* have narrow triangular and 13 ribs (Fig. 1A to 1E) while the calyx of *S. angustifolia* have linear and 15 ribs (Fig. 2A to 2E). In *S. todgarhica* one calyx rib terminating the tip of each lobe and others in the sinus (Fig. 1D & 1E) while *S. angustifolia* all three calyx rib terminating the tip of each lobe (Fig. 2-4). Corolla tube of *S. todgarhica* is up to 2 cm long, tube apically strongly curved; Upper lip 2-partite and lower lip tri to tetra-partite (Fig. 1A & 1B) while the corolla tube of *S. angustifolia* is 0.8–1.5 cm long, tube abruptly incurved at or above the middle, upper lip 2-partite and lower lip outside in bud, tripartite (Fig. 3A & 3B; Fig. 4B & 4E). Study of herbarium sheets of *Striga angustifolia* in BSJO collected by various scientists of this office (Fig. 2 & 5) in different places and found same characters as observed above. Considering these significant differences, the name *Striga todgarhica* should be reinstated and characters of the species amended.

### Taxonomic treatment

**Striga todgarhica** C.S. Purohit in J. New Biol. Repor. 10(2): 89–94. 2021.

**Type:** INDIA. Rajasthan, Todgarh-Raoli Wildlife Sanctuary, Jojawar Range, Uperli Babhan, 460m, (25°46.306' N; 73°56.897' E), 12 November 2017, C.S.Purohit 33306A, (Holo BSJO!); C.S.Purohit 33306B (Iso BSJO!).

Annual, erect herb, 10–25 cm tall, entirely hirsute. Stem erect, branched ribbed, hispidly hairy. Leaves green, linear to narrowly lanceolate, with subacute apex, alternate, sessile, 2–4 × 0.2–0.4 cm, densely hairy on dorsal side, flowers in lax, leafy terminal spikes. Bracts similar to leaf in shape and size, densely hairy; Bracteole 2, equal, linear, densely hairy on both side, 6–8 mm long. Calyx 6–8 mm long; calyx lobes 5, narrow triangular, hispid, calyx lobe half the length of calyx tube [lobe 2–3 mm long; tube 5–6 mm long]; calyx 13-ribbed, densely hispid along ribs, one calyx rib terminating the tip of each lobe and others in the sinus. Corolla white, salverform, tube 1.5–2 cm, apically strongly curved, lower lip tri-tetra partite, outside in budding; upper lip bilobed, 2.2–2.6 × 2.2–2.4 mm. Stamen 4, attached to distal end of the tube, just below the throat. Ovary 1.5–2 mm long, glabrous; style long upto 4–5 mm long, glabrous, brown colour at the tip. Capsule ovoid, 8–10 mm long, enveloped in persistent calyx. (Fig. 1)

**Flowering & Fruiting:** October – December.

**Distribution:** Endemic, India, Rajasthan, Todgarh-Raoli wildlife sanctuary.

**Note:** This species is strictly parasitic on the grass roots of *Sorghum halepense* (L.) Pers. belonging to family Poaceae. It grows on transition area of gravel and sandy habitat in associated with *Aerva javanica* (Burm.f.) Juss. ex Schult., *Cordia sinensis* Lam., *Bergia ammannioides* Roxb. ex Roth., *Butea monosperma* (Lam.) Taub., *Calotropis procera* (Aiton) Dryand., *Commelina difusa* Burm.f., *Eragrostis amabilis* (L.) Wight & Arn., *Ipomoea nil* (L.) Roth, *Lindenbergia indica* Vatke, *Prosopis juliflora* (Sw.) DC., *Senna siamea* (Lam.) H.S.Irwin & Barneby, *Sorghum halepense* (L.) Pers. and *Trichosanthes tricuspidata* Lour.

### Key to the *Striga todgarhica* and morphologically similar congenis

- 1a. Calyx 5-ribbed 2
- 1b. Calyx 10-15 ribbed 3
- 2a. Leaves normal, linear, green *S. densiflora*
- 2b. Leaves reduced to scales, purple *S. gesnerioides*
- 3a. Three calyx-ribs terminating the tip of each lobe, Calyx 15-ribbed *S. angustifolia*
- 3b. One calyx-rib terminating the tip of each lobe and others in sinus, Calyx 10-13 ribs 4
- 4a. Leaves smooth or villous; Calyx 10-ribbed, hispidulous along ribs, calyx lobe linear, as long as calyx tube; petals lower lip tripartite, yellow, white or red; *S. asiatica*
- 4b. Leaves densely hairy on dorsal side; Calyx 13-ribbed, densely hispid along ribs, calyx lobe narrow triangular, half the length of calyx tube; petals lower lip tri – tetra partite, white; *S. todgarhica*

**Table 1: Distinguished characters between *Striga todgarhica*, *S. angustifolia* & *S. asiatica*.**

Characters	<i>S. angustifolia</i>	<i>S. todgarhica</i>	<i>S. asiatica</i>
Stem	Erect, sub-quadrangular, simple, rarely apically branched, hispidulous	Erect, sub-quadrangular, simple, branched from base, hispidulous	Erect, one rarely branched, ribbed, hispidly hairy
Leaf	Linear, smooth or slightly villous	Linear, densely hairy on dorsal side	Linear, smooth or slightly villous
Bracts	Similar to leaf in shape and size, slightly hairy	Similar to leaf in shape and size, densely hairy	Similar to leaf in shape and size, slightly hairy
Bracteoles	Less than 1 mm long, linear, slightly hairy	Less than 1 mm long, linear, densely hairy	Less than 1 mm long, linear, slightly hairy
Flower	Flowers sub-sessile, axillary, passing into terminal spike, 3 to numerous flowers	Flowers sub-sessile, in lax, leafy, terminal spike, 8–16 flowers	Flower axillary solitary or in a spike upwards, 3 – numerous flowers
Calyx	Calyx 9–10 mm; Calyx lobe: 5, calyx lobe as long as tube, subulate, linear	Calyx 6–8 mm; Calyx lobe: calyx lobe half the length of tube, narrow triangular, hispid	Calyx 4–8 mm; Calyx lobe: calyx lobe one third as long as tube linear
Calyx ribs	15-ribbed; three calyx ribs terminating the tip of each calyx lobe, hispidulous along ribs	13-ribbed, one calyx rib terminating the tip of each calyx lobe and others in the sinus, densely hispid along ribs	10-ribbed, one calyx rib terminating the tip of each calyx lobe and others in the sinus
Corolla	Corolla white; Corolla tube 0.8–1.5 cm long, tube abruptly incurved at or above the middle; upper lip 2-partite, lower lip outside in bud, tri-partite;	Corolla white; Corolla tube longer up to 2 cm, apically strongly curved. upper lip 2-partite, lower lip tri to tetra-partite.	Corolla bright yellow, white or red; Corolla tube 0.8–1.5 cm long, apically strongly curved; upper lip 2-partite, lower lip tri-partite;

### Similar species

*Striga asiatica* (L.) Kuntze – Linnaeus first described this species as *Buchnera asiatica* based on material from the Comoros Islands near Madagascar. Loureiro separated *Striga* from *Buchnera* in 1790 and applied the name *Striga lutea* to yellow-flowered specimens collected in India. Many other names were subsequently applied to different variants in different regions, including *Striga hirsuta* by Bentham in 1846. Saldanha (1963) expressed doubts whether Linnaeus' original name was valid and proposed that it should be abandoned in favour of *Striga lutea* Lour. However, Hepper (1974) confirmed that the original name was valid and the combination *Striga asiatica* (L.) Kuntze is now generally accepted. The species shows great variability in stature, branching, hairiness, flower colour and host preference (Fig. 6A to 6E), and different forms may occur in the same region. *Striga asiatica* has close relationship with *S. todgarhica* and *S. angustifolia*. The diagnostic characteristics of the three species are shown in Table 1.

### Specimen examined during study:

***Striga todgarhica* C.S. Purohit: Rajasthan:** Todgarh-Raoli Wildlife Sanctuary, Jojawar Range, Uperli Babhan, 25°46.306' N; 73°56.897' E, 460m, 12.11.2017, C.S. Purohit 33306A, 33306B;

***Striga asiatica* (L.) Kuntze: Gujarat:** Near nani Singhloti, Narmada, 08.09.2017, 20°36'00.4", 73°15'28.2", 125 m, Vinod Maina 35098; Near nani Singhloti, Narmada, 08.09.2016, 21°39'01.1", 73°37'49.1", 211 m, S.L. Meena 30600; Near Singhloti checkpost, Narmada, 29.08.2013, 21°39'54.2", 73°40'75.9", 277 m, S.L. Meena 27676;

***Striga angustifolia* (D.Don) Saldanha: Rajasthan:** Dabla, Jaisalmer, 02.10.2017, C.S. Purohit 33609; Aasan, Bijaguda, Todgarh- Raoli, 03.09.2018, 25°57.1'17", 74°07.9'38", 442 m, C.S. Purohit 33562; Plantation near hampi, Along Desuri-Phulad Road, Pali, 24.08.1975, 335 m, B.V. Shetty 1867; On the way to Rajmahal R.F. Deoli, Tonk, 11.09.1974, 330 m, B.V. Shetty 1215; Sunda hill, Rajpura, Jalore, 15.09.1979, 340 m, B.L. Vyas 6863; Near Bhadrajun along Jodhpur- Jalore road, Jalore, 23.09.1979, 250 m, B.L. Vyas 7345; Barli, Jodhpur, 07.10.1976, 375 m, A.N. Singh 3119; CAZRI afforestation area, Beriganga compt. I, Jodhpur, 04.10.1972, 165 m, B.V. Shetty 165; Bhaduwala, Ganganagar, 08.10.1977, 122 m, G.P. Roy 4951; Parah village, Bhilwara, 22.09.1978, A.N. Singh 6498; Stream in Naga R.F., Siwana Barmer, 17.10.1975, 385 m, B.V. Shetty 2233; Near Forest Route house, Kusalgargh, Banswara, 24.08.1976, 850 m, V. Singh 3031; Lasara village, Banswara, 14.10.1976, 600 m, V. Singh 3620; Bhuja ki Dhani, on the way to Dipas, Sikar, 20.09.1992, P.J. Parmar 10531. **Gujarat:** Umrat, Navsari, 31.10.2015, 21°02'79.8", 72°44'90.0", 2.3m., Vinod Miana 31759; Palsava-Village, Kachchh, 06.10.2004, 250 m., R.P. Pandey 19902; NSWS-near

Halapur, Kachchh, 14.10.2002, 60m., R.P. Pandey 13933; Taranga hills, Kheralu, Mehsana, 28.09.2003, P.J. Parmar 12934; Narayan sarovar, Haman khudi, Kutchh, 27.09.2000, V. Singh 15940; Jessoré wildlife sanctuary, Banaskantha, 07.09.2004, 290m., S.L. Meena 20796; Aaji dam, Rajkot, 23.08.2008, 150m., P.M. Padhye 23757; Nalsarovar Birds Sanctuary, Amdavad, 29.08.2005, 65m., S.L. Meena 21283; Pond near village sapar, Jamnagar, 15.02.2004, 50 ft., P.M. Padhye 18895; Dhowadri- Bhatiya, Jamnagar, 14.10.2003, 3000 ft., P.M. Padhye 18678; Near Shawlaji, Sabarkantha, 24.09.2005, P.J. Parmar 22784; Velavadar Blackbuck National Park, Bhavnagar, 12.08.2009, S.L. Meena 24362; Jessoré wildlife sanctuary, Banaskantha, 21.08.2003, 450m., S.L. Meena 18124; Goraj, Amdavad, 30.08.2005, 65m., S.L. Meena 21400; Umrat, Navsari, 31.10.2015, 21°02'79.8", 72°44'90.0", 2.3m., Vinod Maina 31750; Gichand forest, Narmada, 01.09.2013, 21°43'38.4", 73°40'95.2", 343m., S.L. Meena 27752; Junaraj Forest, Narmada, 02.09.2014, 21°46'24.1", 73°38'18.1", 430m., S.L. Meena 30274; Bansda national park, Navsari, 07.09.2017, 20°46.3'74.7", 73°29.1'12.1", 127m., Vinod Maina 35042.

**Synonymizing *Striga khordae* Sanjeet Kumar & Devi with *Striga angustifolia* (D.Don) Saldanha var. *masuria* (Buch.-Ham. ex Benth.) Omalsree & V.K. Sreenivas**

Described characters of a recently published new species *S. khordae* by Devi & Kumar (2024) i.e. smaller calyx, calyx lobes unequal to subequal, length one third as long as tube, each lobe of lower lip of corolla lobe is divided up to the half-length, longer capsule and seed surface irregularly pitted are developed in climate variation and all these characters reported in *S. angustifolia* var. *masuria* by Omalsree & Sreenivas (2018). According to Hepper (1963) characters of diagnostic value in separating species include number of ribs in the calyx and its clear-cut show the calyx having 15 ribs in *S. khordae* (photos of published paper) as well as *S. angustifolia* var. *masuria* and it revealed that *S. angustifolia* var. *masuria* shows characters i.e. pubescent nature of the plant, white corolla lobes, 15-ribbed calyx lobes, length of the plant, the loosely arranged flowers and closely positioned ribs on calyx lobes. Based on study of protoglosses and type specimens, author found that all the taxonomic characters of *S. khordae* (Devi & Kumar, 2024) are very similar and fall within the circumscription of *S. angustifolia* and treated as synonym of it.

### Taxonomic treatment

***Striga angustifolia* (D. Don) Saldanha var. *masuria* (Buch.-Ham. ex Benth.) Omalsree & V.K. Sreenivas, J.O.T.T. 10(9): 12294–12297. 2018.**

= *Striga masuria* (Buch.-Ham. ex Benth.) Benth. Compan. Bot. Mag. 1: 364. 1836.

= *Buchnera masuria* Buch.-Ham. ex Benth. in Scroph. Ind.: 41. 1835.

= *Buchnera wallichii* Benth. in N. Wallich, Numer. List: n.º 3876 (1831), nom. nud.

Lectotype: Nepal, Morang hills, 28.vii.1810, Francis (Buchanan) Hamilton, 1419 (E barcode: E00273651); Isolectotypes, K000899664, K001117506.


Syntypes: Myanmar, Prome, 09.x.1826, N. Wallich, 1161 (K K001117505, K000899663, E00273649).

= *Striga khordae* Sanjeet Kumar & Devi, J. Biodivers. Conservation 8(2): 23-27. 2024.

Holotype: India, Odisha, Khordha district, Godibari, 20°32' 25.71" N, 85°74' 39.92" E, c 48 m, 3 August 2020, S. Kumar & R.S. Devi 30 (Herbarium of Ambika Prasad Research Foundation).

Erect annual, chlorophyllous herbs, 15–68 cm tall. Stem densely hispid, quadrangular, green, branched from middle to apex or unbranched. Leaves opposite at base, alternate towards apex, sessile, 10–40 × 1–4 mm, linear-lanceolate, acute at apex, cuneate at base, hairy on both surfaces especially on mid-rib; margins entire


or ciliate with strigose hairs; mid-rib prominent. Inflorescence a raceme, terminal or from axils of upper leaves, 15–45 cm long. Rachis angular, strigose hairy. Flowers zygomorphic, sessile or subsessile, hypogynous, lax, alternate, 22–34 per inflorescence. Bract 1, 2.6–2.9 mm long, linear-lanceolate, hairy. Bracteoles 2, 1.8–2.1 mm long, linear-lanceolate, hairy. Calyx 6–8 mm, tubular, 15-ribbed; ribs distantly arranged and ending up to the teeth; lobes 5, 2–2.8 mm long, linear-lanceolate, hairy, green or with a brownish tinge. Corolla bilabiate; tube 8.8–9.8 mm long, prominently curved above the middle, greenish; lobes creamy-white, 3.8–4.4 mm long, broadly obovate, hairy; throat 4–4.2 mm long, hairy. Stamens 4, included, didynamous; filaments 1.5–3.2 mm long, attached to the distal end of the corolla tube; anthers 1-celled. Ovary superior, 2-celled, 2–2.2 mm long, oblong to ellipsoid, glabrous; ovules many, axile; style 4.8 mm long, white, brown at apex, glabrous; stigma brown. Fruit 5.5 mm long, ellipsoid, beaked. Seeds numerous, 0.3–0.4 mm long, ellipsoid, irregularly pitted, glabrous.




**Fig. 1.** *Striga todgarhica* C.S. Purohit: (A & B) – Whole plant with flowering; (C) – Calyx with nerves; (D) – Calyx ventral view showing 13 ribs; (E) Calyx dorsal view showing 13 ribs with one nerves reach on tip.




**Fig. 2. *Striga angustifolia* (D.Don) Saldanha:** (A) – Herbarium-collected from Katchchh, Gujarat; (B & C) – Calyx with ribs; (D & E) – Calyx close-up showing three ribs reach up to tip of calyx.




**Fig. 3. *Striga angustifolia* (D.Don) Saldanha** flower & bud shows three nerves in each calyx rich up to tip: (A & B) – Plant at Todgarh-Raoli wildlife sanctuary, Rajasthan; (C & D) – Plant at Mount Abu wildlife sanctuary, Rajasthan; (E) – Plant at Pali, Rajasthan.



**Fig. 4.** *Striga angustifolia* (D.Don) Saldanha flower & bud shows three nerves in each calyx rich up to tip: (A & B) – Plant at Desert National Park, Jaisalmer, Rajasthan; (C & D) – Plant at, Machiya Biological Park, Jodhpur, Rajasthan; (E & F) – Plant at Viratra Mata, Barmer, Rajasthan.



**Fig. 5.** *Striga angustifolia* (D.Don) Saldanha: (A) – Herbarium sheets, collected from Jalore, Rajasthan; (B) – Herbarium sheets, collected from Barmer, Rajasthan; (C) – Herbarium sheets, collected from Bhilwara, Rajasthan; (D) – Herbarium sheets, collected from Pali, Rajasthan; (E) – Herbarium sheets, collected from Tonk, Rajasthan.



**Fig. 6.** *Striga asiatica* (L.) Kuntze: (A) – Herbarium sheets, collected from Bansda National Park, Gujarat; (B & C) – Calyx close-up showing one ribs reach up to tip of calyx; (D) – Buds; (E) – Close-up of flower with leaf.

**Acknowledgements.** The Author's grateful to Director, Botanical Survey of India, Kolkata and Head of Office BSI, AZRC, Jodhpur for provide facilities during study and encouragement. Thanks are due to the officials of Forest Department of Rajasthan and authorities of Todgarh–Raoli wildlife sanctuary, Rajasthan for rendering the logistic support during field explorations. Thanks to all officials & staffs for help in plant collection and study of material.

## REFERENCES

Arigela, R. N., Singh, R. K., Kumar, S. & Reddy, C. S. (2024). Morphological variability, typification and a new synonym of *Striga angustifolia* (Orobanchaceae). *J. Biodivers. Conservation*, 8(4), 120–129.

Berner, D. K., Winslow, M. D. Awad, A. E., Cardwell, K. F., Mohan Raj, D. R. & Kim, S. K. (1997). *Striga* research methods – a manual. 2<sup>nd</sup> Edition. IITA Striga research group and International Institute of Tropical Agriculture, PMB, Ibadan, Nigeria.

Bhandari, M. M. (1978). *Flora of the Indian Desert*, Scientific publishers, pp- 107.

Blatter, E. & Hallberg, F. (1918). The Flora of the Indian Desert (Jodhpur and Jaisalmer). *J. Bomb. Nat. Hist. Soc.*, 26(1), 218-246.

Devi, R. S. & Kumar, S. (2024). A new species of *Striga* (Orobanchaceae) from Odisha state,

India. *Journal of Biodiversity and Conservation*, 8(2), 23–27.

Ejeta, G. & Gressel, J. (2007). Integrating New Technologies for *Striga* Control: Towards Ending the Witch-Hunt. Singapore; Hackensack, NJ: World Scientific.

Fischer, E., Lobin, W. & Mutke, J. (2011). *Striga barthlottii* (Orobanchaceae), a new parasitic species from Morocco. *Willdenowia*, 41, 51–56.

Hepper, F. N. (1963). Scrophulariaceae. In Flora of West Tropical Africa (2nd ed.) vol. 2 (eds. J. Hutchinson and J.M. Dalziel). London, UK: Crown Agents for Overseas Governments.

Hepper, F. N. (1974). Parasitic witchweed: *Striga asiatica* versus *S. lutea* (Scrophulariaceae). *Rhodora*, 76(805), 45–47.

Jayanthi, P., Prabhukumar, K. M., Rajendran, A., Thomas, B., Sabu, M. & Pradeep, A. K. (2014). *Striga indica* (Orobanchaceae) - A new parasitic species from Southern Western Ghats of India. *Feddes Reperorum*, 123, 283–290.

Jeeva, S., Shynin Brintha, T. S. & Rasingam, L. (2012). *Striga scottiana* (Scrophulariaceae) - a new species from southern Western Ghats of Tamilnadu, India. *Journal of Basic and Applied Biology*, 6(3&4), 79–82.

Kumar, S. & Purohit, C. S. (2015). Conservation of Threatened Desert Plants. pp-148, Scientific publishers, Jodhpur.

Mao, A. A. & Dash, S. S. (2020). Flowering plants of India: An annotated checklist (Dicotyledons), Botanical Survey of India 2: 219.

Mohamed, K. I., Musselman, L. J. & Riches, C.R. (2001). The genus *Striga* (Scrophulariaceae) in Africa. *Annals of the Missouri Botanic Garden*, 88, 60–103.

Omalsree, M., Pradhukumar, K. M., Sabu, M., Sunojkumar, P., Thomas, B., Rajendran, A. & Sreenivas, V. K. (2015). A new species of *Striga* section *Polypleurae* (Orobanchaceae) from southern western Ghats of India. *Phytotaxa*, 212(2), 163–168.

Omalsree, M. & Sreenivas, V. K. (2018). A reassessment and lectotypification of the name *Striga Masuria* (Buch.-Ham. ex Benth.) Benth. (Orobanchaceae) and its collection from the western ghats of India. *J.O.T.T.*, 10(9), 12294–12297.

Purohit, C. S. (2021). *Striga todgarhica* (Orobanchaceae)- A new parasitic species from Todgarh-Raoli Wildlife Sanctuary, Rajasthan, India. *J New Biol Rep.*, 10(2), 89–94.

Purohit, C. S. (2020a). Project report on Flora of Todgarh-Raoli wildlife sanctuary with GIS mapping of EET species. Botanical Survey of India, Kolkata. 581pp.

Purohit, C. S. (2020b). Dalechampia & Micrococoa—Two generic addition for flora of Aravalli range, India with Status of Family-Euphorbiaceae of Todgarh-Raoli wildlife sanctuary, Rajasthan, India. *J. New Biol Rep* 9(2), 209–219.

Purohit, C. S., Maina, V. & Kumar, R. (2020). Project report on GIS mapping of Endemic, Endangered and Threatened plants of Rajasthan. Botanical Survey of India, Kolkata. 385pp.

Purohit, C. S., Vivek, C. P., Dey, B. C. & Singh, L. J. (2023). *Hygrophila auriculata* (Acanthaceae): A new addition for Flora of Andaman & Nicobar Islands. *Journal of J. New Biological Reports*, 12(1), 5–8.

Purohit, C. S. & Meena, S. L. (2025). Reinstatement of the name *Alysicarpus monilifer* var. *venosa* (Fabaceae) and their IUCN Red List Assessment. *Species 2025*, 26, e18s3101.

Saldanha, C. J. (1963). The genus *Striga* Lour. *Bulletin of Botanical Survey of India* 5 (1), 67–70.

Shetty, B. V. & Singh, V. (1991). *Flora of Rajasthan* Vol-2, Botanical Survey of India, Howrah.

**How to cite this article:** C.S. Purohit (2025). Reinstatement of the Name *Striga todgarhica* (Orobanchaceae) and a New Synonym of the Name *Striga angustifolia* var. *Masuria*. *International Journal of Theoretical & Applied Sciences*, 17(2): 01–08.