



A little known orchid *Habenaria longicorniculata* (Orchidaceae) from Mount Abu wildlife sanctuary, Rajasthan and its IUCN status

C.S. Purohit¹, R.N. Kulloli^{2*}, V. Maina³, R. Kumar⁴, Balaji Kari⁵

¹Botanical Survey of India, Andaman & Nicobar Regional Centre, Port Blair, India- 744 102

²Arid Forest Research Institute, Jodhpur, Rajasthan, India-342 005

³Botanical Survey of India, Arid Zone Regional Centre, Jodhpur, Rajasthan, India-342 014

⁴Botanical Survey of India, CBL, Howrah, West Bengal, India-711 103

⁵Mount Abu wildlife sanctuary, Forest Department, Sirohi, Rajasthan

Corresponding author: r.kulloli@gmail.com

| Received: 31 August 2020 | Accepted: 15 October 2020 |

How to cite: Purohit CS, Kulloli RN, Maina V, Kumar R, Kari B. 2020. A little known orchid *Habenaria longicorniculata* (Orchidaceae) from Mount Abu wildlife sanctuary, Rajasthan and its IUCN status. J New Biol Rep 9(3): 299 – 305.

ABSTRACT

During the survey of Mount Abu wildlife sanctuary, Sirohi, Rajasthan, a little known rare terrestrial orchid species *Habenaria longicorniculata* was observed from four locations in this sanctuary. Sharma (2003) reported this species from Phulwari wildlife sanctuary, Udaipur. Mount Abu wildlife sanctuary is newly reported location of this species and also addition for flora of Sirohi district. Based on our observations and assessment using IUCN criteria, this species places under the category Vulnerable regionally [criteria – Vu/B₁b (i, v) D₁ (IUCN version 3.1)]. The identification characters, distribution map along with live macro and microscopic photographs are provided in the present study. The current status of *Habenaria longicorniculata* in Rajasthan with the help of IUCN criteria sheets is also provided.

Key words: *Habenaria longicorniculata*, IUCN, status, terrestrial orchid.

INTRODUCTION

Habenaria Willdenow (1805: 5) is one of the largest genera of the family Orchidaceae having c. 883 terrestrial species (Govaerts et al. 2018). *Habenaria* has wide distribution throughout the tropical and subtropical regions of the old and new world (Pridgeon et al. 2001; Batista et al. 2013), with centers of diversity in Brazil, southern and central Africa and East Asia (Batista et al. 2013). In India, it is represented by 84 species; 39 of them are found in the Western Ghats, with 22 being endemic (Misra 2007; Nayar et al. 2014, Kumar et al. 2016), while, in

Rajasthan 4 species occurs (Choudhary et al. 2011; Sharma 2003; Shetty and Singh 1991; Kulloli and Purohit 2020).

Habenaria longicorniculata J. Graham was first collected by John Graham from Khandala (Graham 1839) and then near Belgaum (Dalzell and Gibson 1861). As per the literature distribution of the species is reported as from Konkan to Travancore (Hooker 1890), Malabar and Konkan (Cooke 1907), Khandala and Deccan (Santapau and Kapadia, 1966). Further it is reported from other districts of

Maharashtra i.e. Akola, Aurangabad, Amravati, Pune, Raigad, Satara, Kolhapur, Ratnagiri and Sindhudurg (Lakshminarasimhan 1996; Almeida 2009; Shivkar and Shinde 2015). From Gujarat state this species was reported from three localities including Dang district (Suryanarayan 1968; Shah and Suryanarayana 1969; Shah 1978; Tadvi 2013), Jessore Wildlife Sanctuary of Banaskantha district (Desai 2013) and recently, from Sabarkantha (Punjani et al. 2019). In Rajasthan, state it was earlier reported from single locality from Phulwari Wildlife Sanctuary, Udaipur (Sharma 2003). While working on the GIS mapping of threatened plants of Rajasthan during 2017 to 2019, authors were conducted field survey of Mount Abu wildlife sanctuary during August, 2019 (Fig.1) and collected unidentified herbarium samples of family Orchidaceae. These collected herbarium samples deposited at Botanical Survey of India, Jodhpur (BSJO). After critical study of literature (Shetty and Singh 1991; Sharma 2003; Choudhary et al. 2011; Nayar et al. 2014, Kumar et al. 2016) and herbaria (BSJO, BSA, RUBL, JAC, BLAT, DCH, CAL, K), it was identified as *Habenaria longicorniculata* J. Graham. It also noticed that earlier this species was not reported from Sirohi district. So this species is addition the flora of Sirohi district.

Habenaria longicorniculata J. Graham, Cat. Bombay Pl. 202. 1839; Santapau and Kapadia, Orch. Bombay 29, t.1, f.1, 1966; *Habenaria longecalcarata* A. Rich in Ann. Sci. nat. Ser. II 15:71. T 3B. 1841; Wt. Icon t. 925. *Habenaria longecalcarata* A. Rich var. *viridis* Blatt. & Mc Cann in Journ. Bombay nat. Hist. Soc. 36:20, 1932, Shah FLG 658. 1978.

Morphological Description: Terrestrial herb, up to 1m high, subglobose tubers with several slender roots. Leaves clustered at the base of the short stem with amplexicaul sheaths. Inflorescence racemose, few flowered. Flowers pedicellate, green in colour and fragrant. Dorsal sepal green, 3-nerved, ovate, concave. Lateral sepal opened backwards, longer than the dorsal sepal, white with greenish tinge on the margins. Petal greenish, forms hood along with dorsal sepal, somewhat ligulate; Anther pollinia 2, caudicle longer than the pollinia, viscidium inconspicuous. Rostellum

little shorter than the anther. Stigmatic processes 2, dark green (darkest in the flower), ovary ribbed. Spur 10-15 cm long, whitish at base becomes darker green at the apex, slightly pointed at apex, bulges at the opening. Labellum white, side lobes slightly larger than the midlobe. Sidelobes deflexed outwards (Fig.2).

Fl. & Fr.: August – September.

Distribution: India: Maharashtra, Karnataka, Andhra Pradesh, Tamilnadu, Goa, Chhattisgarh, Jharkhand, West Bengal, Gujarat (Sabarkantha, Banaskantha, Dang), Rajasthan (Sirohi and Udaipur).

Habitat and Ecology: This species has its typical habitats in rocky and gravelly area and undergrowth in forest areas, often on moist slopes in core zone of Mount Abu wildlife sanctuary (Fig.3), associated with *Carvia callosa* (Nees) Bremek., *Flueggea leucopyrus* Willd., *Lantana camara* L., *Euphorbia nerifolia* L., *Chlorophytum tuberosum* (Roxb.) Baker, *Senna insularis* (Britton & Rose) H.S. Irwin & Ba and *Carissa carandas* L. It prefers high altitudes between 900 – 1200 m.s.l.

Threats: Major threat to decline of this species is consumption of tubers by wild animals like bear and wild boar as food. The remaining population is perpetuating by means of the tubers, having edible and medicinal properties which are collected by local people. Habitat destruction and developmental activities for tourism purpose is also threat to decline the population.

Specimen examined: INDIA: RAJASTHAN: Mount Abu wildlife sanctuary, Adhar Devi, 26 Sept. 2018, C.S. Purohit & R. Kumar 36928 (BSJO).

Status: Based on our observations area of occupancy <500 km² (1 km²) and population size < 1,000 with mature individuals (500-650 individuals at four locations) *Habenaria longicorniculata* should be placed in the Vulnerable threat category of IUCN regionally. This needs to be further confirmation.

Detailed survey of areas of occurrence confirmed its rarity. Hence its assessment using IUCN criteria placed it under the category Vulnerable regionally [**criteria- Vu/ B₁b (i, v); D₁ (IUCN version 3.1)**]. Therefore it needs immediate efforts to rehabilitate this species in nature.

SHEET OF IUCN CRITERIA-A

Use of the criteria A	Critically Endangered	Endangered	Vulnerable
A. Population reduction	Declines measured over the longer of 10 years or 3 generations		
A1	≥ 90%	≥ 70%	≥ 50% X
A2, A3 & A4	≥ 80%	≥ 50%	≥ 30% X

A1. Population reduction observed, estimated, inferred, or suspected in the past where the causes of the reduction are clearly reversible **AND** understood **AND** have ceased, based on and specifying any of the following

	Tick right sign.	others
(a) direct observation	X	
(b) an index of abundance appropriate to the taxon	X	
(c) a decline in area of occupancy (AOO), extent of occurrence (EOO) and/or	X	

(iv) number of locations or subpopulations	X	
(v) number of mature individuals	√	

B1 OR B2. (c) Extreme fluctuations in any of		
	Tick right sign.	others
(i) extent of occurrence	X	
(ii) area of occupancy	X	
(iii) number of locations or subpopulations	X	
(iv) number of mature individuals	X	

IUCN Assessment	=	Vu/ B₁b(i, v)
------------------------	---	---------------------------------

SHEET OF IUCN CRITERIA-C

Use of the criteria C	Critically Endangered	Endangered	Vulnerable
C. Small population size and decline			
Number of mature individuals	< 250	< 2,500	< 10,000 X
C1. An estimated continuing decline of at least:	25% in 3 years or 1 generation	20% in 5 years or 2 generations	10% in 10 years or 3 generations
C2. A continuing decline	(up to a max. of 100 years in future)		
C2. (a) (i) Number of mature individuals in each subpopulation	< 50	< 250	< 1,000 X
C2. (a) (ii) % individuals in one subpopulation =	90–100%	95–100%	100% X
C2. (b) Extreme fluctuations in the number of mature individuals.			

IUCN Assessment	=	Vu/ B₁b(i, v)
------------------------	---	---------------------------------

SHEET OF IUCN CRITERIA-D

Use of the criteria D	Critically Endangered	Endangered	Vulnerable
D. Very small or restricted population			
Number of mature individuals	< 50	< 250	D1 < 1,000 √
VU D2. Restricted area of occupancy or number of locations with a plausible future threat that could drive the taxon to CR or EX in a very short time			D2. typically: AOO<20 km ² or number of locations ≤ 5

IUCN Assessment	=	Vu/ B₁b(i, v); D₁
------------------------	---	--

SHEET OF IUCN CRITERIA-E

Use of the criteria E	Critically Endangered	Endangered	Vulnerable
E. Quantitative Analysis			
Indicating the probability of extinction in the wild to be	≥ 50% in 10 years or 3 generations (100 years max.)	≥ 20% in 20 years or 5 generations (100 years max.)	≥ 10% in 100 years X

IUCN Assessment	Vu/ B₁b(i, v); D₁
------------------------	--




Fig.1 Distribution map of *H. longicorniculata*.



Fig. 2 A. Habitat and B. Habit of *H. longicorniculata* at Mount Abu wildlife sanctuary, Rajasthan.




Fig.3. *Habenaria longicorniculata* J. Graham. (A) Habit; (B) Bud; (C) Bract; (D) Flower front view; (E) Flower side view; (F) Lip; (G) Dorsal sepal; (H) Lateral sepals; (I) Lateral petals; (J) Gynoecium; (K) Close up of stigmatic lobes; (L) Pollinaria and (M) Fruit- capsule.

REFERENCES

- Abraham A, Vatsala P. 1981. An introduction to orchids with Illustrations and description of 150 South Indian Orchids. Tropical Bot. Garden and Research Institute, Trivandrum.
- Almeida MR. 2009. Flora of Maharashtra VA: 59. Orient Press, Mumbai.
- Batista JA, Karina SB, Marina WF, de Faria K, Aline P, Gerardo JR. 2013. Molecular phylogenetics and morphological reappraisal of the Platanthera clade (Orchidaceae: Orchidinae) prompts expansion of the generic limits of Galearis and Platanthera. *Annals of Bot* 104: 431-445.
- Batista JAN, Borges KS, de Faria MWF, Proite K, Ramalho AJ, Salazar GA, Van den Berg C. 2013. Molecular phylogenetics of the species-rich genus *Habenaria* (Orchidaceae) in the New World based on nuclear and plastid DNA sequences. *Molr Phylogene Evol* 67: 95–109.
- Bhatt MR, Nagar PS. 2014. *Habenaria foliosa* A. Rich. (Orchidaceae) – A new record for the flora of Saurashtra, Gujarat. *J Econ Taxon Bot*38(3-4): 552 – 554.
- Bhatt RP, Sabnis SD. 1987. Contribution to the Ethnobotany of Khedbrahma region of North Gujarat. *J Econ TaxonBot*9: 139–145.
- Bhatt RP, Bedi SJ. 1969. A study of the vegetation and flora of Khedbrahma region of North Gujarat. BSI, India J J: 3 J 1–321.
- Cerejo-Shivkar S, Shinde RD. 2015. A brief account of Orchidaceae in Sanjay Gandhi National Park, Mumbai, India. *J Threat Taxa* 7(6): 7287–7295.
- Choudhary C, Mukharjee SK, Chowdhary HJ. 2011. Distribution and Diversity of the Genus

- Habenaria* Willdenow (Orchidaceae) in India. In: Chandra Ghosh & A.P. Das (Eds.), Recent Studies in Biodiversity and Traditional Knowledge in India. Publisher- Gour Mahavidyalaya, Malda.
- Cooke T. 1907. The Flora of the Presidency of Bombay III: 223. Botanical Survey of India.
- Dalzell NA, Gibson A. 1861. The Bombay Flora. Education Society's Press, Calcutta.
- Desai PR. 2013. Studies on Floristic Diversity, Ethnobotany and Resources Survey in Jessore Wildlife Sanctuary of Banaskantha District, Gujarat, Ph.D. thesis, HNG University, Patan, Gujarat.
- Deva S, Naithani HB. 1986. Orchid Fl. North West Himalayas. Print and Media Associates, NewDelhi, 1–459. Government Press, Bombay.
- Graham J. 1839. A catalogue of the plants growing in Bombay and its vicinity pp 202.
- Hooker JD. 1890. The Flora of British India VI: 141. L. Reeve & Co., London.
- Jain SK, Rao RR. 1977. Field and Herbarium Methods. Today and Tomorrow's Printers and Publishers. New Delhi, India.
- Kulloli RN, Purohit CS. 2020. *Habenaria gibsonii* var. *foetida* (Orchidaceae): An addition to the flora of Rajasthan. Species 21(67): 202 – 207.
- Kumar CS, Shetty BV, Bennet SSR, Rao TA, Molur S, Walker S. (eds.), 2001. Endemic Orchids of the Western Ghats - Conservation Assessment and Management Plan (C.A.M.P.) Workshop. Wildlife Information Liaison Development (WILD) Society and Zoo Outreach Organisation Coimbatore, India.
- Kumar P, Jalal JS, Rawat GS. 2007. Orchidaceae, Chotanagpur, state of Jharkhand, India. Check List 3(4): 297–304.
- Kumar P, Prabhukumar KM, Nirmersh TK, Sreekumar VB, Hareesh VS, Balachandran I. 2016. *Habenaria sahyadrica* (Orchidaceae, Orchideae) a new species from the Western Ghats (India) with critical notes on allied taxa. Phytotaxa 244 (2): 196–200.
- Kumar P, Gale SW, Pedersen HE, Phaxaysomba T, Bouamanivang S, Fischer GA. 2018. Addition to the Orchid Flora of Laos and taxonomic notes on orchids of the Indo-Burma region. Taiwania 63(1): 61 – 83.
- Lakshminarasimhan P. in Sharma, 1996. Flora of Maharashtra State Monocotyledons. Botanical Survey of India, Calcutta.
- Misra S. 2004. Orchids of Orissa. Bishen Singh Mahendra Pal Singh, Dehra Dun.
- Misra S. 2007. Orchids of India - A Glimpse. Bishen Singh Mahendra Pal Singh, Dehradun, v+402 pp.
- Nayar TS, Sibi M, Beegam AR. 2014. Flowering plants of the Western Ghats, India. Palode, India: Jawaharlal Nehru Tropical Botanic Garden and Research Institute.
- Pandey VB. 2011. Plant Species Diversity and their Ethnobotanical study in Tribal area of Bhiloda (West) Forest range of Sabarkantha district (North Gujarat). Ph.D. thesis HNG University, Patan, Gujarat.
- Parmar PJ. 2012. A checklist of the Vascular Plants of Sabarkantha District, Gujarat, India. Nelumbo 54: 92-137.
- Patel SK. 2003. A contribution to the Flora of Meghraj and Malpur talukas of Sabarkantha District of North Gujarat. Ph.D. thesis, HNG University, Patan, Gujarat.
- Pridgeon AM, Cribb PJ, Chase MC, Rasmussen FN. 2001. Genera *Orchidacearum*, Vol. 2, Orchidoideae (part 1). Oxford: Oxford University Press.
- Punjani B, Patel S, Desai P, Chaudhary Y, Pandey V. 2019. *Habenaria longicorniculata* J. Graham (Orchidaceae), a new distributional record from Sabarkantha district for Gujarat state. Bioscience Discovery 10(2):53–57.
- Punjani BL. 1997. An Ethnobotanical Study of Tribal Areas of District Sabarkantha (Gujarat). Ph.D. thesis, HNG University, Patan, Gujarat.
- Santapau H. 1955. Botanical collectors manual. Min. of Natural Resources and Science Research New Delhi, pp 62.
- Santapau H, Kapadia Z. 1966. The Orchids of Bombay. Govt. of India Press, Calcutta, vi+ 239pp.
- Saxton WT. 1922. Additional notes on plants of Northern Gujarat. Rec. Bot. Survey of India, 9: 251- 262. <http://biosciencediscovery.com> 57 ISSN: 2231-024X (Online) Bioscience Discovery, 10(2): 53 – 57.
- Saxton WT, Sedgwick LJ. 1918. Plants of Northern Gujarat. Bot. Survey of India 6 (7): 209– 323.
- Shah GL. 1978. Flora of Gujarat State, Part I and II. Sardar Patel University Press, Vallabh Vidyanagar.
- Shah GL, Suryanarayana B. 1969. Further contribution to the flora of Dangs forest in Gujarat. Bulletin of Botanical Survey of India 11 (3 & 4): 290–300.
- Sharma SK. 2003. Flora of protected areas-I. Orchid Flora of Phulwari Wildlife Sanctuary Udaipur district, Rajasthan. Zoo Print J, 18(10):1227– 1228.
- Shetty BV, Singh V. 1991. Flora of Rajasthan Vol-2. Botanical Survey of India, Howrah.
- Suryanarayana B. 1968. A contribution to the flora of Dangs forest, Gujarat, Part I, II, and III. Ph.D. thesis, Sardar Patel University, Vallabh Vidyanagar, Gujarat.
- Tadvi DS. 2013. Floristic diversity of Dangs. Ph.D. thesis, M.S. University, Baroda, Gujarat.
- Yogi DV. 1970. A contribution to the flora of North Gujarat. Ph. D. thesis, SP University Vallabh Vidhyanagar, Gujarat.